

of transitional shelters to environmental tobacco control interventions.

A telephone survey of all long-term transitional shelters in Los Angeles County was conducted from March to June 2005. The 20 min institutional survey utilised a pretested 31-item questionnaire to query administrative officials at each participating shelter about resident demographics, institutional characteristics (eg, size, average length of stay, etc), inventory of existing "no smoking" policies and the willingness of each facility to participate in environmental tobacco control efforts. The study inclusion criteria included that (1) each facility must be operational at the time of the survey; (2) each facility must offer at least a 12-month length of stay to their homeless residents to qualify as a "long-term" transitional shelter; and (3) each facility must be a transitional housing programme based on the continuum of care model for rehabilitating homeless individuals back to mainstream society.⁹

Bivariate and multivariate analyses were conducted using the STATA 9.0 statistical software package to evaluate the relationships between shelter receptivity to environmental tobacco control interventions (dependent variable) and institutional-resident characteristics (independent variables). To understand whether tobacco control policies varied by size of facility, subgroup analyses were conducted by facility-size categories based on the maximum number of beds that were available at each facility: small (<50 beds), medium (50–199 beds) or large (>200 beds). To estimate individual-level statistics from these facility-specific data, some sociodemographic measures were weighted by the number of beds reported for the facility; other statistics were computed directly from shelter-level data.

In all, 76 transitional shelters met the study inclusion criteria and were contacted (table 1). Of these, 71 (93.4%) participated in the survey. Nearly a quarter (23%) had a programme that focused on helping individuals who were addicted to drugs or to alcohol to reduce their dependence on these substances. Another third (30%) had programmes that focused on helping mothers with small children transition to more permanent housing; eight (11%) were dedicated to housing the mentally ill; three (4%) focused on helping war veterans, emancipated youth or gay men find more permanent housing. A majority of the transitional shelters reported having an indoor "no smoking" policy (75%) and designated smoking areas (78%). A total of, 72% and 95%, respectively, reported that they would be open to adopting new anti-smoking policies and environmental tobacco

control interventions designed to reduce resident smoking.

Results from comparative analyses suggest that the size of the facility was inversely related to the likelihood that the facility would report having a current indoor no smoking policy ($p<0.001$), a designated no smoking area ($p<0.001$) and a policy to limit smoking among its residents ($p<0.001$) (table 1). The size of the facility and the service mission of the organisation (eg, those with substance abuse treatment programmes) also predicted greater receptivity towards environmental tobacco control interventions ($p<0.001$). These findings suggest that enforcement of a smoke-free environment may be more difficult at larger facilities or in facilities without experience in treatment for substance misuse.

Contrary to the prevailing perception among tobacco control advocates that homeless service providers have casual attitudes towards client smoking behaviour,³ the present data showed that homeless service facilities, such as long-term transitional housing, recognise the importance of intervening on this high-risk behaviour; they are highly receptive to tobacco control efforts designed to reduce the prevalence of smoking in this population. This emerging viewpoint in the homeless community would suggest that greater investment of tobacco-control resources, tailored to the unique needs of the homeless, would be justified and deserving of further investigation.^{3 10}

Lisa Arangua

Department of Family Medicine, David Geffen School of Medicine at University of California, Los Angeles, California, USA

William J McCarthy

Department of Health Services, University of California, Los Angeles School of Public Health, Los Angeles, California, USA

Rebecca Moskowitz

University of California, Los Angeles School of Law, Los Angeles, California, USA

Lillian Gelberg

Department of Family Medicine, David Geffen School of Medicine at University of California Los Angeles, Los Angeles, California, USA

Tony Kuo

County of Los Angeles, Department of Public Health, Department of Family Medicine, David Geffen School of Medicine at University of California Los Angeles, Los Angeles, California, USA

Correspondence to: Dr T Kuo, 10880 Wilshire Blvd, Suite 1800, Los Angeles, California 90024-4142, USA; tkuo@mednet.ucla.edu

doi: 10.1136/tc.2006.018697

Competing interests: None declared.

References

- 1 US Department of Health and Human Services. (USDHHS). *The health consequences of smoking: a report of the surgeon-general*. Washington, DC: US Printing Office, 2004.
- 2 Okuyemi KS, Caldwell AR, Thomas JL, et al. Homelessness and smoking cessation: insights from focus groups. *Nicotine Tob Res* 2006;**8**:287–96.
- 3 Apollonio DE, Malone RE. Marketing to the marginalized: tobacco industry targeting of the homeless and mentally ill. *Tob Control* 2005;**14**:409–15.
- 4 Arnsten JH, Reid K, Bierer M, et al. Smoking behavior and interest in quitting among homeless smokers. *Addict Behav* 2004;**29**:1155–61.
- 5 Butler J, Okuyemi KS, Jean S, et al. Smoking characteristics of a homeless population. *Subst Abuse* 2002;**23**:223–31.
- 6 Connor SE, Cook RL, Herbert MI, et al. Smoking cessation in a homeless population: there is a will, but is there a way? *J Gen Intern Med* 2002;**17**:369–72.
- 7 Greene JM, Ennett ST, Ringwalt CL. Substance use among runaway and homeless youth in three national samples. *Am J Public Health* 1997;**87**:229–35.
- 8 McCarthy WJ, Collins C, Hser YI. Does cigarette smoking affect drug abuse treatment? *J Drug Issues* 2002;**32**:61–79.
- 9 Robertson MJ. The prevalence of mental disorder among homeless people. In: Jahiel, R, eds. *Homelessness: a prevention-oriented approach*. Baltimore and London: Johns Hopkins University Press, 1992:57–86.
- 10 Rohrbach LA, Howard-Pitney B, Unger JB, et al. Independent evaluation of the California Tobacco Control Program: relationships between program exposure and outcomes, 1996–1998. *Am J Pub Health* 2002;**92**:975–83.

CORRECTION

In the February issue of *Tobacco Control* the cover credit was incorrect. Please find the correct credit below.

Cover credit: Miss America Chesterfield's Valentine Girl Patricia Donnelly of Detroit. A Pocketful of Pleasure Chesterfield. The real reason why Chesterfields are in more pockets every day is because Chesterfield's Right Combination of the world's best cigarette tobaccos gives you a better smoke. . .definitely milder, cooler and better-tasting. You can't buy a better cigarette. Make your next pack Chesterfield. They Satisfy.