- Siegel M. Involuntary smoking in the restaurant workplace. A review of employee exposure and health effects. JAMA 1993;270:490—3.
- Stillman F, Navas-Acien A, Ma J, et al. Second-hand tobacco smoke in public places in urban and rural China. Tob Control 2007;16:229—34.
- Semple S, Maccalman L, Naji AA, et al. Bar workers' exposure to secondhand smoke: the effect of Scottish smoke-free legislation on occupational exposure. Ann Occup Hyg 2007;51:571—80.
- Pilkington PA, Gray S, Gilmore AB. Health impacts of exposure to second hand smoke (SHS) amongst a highly exposed workforce: survey of London casino workers. BMC Public Health 2007;7:257.
- Gee IL, Watson AF, Carrington J, et al. Second-hand smoke levels in UK pubs and bars: do the English Public Health White Paper proposals go far enough? J Public Health (Oxf) 2006;28:17—23.
- Ho SY, Lam TH, Chung SF, et al. Cross-sectional and prospective associations between passive smoking and respiratory symptoms at the workplace. Ann Epidemiol 2007;17:126—31.
- Burney PG, Laitinen LA, Perdrizet S, et al. Validity and repeatability of the IUATLD (1984) Bronchial Symptoms Questionnaire: an international comparison. Eur Respir J 1989;2:940-5.
- Repace J, Hughes E, Benowitz N. Exposure to second-hand smoke air pollution assessed from bar patrons' urinary cotinine. Nicotine Tob Res 2006:8:701—11.

- Wakefield M, Cameron M, Inglis G, et al. Secondhand smoke exposure and respiratory symptoms among casino, club, and office workers in Victoria, Australia. J Occup Environ Med 2005;47:698

 —703.
- Huss A, Kooijman C, Breuer M, et al. Fine particulate matter measurements in Swiss restaurants, cafes and bars: What is the effect of spatial separation between smoking and non-smoking areas? Indoor Air 2010;20:52—60.
- World Health Organization. Report on the Global Tobacco Epidemic, 2009: implementing smoke-free environments. Geneva, Switzerland: WHO, 2009. http://www.who.int/tobacco/.
- Zheng P, Fu H, Li G. Smoke-free restaurants in Shanghai: should it be mandatory and is it acceptable? Health Policy 2009;89:216—24.
- Jaakkola MS, Jaakkola JJ, Becklake MR, et al. Effect of passive smoking on the development of respiratory symptoms in young adults: an 8-year longitudinal study. J Clin Epidemiol 1996;49:581—6.
- ANSI/SHARE Standard 62.1-2004. Ventilation for acceptable indoor air quality. American Society of Heating Refrigerating and Air-Conditioning Engineers, Atlanta, GA., USA, 2004.
- Office of Environmental Health Hazard Assessment. Health effects of exposure to environmental tobacco smoke. Sacramento: Environmental Protection Agency, 1997. http://www.oehha.ca.gov/air/environmental tobacco/2005etsfinal.html.
- World Health Organization. Workers' health: global plan of action. Sixtieth World Health Assembly, 2007. http://www.who.int/occupational_health/WHO_health_ assembly en web.pdf (accessed 15 Jan 2008).

Correction

Lee JGL, Goldstein AO, Kramer KD, Steiner J, Ezzell MM, Shah V. Statewide diffusion of 100% tobacco-free college and university policies. *Tob Control* 2010;**19**:311–17. The authors should read as follows: Lee JGL, Goldstein AO, Kramer KD, Steiner J, Mathew M, Ezzell MM, Shah V.

Tobacco Control 2011;20:63. doi:10.1136/tc.2009.032888corr2