

A review of risk perception measurement in tobacco control research

Annette R Kaufman ¹, Alexander Persoskie,² Jenny Twesten,³ Julie Bromberg³

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/tobaccocontrol-2017-054005>).

¹Tobacco Control Research Branch, Behavioral Research Program, Division of Cancer Control and Population Sciences, National Cancer Institute, National Institutes of Health, Rockville, Maryland, USA
²Office of Science, Center for Tobacco Products, US Food and Drug Administration, Silver Spring, Maryland, USA
³BLH Technologies, Inc., Rockville, Maryland, USA

Correspondence to

Dr Annette R Kaufman, Tobacco Control Research Branch, Behavioral Research Program, Division of Cancer Control and Population Sciences, National Cancer Institute, National Institutes of Health, Rockville, MD 20850, USA; kaufmana@mail.nih.gov

Received 22 August 2017
Revised 22 November 2017
Accepted 28 November 2017
Published Online First
6 February 2018

ABSTRACT

Objective To describe the characteristics of risk perception measures used in tobacco control research and to evaluate whether these measures incorporate measurement suggestions put forward by risk perception measurement scholars.

Data sources Three databases (PubMed, PsycINFO and Web of Science) were searched in March 2015 for published English language peer-reviewed articles measuring tobacco risk perceptions (n=2557). The search string included terms related to tobacco products, perceptions and risk.

Study selection Three coders independently coded abstracts for initial inclusion. In total, 441 articles met the initial inclusion criteria, and 100 were randomly selected for a full-text review.

Data extraction A codebook was developed and tested through a training phase. Three coders independently coded the characteristics of each article (eg, population), multi-item measure (eg, validity) and item (eg, likelihood, affect, health outcome). Fifty-four articles, 33 measures and 239 items were coded.

Data synthesis Twenty-one articles had a multi-item risk perception measure, and 12 articles had one risk perception item. Many of the items asked about general health outcomes (36%), did not specify the person for whom risk was being judged (44%; eg, self, average person) or did not specify the conditions of use (27%; eg, the product used, intensity of use).

Conclusions There is little consistency across risk perception measures in tobacco research. There may be value in developing and disseminating best practices for assessing tobacco risk perceptions. A set of risk perception consensus measures may also benefit researchers in the field to help them consistently apply measurement recommendations.

INTRODUCTION

Risk perceptions—subjective judgements about the potential harm to health related to a hazard—are important constructs in health behaviour and decision-making theories.^{1–4} Risk perceptions can motivate people to avoid health hazards and to engage in health-protective behaviours.⁵ The strength of these motivational effects may depend on other factors such as whether risk appraisals, response efficacy and self-efficacy are high.⁶ Risk perceptions play a role in the experimentation with and initiation of tobacco use,^{7–8} cessation of tobacco use,^{9–11} the frequency and intensity of tobacco use,¹² and switching from one tobacco product to another.^{13–14} In tobacco control research, risk perceptions are often evaluated to examine the impact of advertising (eg, descriptors^{15–16},

packaging,^{17–20} warning labels,^{21–22} flavours (eg, menthol²³), educational interventions (eg, for water pipe²⁴) and product design (eg, filters²⁵) on users and non-users of tobacco products. Perceptions of risk of tobacco products are also relevant to how physicians address the treatment of tobacco use among their patients (eg, recommending cessation and delivering the five As¹³).

Despite the theoretical importance of risk perceptions, measures of perceived risk are not always associated with tobacco use behaviours. For example, among current cigarette smokers, some studies have found no association between risk perceptions and quit intentions and subsequent quitting behaviour, and others have found only weak associations.^{9–10–26–30} Moreover, some studies have observed a lack of association, small associations, and even reverse associations between risk perceptions and smoking behaviour.^{31–33} For instance, one study concluded that ‘the ability of these variables to explain individual variation is small’, with risk perceptions explaining only 5% of the variance in smoking behaviour beyond socio-demographic factors (Cutler and Glaeser, p241).³¹

One explanation for why risk perceptions are not always associated with tobacco use behaviours is that this association may be moderated by other factors. For example, one study found that the association between risk perceptions and smoking behaviour in youth and young adults was moderated by addiction beliefs and immediacy of health effects.⁷ Perceived health risk was a strong deterrent for those who believed that smoking was addictive and that smoking’s health effects occur rather immediately but not for those who viewed the addiction risk as low and the health effects as not immediate.⁷ The link between risk perceptions and smoking behaviour has also been shown to vary based on other factors, such as the extent to which people worry about their risk,³⁴ the extent to which people have high response efficacy and self-efficacy for protective behaviours,⁶ and the complexity of the smoking behaviour (eg, planning to quit vs sustained quitting¹⁰).

Another potential explanation for the variability of findings is that some measures of risk perception may be more or less valid than others. Early research in the field of risk perception revealed that risks are more likely to motivate protective actions depending on their characteristics, such as whether the hazard evokes *dread* (eg, by being uncontrollable) and whether the hazard is *unknown* (eg, by being new and not yet understood by science).³⁵ Dual-process models of decision-making propose that humans have two parallel systems for processing and responding to risk³⁶: *system 1*



To cite: Kaufman AR, Persoskie A, Twesten J, et al. *Tob Control* 2020;**29**:s50–s58.

uses heuristics and automatic associations to make rapid, low-effort judgements, whereas *system 2* uses controlled, deliberative reasoning to make rule-based, effortful judgements. In general, *system 1* is the default processing system but can be overridden with conscious and effortful *system 2* processing. A key component of *system 1* is *affect* (ie, the immediate, reflexive feeling of the goodness or badness of a stimulus³⁷), which plays an important role in guiding decision-making.³⁸ Research has uncovered an important bias known as *unrealistic optimism*, which is the tendency for people to minimise the extent to which they see themselves as at risk even when they recognise risks to other people.³⁹

Based on advances in risk perception research and the field of survey design, researchers have identified important considerations for assessing risk perceptions in applied contexts. Brewer and colleagues¹ made suggestions for designing risk perception measures using influenza as the example health risk, including (1) focusing on specific risks and outcomes rather than vague ones, (2) identifying the person for whom the risk is being evaluated, (3) making the question contingent on behaviour (ie, specifying the level of risk exposure) and (4) identifying a risk timeframe. These recommendations are aimed at standardising the risk that people are evaluating (ie, removing ambiguity) and increasing the likelihood that people provide perceptions of the same risk when responding to risk perception questions. For example, when asked a non-specific question about one's own risk of developing lung cancer, a current smoker may estimate her risk as low, because she is expecting to quit in the near future. In contrast, another current smoker may estimate her own risk as high because she does not expect to quit. This inconsistency can be resolved by instead asking people to evaluate their risk of developing lung cancer if they continue smoking at their current rate. Failure to adhere to such measurement suggestions may lead to weak and inconsistent associations between risk perceptions and behaviour. Indeed, Brewer and colleagues¹ found that risk perceptions for getting influenza were a stronger predictor of influenza vaccination behaviour when the measures adhered to their measurement suggestions.

A report by the Institute of Medicine (IOM) on studies of modified risk tobacco products also offers suggestions for how to measure risk perceptions.⁴⁰ Consistent with Brewer and colleagues,¹ the IOM suggested using *conditional* risk scenarios rather than *unconditional* risk assessments and assessing perceptions of *specific* tobacco-related outcomes rather than *general* perceptions of harm.⁴⁰ The IOM report also suggested assessing *affective* reactions to risk⁴⁰ consistent with research demonstrating the importance of affect in judgement and decision-making.^{38 41} Moreover, this suggestion is supported by research indicating that risk perception measures are more predictive of health-protective behaviours when measures ask about people's *feelings* of risk rather than simply their cognitive probability judgements.^{42–44}

Finally, general scientific principles also stress the need to empirically test the reliability and validity of measures of psychological constructs such as risk perception.^{45 46} Measures should be reliable over time, internally consistent (in the case of multi-item scales) and valid for assessing perceived risk. In general, there are benefits to using multi-item scales rather than single items to measure psychological constructs, such as risk perception, including reduced measurement error and the ability to assess internal consistency. Also, given that multi-item scales tend to outperform single-item scales in terms of their predictive validity,⁴⁷ developing measures with multiple items for assessing risk perceptions is advisable.

Despite the importance of perceived risk and the breadth and volume of tobacco control research assessing risk perceptions, no studies have been conducted to review how this construct is measured across tobacco studies. Thus, the extent to which advances in the scientific understanding of risk perception have been incorporated into the field of tobacco control research remains unclear. Moreover, given that risk perception researchers have suggested ways to improve risk perception measures, there is an opportunity to evaluate whether tobacco studies typically reflect these considerations by incorporating the emerging guidance. This literature review is an effort to fill this gap by describing and evaluating the characteristics of risk perception items and measures used in tobacco control research.

METHODS

Data sources

In March 2015, we conducted a search of three databases (PubMed, PsycINFO and Web of Science) for published, peer-reviewed articles measuring tobacco risk perceptions with no date restrictions. The search string (see online supplementary file: Supplement A) included terms related to tobacco products, perceptions and risk.

Study selection

Figure 1 shows the article inclusion flowchart. The initial search produced 5478 articles from the three databases, of which 2927 (53.4%) were duplicates and 2551 were unique. Three reviewers independently coded abstracts for initial inclusion, double coding approximately 10% of the abstracts (n=251), with 89.6% agreement among reviewers on average. Articles were included for further review if they appeared to describe research assessing people's perceptions of risk in the context of tobacco products or tobacco use. Articles were excluded if they reported only qualitative data (n=67) or tests of tobacco-related knowledge (n=69) or were not assessing risk perceptions related to tobacco products or their use by an individual (n=1974). A total of 441 (17.2%) articles met the initial inclusion criteria.

Next, given the large number of articles (n=441), we randomly selected 100 (22.7%) articles for full-text review. We liken this approach to recruiting a random sample of human participants or taking a random sample of media or web content^{48–50} to draw inferences about a larger population. We included an article for coding if it stated that it assessed people's perceptions of risk or if any of the items met our definition of tobacco risk perception as *a participant's subjective judgement about the potential harm to health related to tobacco products or their use*. Examples of potential harms to health included cancer, heart disease, respiratory effects, oral effects and the risk of becoming addicted. After full-article review, 29 articles were excluded (see footnote in figure 1). Thus, 71 articles were identified as being related to tobacco risk perceptions and included in the review. If an article did not provide the risk perception survey item(s) or an adequate description of the item(s), the authors of the publication were contacted to provide this information. For measure and item-level coding, an additional 17 articles were excluded because of insufficient information.

Data extraction

We coded at the article, multi-item measure and item levels. Multi-item measures and items were coded only if they were about risk perceptions. Within a multi-item measure, items that did not meet the inclusion criteria were excluded from coding (eg, an item assessing harm from secondhand smoke).

Review

We developed a codebook and tested it through a training phase in which all authors coded the first 10 articles. Three reviewers then independently coded 20% (n=11) of the articles, and 90% agreement among reviewers was achieved across all codes. Discrepancies were resolved through discussions among the reviewers and lead authors.

Data synthesis

Population

Articles were coded based on the study populations and included youth (ages younger than 18), adults (ages 18 and older) or both.

Research design

Articles were coded for being prospective (same participants assessed more than once over time), cross-sectional (assessed at a single point in time), experimental (random assignment, including both prospective and cross-sectional designs) or other.

Citation of source


Items and measures were coded based on whether the article cited the source of the item or measure (eg, a prior study) or stated that it had been adapted or used verbatim from a previous study.

Validity and reliability

Items and measures were coded as valid or reliable if the study completed validity or reliability testing or if the authors stated that the items or measures were previously shown to be valid or reliable.

Exact wording

Items were coded based on the amount of information provided about the questions and response options. (For example, did the article include the exact wording of the item, partial wording or no wording? If the latter two, we coded the item if the author sent exact wording after being contacted for the information.)



*Twelve articles did not include survey items or measures about tobacco risk perceptions, 6 articles only assessed knowledge of tobacco health risks, 1 article used qualitative measurement, 10 articles included tobacco risk perception items but did not meet other inclusion criteria (e.g., conference abstract, literature review)

Figure 1 Risk perception literature review flow chart.

Table 1 Risk perception item characteristics

Risk perception item characteristics	Description, (code)	Example (response options) (code)
Likelihood, absolute	Perceived probability that one will be harmed by tobacco product use (yes/no)	How likely are you to get lung cancer? (very low, somewhat low, moderate, somewhat high, very high) ⁵¹ (yes)
Likelihood, comparative	Perceived probability that one will be harmed by tobacco product use compared with another person, another product or another behaviour (yes/no)	Compared with others your same age and sex, how would you rate your risk of having a heart attack within the next 10 years? (1. much lower than average, 2. lower than average, 3. about average, 4. higher than average, 5. much higher than average) ⁵² (yes)
Conditional	Consideration of future outcomes with respect to tobacco product use (often hypothetical) in which they might engage, with the item being contingent on certain factors (frequency/time/product/some combination of these/no)	I can smoke a couple of cigarettes a day and still not harm my health (strongly agree, agree, neither agree nor disagree, disagree, strongly disagree) ⁵³ (frequency and product)
Risk target	The person who experiences the harm (self (eg, what is your risk)/specific other (eg, 50-year-old male's risk)/average other (eg, average person's risk)/general (eg, what is the risk?))	How concerned are you about getting lung cancer in your lifetime? (four-point Likert scale, 1—not at all to 4—very much) ⁵⁴ (self)
Severity	Perceived extent of harm that tobacco product use would cause (yes/no)	How serious would the health consequences be if you developed lung cancer? (1=not at all, 2=a little, 3=somewhat, 4=quite, 5=extremely) ⁵⁵ (yes)
Affect	Emotional response to tobacco product use (eg, fear, worry, disgust) (yes/no)	How often do you worry about getting lung cancer? Would you say... (rarely or never, sometimes, often, all the time) ⁵⁶ (yes)
Controllable	Ability to take action to reduce harm (yes/no)	Menthols are less harmful to me than non-menthols. (each (scale) rated on a four-point scale ranging from strongly agree to strongly disagree) ⁵⁷ (yes)
Known	Harms being well known to experts or others (yes/no)	The evidence indicating that smoking causes serious illness is very convincing (strongly agree, agree, don't know, disagree, strongly disagree) ⁵⁸ (yes)
Health outcome	Health effect caused by tobacco (specific (eg, cancer, heart disease)/general (eg, harm, risk, risk of disease)/hybrid (eg, risk of disease such as cancer))	What is the likelihood of getting addicted when using a water pipe socially? (none, low, medium, high) ⁵⁹ (specific)

Risk perception characteristics

Items were coded based on their risk perception characteristics. These characteristics included whether the items assessed absolute or comparative likelihood; were contingent on behaviour (ie, particular product, frequency of use, timeframe or some combination of these); identified the risk target; and assessed risk severity, affective responses to risk, the controllability of the risk, the extent to which the risk is known to experts or others and general perceptions of harmfulness. Table 1 provides descriptions for each of these characteristics.

Response options

The response options for each item were coded based on whether they used a numeric judgement scale (eg, “out of 100 smokers, how many do you think will get lung cancer due to their smoking?”) or a Likert-type rating scale (ie, any ordinal scale response with verbally labelled options). ‘Other’ response options included dichotomous (eg, yes/no) or non-ordinal categories (eg, males, females, both equally likely). We also created a response transformation code to capture whether the response options from one or more items were combined by taking the difference or product between items. This code did not include collapsing response options or instances in which items were simply combined into a risk perception index, which we refer to as a measure.

Additional analyses

Analyses were conducted to examine different combinations of item characteristics. To describe items that assessed product

perceptions, we examined items that were coded as both conditional on product and general health outcome harm. We examined severity items that were also coded with the following item characteristics: likelihood, health outcome and risk target. We examined the items assessing affect and their overlap with health outcome and risk target.

RESULTS

Study characteristics

Fifty-four studies published between 1981 and 2015 were included in the review and can be found in online supplementary file: Supplement B.^{12 51–103} Sixty-nine per cent (n=37) of the articles focused only on adults, 19% (n=10) of the articles focused only on youth, and 13% (n=7) of the articles included both youth and adults. Seventy-four per cent (n=40) of the studies were cross-sectional, 18.5% (n=10) were prospective, 5.5% (n=3) were experimental and 1.9% (n=1) were other (repeated measures).

Multi-item measure descriptive statistics

Of the 54 articles, 21 (39%) reported using a multi-item risk perception measure (ie, two or more items combined into a single scale or index). Across the 21 articles that used multi-item risk perception measures, a total of 33 measures were coded. Six of these measures (18%) were used exactly and four (12%) were adapted from a previously published study. For the other 23 measures (70%), the article did not provide a citation or source for the origin of the measure. Only four of the measures

Table 2 Number of risk perception items across articles

Number of risk perception items	Number of articles
One item	12 (22%)
Two items	8 (15%)
Three items	9 (17%)
Four items	5 (9%)
Five items	5 (9%)
Six items	4 (7%)
Seven items	2 (4%)
Eight items	2 (4%)
Nine items	2 (4%)
Ten or more items	5 (9%)
Total=239	Total=54

(12%) in two separate articles were described as being previously validated; neither reported any validity testing. Twenty-three measures (70%) across 14 studies assessed the reliability of the measure using Cronbach's alpha.

Characteristics of risk perception items

Table 2 shows the number of risk perception items included across articles. Many articles included multiple risk perception items, such that 239 items were coded across the 54 articles. The number of items per article ranged from 1 to 18, with the average number of risk perception items being 4.4 per article. Forty-one items (17%), including those within and not within multi-item measures, were used exactly as used in another cited study.

Table 3 presents the risk perception item descriptive statistics. More than half (57%) of the risk perception items included a specific health outcome. Slightly less than half (48%) of the items assessed risk for the self, and 42% (n=100) assessed risk generally (eg, 'It seems like almost everything causes cancer'). Three quarters of the items (n=179) assessed absolute risk and 22% (n=52) assessed comparative risk. Few items assessed severity (n=27, 11%), affect (n=22, 9%), controllability (n=19, 8%) or known risk (n=8, 3%).

Nearly 73% (n=173) of the items included consideration of specific (often hypothetical) behaviours, such as the use of a particular product, frequency of use, timeframe or some combination of these. Seventy per cent (n=167) of the items asked about risk conditional on using a particular product (eg, 'If I smoke cigarettes, I will live for a long time'). Only 13% (n=31) of the items specified a frequency of use (eg, 'I can smoke a couple of cigarettes a day and still not harm my health') and 8% (n=20) specified the timeframe (eg, 'Compared with others your same age and sex, how would you rate your risk of having a heart attack within the next 10 years?' or 'There's no risk of getting cancer if someone only smokes a few years'). Only 5% (n=12) of items specified all of the above (ie, product, frequency and timeframe) (eg, 'Now imagine that you continued to smoke about 2 OR 3 cigarettes each day for the rest of your life. What is the chance that you will get lung cancer?').

Most (n=183, 77%) of the item responses used a Likert-type rating scale and few items (n=15, 6%) used a numeric judgement scale. All numeric risk estimates were for absolute risk perception items. Of these, more than half (n=8, 53%) were for risk estimates for the self, whereas 20% (n=3) were for specific others and 27% (n=4) were for general risk. In addition, 73% of the numeric risk estimate response options were for specific risks rather than general risks (n=4, 27%). Few articles reported

Table 3 Risk perception item descriptive statistics

Variable	Percentage	N
Health outcomes		
Specific health outcome	57	137
General health outcome	36	85
Hybrid	5	11
Not applicable	3	6
Risk type		
Self	48	114
Specific other	8	19
Average other	<1	1
General	42	100
Not applicable	2	5
Absolute or comparative		
Absolute	75	179
Comparative	22	52
Not applicable	3	8
Conditional		
Product*	54	129
Time	1	3
Other	1	3
Frequency and product	8	19
Time and product	2	5
Frequency, time and product	5	12
Product and other	1	2
Unknown/not applicable	<1	1
Not assessed	27	65
Severity	11	27
Affect	9	22
Controllability	8	19
Risks known to experts or others	3	8
Numeric risk estimate	6	15
Rating scale (Likert type)	77	183
Other response option	17	41
Transformation/combination of items	12	28

*Sixty-two per cent (n=148) of all items coded assessed the risk associated with cigarette smoking, and 26% did not include a product (eg, 'How likely are you to get lung cancer?'). The remaining items assessed the risk of hookah smoking (n=12, 5%); menthol cigarettes (n=5, 2%); e-cigarettes (n=3, 1%); bidis (n=3, 1%); roll-your-own, pipes and cigars (n=3, 1%); reduced-risk-labelled cigarettes (n=2, 0.8%); or general tobacco (n=1, 0.4%) (data not shown). In some instances, both cigarettes and other products were assessed together (n=12, 5% eg, 'How do you compare Shisha and cigarette smoking considering their health effects?').

combining responses across different items (eg, by taking the difference or product between items) (n=8, 15%). Among those items that were transformed (n=28), 57% (n=16) used a rating scale whereas 11% (n=2) used a numeric estimate.

Further analyses revealed that 32% (n=82) of the items assessed perceived product harm (ie, items that were coded as both conditional on product and general health outcome: 'Do you think smoking tobacco is harmful to your health?') and that 55% (n=132) of items assessed risk perceptions of health outcomes (ie, items that were coded as self-risk, specific other risk, or average other risk and all health outcomes). Seventy-eight per cent (n=21) of the severity items assessed absolute risk, whereas 15% (n=4) assessed comparative risk. Of the articles that assessed severity (n=27), 63% (n=17) assessed a specific risk, 26% (n=7) assessed a general risk and 11% (n=3) assessed a hybrid of both. In addition, 44% (n=12) of those that assessed severity were for risk for the self and 41% (n=11) assessed

general risk. All items assessing affect were for the self. For those items assessing affect ($n=22$), 64% ($n=14$) assessed a specific risk, 23% ($n=5$) assessed a general risk and 9% ($n=2$) assessed a hybrid of both.

DISCUSSION

To our knowledge, this is the first review examining how risk perceptions are measured in tobacco control research. It is clear that risk perceptions are indeed measured in a large number of studies and that there is wide variation in the types of risk perception items used. There is a wealth of scientific research identifying important aspects of how people perceive risk and the components of risk perceptions that motivate behaviour. This study revealed that, among the studies reviewed, many did not incorporate advances from the field of risk perception.^{1 40}

More than half of the studies used single items to assess risk perception, even though using multi-item measures can increase statistical power.⁴⁷ The reliability and validity of measures are integral to the interpretability of research involving psychological constructs such as risk perception. In studies of tobacco risk perception, when small or no effects are observed, the reason may be the use of single-item measures with low reliability and validity. In this review, we found almost no validity testing and minimal reliability testing. Items that assess different aspects of risk perception (eg, absolute vs comparative) should not be combined into measures without psychometric or other testing showing that they can be combined.

The items and multi-item measures used in the studies in this review generally did not comply with the recommendations put forward by Brewer and colleagues¹ and the IOM,⁴⁰ although some recommendations were more likely to be met than others. The extent to which items specified various aspects of the risk to be judged (eg, the heaviness of product use) was highly variable across items. The health risk or outcome was specified in 57% of the items analysed in this review. The tobacco product was specified in more than half of the items (70%), but 27% of the items were not conditional on any aspect of use including the product under examination, heaviness of use or timeframe. Almost half of the risk perception items (48%) assessed risk for the self, but 42% assessed general risk without specifying the person at risk. Clearly, it is not possible to specify all aspects of product use that could contribute to one's risk, as this would cause risk perception items to become too cognitively burdensome. However, ideally, items should be specific about the key aspects of the product use behaviour under consideration (eg, frequency of use) and should assess the risk for the self.^{1 40 104–110} It may also be useful to assess whether particular risk outcomes are salient to individuals.¹¹¹

The IOM report recommended using either response scales with numerical likelihood estimates (ie, 0%–100%) or comparative risk assessments (eg, 'Compared to [another product], is this modified risk tobacco product (MRTP) more or less likely to cause [a specific outcome]?') instead of scales with verbal labels (eg, 'very likely').⁴⁰ This recommendation may have been aimed at reducing noise in responses (eg, differences in interpreting verbal labels) or increasing the interpretability of estimates by using an objectively defined response scale.¹¹² However, a prior study comparing percentage scales with response scales with verbal labels found that the scale with verbal labels was a better predictor of behavioural intentions and individual preferences than the scale with numeric percentage labels.¹¹³ Also, it is not possible to infer the accuracy of risk perceptions by eliciting numeric estimates and comparing them with objective levels

of risk.^{114–116} In our review, few measures included numeric response scales, with more than three-fourths using scales with verbal labels. The IOM report also recommended comparative response scales, but only 22% of the items reviewed here assessed direct comparative risk perception.⁴⁰ The evidence is mixed regarding how and when absolute vs comparative risk perception measurement is most predictive of outcomes of interest, including behaviour.^{117 118} A recent review of the literature on perceptions of relative risk from various tobacco and nicotine products found variation in how the perceptions were measured.¹¹⁹ Future research examining these important issues can inform risk perception measurement in tobacco control research.

Recognising the role of *affect* in risky decision-making⁴¹ has allowed researchers to 'explain a wide range of phenomena that have resisted interpretation in cognitive-consequentialist terms' (Loewenstein *et al*, p267)³⁸ In tobacco control, measures that focus only on cognitive types of perceptions may overlook affect-laden components that are strong motivators of behaviour and are sensitive to factors such as marketing and advertising. However, this review found that only 9% of risk perception items included affect. It is possible that including affect in risk perception measures might improve our ability to predict important tobacco outcomes, including use behaviour.

Although not necessarily recommended as part of the standard assessment of risk perceptions, researchers have also identified certain aspects of risk perceptions that play an important role in motivating protective behaviours.^{5 6} These include whether the hazard is dreaded and unknown. We found that few studies assessed components of these risk perception factors such as the perceived controllability of the risk, its severity, and the extent to which it is seen to be novel and unknown to science. Further research measuring these aspects of perceived risk may be informative.

This review must be viewed in light of several limitations. Although this was not a comprehensive review of all studies, we selected a random sample of studies for analysis to reduce bias. While some studies preceded publication of Brewer and colleagues' paper, when we examined the subset of articles published afterwards, we found that the characteristics of the items were similar to what is reported here.

What this paper adds

- ▶ In tobacco control research, risk perceptions are beliefs about the potential harms to health from using tobacco products and are widely assessed across the field.
- ▶ Risk perceptions are associated with numerous tobacco-related health behaviours, such as initiation, cessation and product switching.
- ▶ There is currently no consensus on how to best measure tobacco risk perceptions, but risk perception researchers have put forward some general guidelines for risk perception measures.
- ▶ This study provides the first review of risk perception measurement in tobacco control research.
- ▶ Previously developed suggestions for risk perception measurement have not been consistently incorporated in tobacco research, making detecting effects and associations concerning this important construct more difficult.

CONCLUSIONS

Suggestions for how to measure risk perceptions^{14 45 46} have not been consistently incorporated in tobacco research, which may make detecting significant associations and effects more difficult. These results indicate that the field may benefit from the development and dissemination of risk perception measurement best practices and specific guidelines for risk perception measurement in tobacco control research. These tobacco-specific guidelines can take into account critical nuances and special considerations. Most importantly, we hope to encourage a dialogue around risk perception measurement among tobacco control researchers. This review highlights the importance and potential benefits of risk perception measurement harmonisation in tobacco control research.

Acknowledgements We thank Alexandra Stern for her contributions to the inclusion criteria, codebook development and initial article coding. We thank William Klein, Michele Bloch and Jerry Suls for their feedback on this manuscript.

Contributors ARK conceptualised the study and led the project and writing. All authors contributed to the development of the coding scheme. JB and JT conducted the coding and analyses and drafted the methods. ARK and AP reviewed the codes and results. All authors contributed to the writing and revision and approved the final version of the manuscript.

Disclaimer The views and opinions expressed in this manuscript are those of the authors only and do not necessarily represent the views, official policy, or position of the U.S. Department of Health and Human Services or any of its affiliated institutions or agencies.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2020. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

ORCID iD

Annette R Kaufman <http://orcid.org/0000-0002-6696-7025>

REFERENCES

- Brewer NT, Chapman GB, Gibbons FX, *et al.* Meta-analysis of the relationship between risk perception and health behavior: the example of vaccination. *Health Psychol* 2007;26:136–45.
- Glanz K, Rimer BK, Viswanath K, eds. *Health behavior and health education: theory, research, and practice*. 4th edn. San Francisco: CA: Jossey-Bass, 2008.
- Janz NK, Becker MH. The health belief model: a decade later. *Health Educ Q* 1984;11:1–47.
- Weinstein ND. The precaution adoption process. *Health Psychol* 1988;7:355–86.
- Weinstein ND. Testing four competing theories of health-protective behavior. *Health Psychol* 1993;12:324–33.
- Sheeran P, Harris PR, Epton T. Does heightening risk appraisals change people's intentions and behavior? A meta-analysis of experimental studies. *Psychol Bull* 2014;140:511–43.
- Gerking S, Khaddaria R. Perceptions of health risk and smoking decisions of young people. *Health Econ* 2012;21:865–77.
- Song AV, Morrell HE, Cornell JL, *et al.* Perceptions of smoking-related risks and benefits as predictors of adolescent smoking initiation. *Am J Public Health* 2009;99:487–92.
- Borrelli B, Hayes RB, Dunsiger S, *et al.* Risk perception and smoking behavior in medically ill smokers: a prospective study. *Addiction* 2010;105:1100–8.
- Costello MJ, Logel C, Fong GT, *et al.* Perceived risk and quitting behaviors: results from the ITC 4-country survey. *Am J Health Behav* 2012;36:681–92.
- Jacobson JD, Catley D, Lee HS, *et al.* Health risk perceptions predict smoking-related outcomes in Greek college students. *Psychol Addict Behav* 2014;28:743–51.
- Hampson SE, Andrews JA, Barkley M, *et al.* Personality traits, perceived risk, and risk-reduction behaviors: a further study of smoking and radon. *Health Psychol* 2006;25:530–6.
- Fiore MC, Jaén CR, Baker TB, *et al.* *Treating tobacco use and dependence: 2008 update. Clinical practice guideline*. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, 2008.
- Tomar SL, Hatsukami DK. Perceived risk of harm from cigarettes or smokeless tobacco among U.S. high school seniors. *Nicotine Tob Res* 2007;9:1191–6.
- Cummings KM, Hyland A, Bansal MA, *et al.* What do Marlboro Lights smokers know about low-tar cigarettes? *Nicotine Tob Res* 2004;6(Suppl 3):323–32.
- Czoli CD, Hammond D. Cigarette packaging: Youth perceptions of “natural” cigarettes, filter references, and contraband tobacco. *J Adolesc Health* 2014;54:33–9.
- Adkison SE, Bansal-Travers M, Smith DM, *et al.* Impact of smokeless tobacco packaging on perceptions and beliefs among youth, young adults, and adults in the U.S: findings from an internet-based cross-sectional survey. *Harm Reduct J* 2014;11:2.
- Bansal-Travers M, Hammond D, Smith P, *et al.* The impact of cigarette pack design, descriptors, and warning labels on risk perception in the U.S. *Am J Prev Med* 2011;40:674–82.
- Hammond D, Parkinson C. The impact of cigarette package design on perceptions of risk. *J Public Health* 2009;31:345–53.
- Stead M, Moodie C, Angus K, *et al.* Is consumer response to plain/standardised tobacco packaging consistent with framework convention on tobacco control guidelines? A systematic review of quantitative studies. *PLoS One* 2013;8:e75919.
- Hammond D, Fong GT, McNeill A, *et al.* Effectiveness of cigarette warning labels in informing smokers about the risks of smoking: findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control* 2006;15(Suppl 3):iii19–25.
- Rodu B, Plurphanswat N, Hughes JR, *et al.* Associations of proposed relative-risk warning labels for snus with perceptions and behavioral intentions among tobacco users and nonusers. *Nicotine Tob Res* 2016;18:809–16.
- Wackowski OA, Delnevo CD, Lewis MJ. Risk perceptions of menthol cigarettes compared with nonmenthol cigarettes among New Jersey adults. *Nicotine Tob Res* 2010;12:786–90.
- Lipkus IM, Eissenberg T, Schwartz-Bloom RD, *et al.* Affecting perceptions of harm and addiction among college waterpipe tobacco smokers. *Nicotine Tob Res* 2011;13:599–610.
- O'Connor RJ, Bansal-Travers M, Cummings KM, *et al.* Filter presence and tipping paper color influence consumer perceptions of cigarettes. *BMC Public Health* 2015;15:1279.
- Avis NE, Smith KW, McKinlay JB. Accuracy of perceptions of heart attack risk: what influences perceptions and can they be changed? *Am J Public Health* 1989;79:1608–12.
- Clark MA, Kviz FJ, Crittenden KS, *et al.* Psychosocial factors and smoking cessation behaviors among smokers who have and have not ever tried to quit. *Health Educ Res* 1998;13:145–53.
- Kreuter MW, Strecher VJ. Changing inaccurate perceptions of health risk: results from a randomized trial. *Health Psychol* 1995;14:56–63.
- Norman P, Conner M, Bell R. The theory of planned behavior and smoking cessation. *Health Psychol* 1999;18:89–94.
- Schnoll RA, James C, Malstrom M, *et al.* Longitudinal predictors of continued tobacco use among patients diagnosed with cancer. *Ann Behav Med* 2003;25:214–21.
- Cutler DM, Glaeser E. What explains differences in smoking, drinking, and other health-related behaviors? *Am Econ Rev* 2005;95:238–42.
- Johnson RJ, McCaul KD, Klein WM. Risk involvement and risk perception among adolescents and young adults. *J Behav Med* 2002;25:67–82.
- Mills B, Reyna VF, Estrada S. Explaining contradictory relations between risk perception and risk taking. *Psychol Sci* 2008;19:429–33.
- Klein WM, Zajac LE, Monin MM. Worry as a moderator of the association between risk perceptions and quitting intentions in young adult and adult smokers. *Ann Behav Med* 2009;38:256–61.
- Fischhoff B, Slovic P, Lichtenstein S, *et al.* How safe is safe enough? A psychometric study of attitudes towards technological risks and benefits. *Policy Sci* 1978;9:127–52.
- Denes-Raj V, Epstein S. Conflict between intuitive and rational processing: when people behave against their better judgment. *J Pers Soc Psychol* 1994;66:819–29.
- Slovic P, Finucane M, Peters E, *et al.* The affect heuristic. In: Gilovich T, Griffin D, Kahneman D, eds. *Heuristics and biases: the psychology of intuitive judgment*. New York: Cambridge University, 2002:397–420.
- Loewenstein GF, Weber EU, Hsee CK, *et al.* Risk as feelings. *Psychol Bull* 2001;127:267–86.
- Weinstein ND. Unrealistic optimism about future life events. *J Pers Soc Psychol* 1980;39:806–20.
- Institute of Medicine. *Methods for studying risk perception and risk communication. scientific standards for studies on modified risk tobacco products*. Washington, DC: The National Academies Press, 2012:191–220.
- Slovic P, Peters E, Finucane ML, *et al.* Affect, risk, and decision making. *Health Psychol* 2005;24:S35–40.
- Ferrer RA, Klein WM, Persoskie A, *et al.* The tripartite model of risk perception (tririsk): distinguishing deliberative, affective, and experiential components of perceived risk. *Ann Behav Med* 2016;50:653–63.
- Taber JM, Klein WM. The role of conviction in personal disease risk perceptions: what can we learn from research on attitude strength? *Soc Personal Psychol Compass* 2016;10:202–18.

- 44 Weinstein ND, Kwitel A, McCaul KD, *et al.* Risk perceptions: assessment and relationship to influenza vaccination. *Health Psychol* 2007;26:146–51.
- 45 Allen MJ, Yen WM. *Introduction to measurement theory*. Prospect Heights, IL: Waveland Press 1979/2002.
- 46 Crano WD, Brewer MB, Lac A. *Principles and Methods of Social Research*. New York, NY: Routledge, 2015.
- 47 Diamantopoulos A, Sarstedt M, Fuchs C, *et al.* Guidelines for choosing between multi-item and single-item scales for construct measurement: a predictive validity perspective. *J Acad Mark Sci* 2012;40:434–49.
- 48 Deephouse DL. Media reputation as a strategic resource: an integration of mass communication and resource-based theories. *J Manage* 2000;26:1091–112.
- 49 Treude C, Barzilay O, Storey MA. *How do programmers ask and answer questions on the web?: NIER track. Paper presented at the 33rd International Conference on Software Engineering*. Honolulu, HI, 2011. https://www.researchgate.net/profile/Christoph-Treude/publication/221556094_How_Do_Programmers_Ask_and_Answer_Questions_on_the_Web_NIER_Track/links/02e7e52320b724ebd0000000/How-Do-Programmers-Ask-and-Answer-Questions-on-the-Web-NIER-Track.pdf (accessed 4 Oct 2017).
- 50 Ghosh S, Zafar MB, Bhattacharya P, *et al.* On sampling the wisdom of crowds: random vs. expert sampling of the twitter stream. *Paper presented at the 22nd ACM International Conference on Information & Knowledge Management*. San Francisco, CA, 2013. <https://people.mpi-sws.org/~gummadi/papers/sampletwitter.pdf> (accessed 4 Oct 2017).
- 51 Dillard AJ, McCaul KD, Klein WM. Unrealistic optimism in smokers: implications for smoking myth endorsement and self-protective motivation. *J Health Commun* 2006;11(Suppl 1):93–102.
- 52 Strecher VJ, Kreuter MW, Kobrin SC. Do cigarette smokers have unrealistic perceptions of their heart attack, cancer, and stroke risks? *J Behav Med* 1995;18:45–54.
- 53 Chassin L, Presson CC, Sherman SJ, *et al.* Long-term psychological sequelae of smoking cessation and relapse. *Health Psychol* 2002;21:438–43.
- 54 Lyna P, McBride C, Samsa G, *et al.* Exploring the association between perceived risks of smoking and benefits to quitting: who does not see the link? *Addict Behav* 2002;27:293–307.
- 55 Park ER, Gareen IF, Jain A, *et al.* Examining whether lung screening changes risk perceptions: National Lung Screening Trial participants at 1-year follow-up. *Cancer* 2013;119:1306–13.
- 56 Parascandola M, Augustson E, O'Connell ME, *et al.* Consumer awareness and attitudes related to new potential reduced-exposure tobacco product brands. *Nicotine Tob Res* 2009;11:886–95.
- 57 Unger JB, Allen B, Leonard E, *et al.* Menthol and non-menthol cigarette use among black smokers in Southern California. *Nicotine Tob Res* 2010;12:398–407.
- 58 Elkind AK. The effect of training on knowledge and opinion about smoking amongst nurses and student teachers. *J Adv Nurs* 1988;13:57–69.
- 59 Abughosh S, Wu IH, Peters RJ, *et al.* Ethnicity and waterpipe smoking among US students. *Int J Tuberc Lung Dis* 2012;16:1551–7.
- 60 Borgan SM, Marhoon ZA, Whitford DL. Beliefs and perceptions toward quitting waterpipe smoking among cafe waterpipe tobacco smokers in Bahrain. *Nicotine Tob Res* 2013;15:1816–21.
- 61 Bresnahan MJ, Zhuang J, Sun S. Influence of smoking norms and gain/loss antismoking messages on young Chinese adults. *Nicotine Tob Res* 2013;15:1564–71.
- 62 Cavazos-Rehg PA, Krauss MJ, Kim Y, *et al.* Risk factors associated with hookah use. *Nicotine Tob Res* 2015;17:1482–90.
- 63 Chassin L, Presson CC, Bensenberg M, *et al.* Predicting adolescents' intentions to smoke cigarettes. *J Health Soc Behav* 1981;22:445–55.
- 64 Choi K, Forster J. Characteristics associated with awareness, perceptions, and use of electronic nicotine delivery systems among young US Midwestern adults. *Am J Public Health* 2013;103:556–61.
- 65 Coleman T, Barrett S, Wynn A, *et al.* Comparison of the smoking behaviour and attitudes of smokers who believe they have smoking-related problems with those who do not. *Fam Pract* 2003;20:520–3.
- 66 Finney Rutten LJ, Augustson EM, Moser RP, *et al.* Smoking knowledge and behavior in the United States: sociodemographic, smoking status, and geographic patterns. *Nicotine Tob Res* 2008;10:1559–70.
- 67 Gerrard M, Gibbons FX, Benthin AC, *et al.* A longitudinal study of the reciprocal nature of risk behaviors and cognitions in adolescents: what you do shapes what you think, and vice versa. *Health Psychol* 1996;15:344–54.
- 68 Giuliani KK, Mire O, Ehrlich LC, *et al.* Characteristics and prevalence of tobacco use among Somali youth in Minnesota. *Am J Prev Med* 2010;39:S48–55.
- 69 Goel S, Singh RJ, D S, *et al.* Public opinion about smoking and smoke free legislation in a district of North India. *Indian J Cancer* 2014;51:330–4.
- 70 Gordon NA, Rayner CA. Smoking practices of dental and oral health students at the University of the Western Cape. *SADJ* 2010;65:304–8.
- 71 Hall MG, Ribisi KM, Brewer NT. Smokers' and nonsmokers' beliefs about harmful tobacco constituents: implications for FDA communication efforts. *Nicotine Tob Res* 2014;16:343–50.
- 72 Holtzman AL, Babinski D, Merlo LJ. Knowledge and attitudes toward hookah usage among university students. *J Am Coll Health* 2013;61:362–70.
- 73 Attah Johnson FY. Attitudes of Nigerian medical students towards use and abuse of tobacco, alcohol and drugs. *Drug Alcohol Depend* 1985;15:323–34.
- 74 Kandra KL, Ranney LM, Lee JG, *et al.* Physicians' attitudes and use of e-cigarettes as cessation devices, North Carolina, 2013. *PLoS One* 2014;9:e103462.
- 75 Keizer I, Gex-Fabry M, Eytan A, *et al.* Smoking in psychiatric inpatients: association with working status, diagnosis, comorbid substance abuse and history of suicide attempts. *Addict Behav* 2009;34:815–20.
- 76 Krosnick JA, Chang L, Sherman SJ, *et al.* The effects of beliefs about the health consequences of cigarette smoking on smoking onset. *J Commun* 2006;56(suppl_1):S18–37.
- 77 Li Q, Dresler C, Heck JE, *et al.* Knowledge and beliefs about smoking and cancer among women in five European countries. *Cancer Epidemiol Biomarkers Prev* 2010;19:2811–20.
- 78 Liu J-T, Hsieh C-R. Risk perception and smoking behavior: empirical evidence from Taiwan. *J Risk Uncertain* 1995;11:139–57.
- 79 Louwagie GM, Ayo-Yusuf OA. Tobacco use patterns in tuberculosis patients with high rates of human immunodeficiency virus co-infection in South Africa. *BMC Public Health* 2013;13:1031.
- 80 Lundborg P, Lindgren B. Do they know what they are doing? Risk perceptions and smoking behaviour among Swedish teenagers. *J Risk Uncertain*;28:261–86.
- 81 Manfredi C, Lacey L, Warnecke R, *et al.* Smoking-related behavior, beliefs, and social environment of young black women in subsidized public housing in Chicago. *Am J Public Health* 1992;82:267–72.
- 82 McBride CM, Halabi S, Bepler G, *et al.* Maximizing the motivational impact of feedback of lung cancer susceptibility on smokers' desire to quit. *J Health Commun* 2000;5:229–41.
- 83 Möller V, Erstad I, Zani D. Drinking, smoking, and morality: do 'drinkers and smokers' constitute a stigmatised stereotype or a real risk factor in the time of HIV/AIDS? *Soc Indic Res* 2010;98:217–38.
- 84 Morrell HE, Song AV, Halpern-Felsher BL. Predicting adolescent perceptions of the risks and benefits of cigarette smoking: a longitudinal investigation. *Health Psychol* 2010;29:610–7.
- 85 Muula AS, Siziya S. Prevalence and determinants of ever smoked cigarettes among school-going adolescents in Lusaka, Zambia. *Afr Health Sci* 2007;7:246–52.
- 86 Ortendahl M, Näsmän P. Perception of smoking-related health consequences among pregnant and non-pregnant women. *Am J Addict* 2007;16:521–7.
- 87 Partos TR, Borland R, Thrasher JF, *et al.* The predictive utility of micro indicators of concern about smoking: findings from the International Tobacco Control Four Country study. *Addict Behav* 2014;39:1235–42.
- 88 Portillo F, Antoñanzas F. Information disclosure and smoking risk perceptions. Potential short-term impact on Spanish students of the new European Union directive on tobacco products. *Eur J Public Health* 2002;12:295–301.
- 89 Price JH, Everett SA. Perceptions of lung cancer and smoking in an economically disadvantaged population. *J Community Health* 1994;19:361–75.
- 90 Pulvers K, Scheuermann TS, Romero DR, *et al.* Classifying a smoker scale in adult daily and nondaily smokers. *Nicotine Tob Res* 2014;16:591–9.
- 91 Rindfleisch A, Crockett DX. Cigarette smoking and perceived risk: a multidimensional investigation. *J Public Policy Mark* 1999;18:159–71.
- 92 Sansone GC, Raute LJ, Fong GT, *et al.* Knowledge of health effects and intentions to quit among smokers in India: findings from the Tobacco Control Policy (TCP) India pilot survey. *Int J Environ Res Public Health* 2012;9:564–78.
- 93 Shi R, Messaris P, Cappella JN. Effects of online comments on smokers' perception of anti-smoking public service announcements. *J Comput Mediat Commun* 2014;19:975–90.
- 94 Siziya S, Rudatsikira E, Muula AS, *et al.* Predictors of cigarette smoking among adolescents in rural Zambia: results from a cross sectional study from Chongwe [corrected] district. *Rural Remote Health* 2007;7:728.
- 95 Song H, Kim J, Kim S-Y. Smoker identity among social smokers: theory-based approaches for anti-smoking interventions. *J Subst Use* 2014;19:346–50.
- 96 Stern MK, Wiens BA. Ethnic differences in adolescent perceptions of and attitudes toward substance use. *J Ethn Subst Abuse* 2009;8:54–69.
- 97 Unger JB, Cruz TB, Rohrbach LA, *et al.* English language use as a risk factor for smoking initiation among Hispanic and Asian American adolescents: evidence for mediation by tobacco-related beliefs and social norms. *Health Psychol* 2000;19:403–10.
- 98 Wakefield M, Ruffin R, Campbell D, *et al.* Smoking-related beliefs and behaviour among adults with asthma in a representative population sample. *Aust N Z J Med* 1995;25:12–17.
- 99 Wright AJ, Whitwell SC, Takeichi C, *et al.* The impact of numeracy on reactions to different graphic risk presentation formats: An experimental analogue study. *Br J Health Psychol* 2009;14(Pt 1):107–25.
- 100 Yeretizian JS, Afifi RA. "It won't happen to me": the knowledge-attitude nexus in adolescent smoking. *J Public Health* 2009;31:354–9.
- 101 Young D, Borland R, Hammond D, *et al.* Prevalence and attributes of roll-your-own smokers in the International Tobacco Control (ITC) Four Country Survey. *Tob Control* 2006;15(Suppl 3):iii76–82.

- 102 Yu C, Shi Y, Kadimpati S, *et al.* Perioperative smoking behavior of Chinese surgical patients. *Anesth Analg* 2013;116:1238–46.
- 103 Zawahir S, Omar M, Awang R, *et al.* Effectiveness of antismoking media messages and education among adolescents in Malaysia and Thailand: findings from the international tobacco control southeast Asia project. *Nicotine Tob Res* 2013;15:482–91.
- 104 Fischhoff B. The real world: what good is it? *Organ Behav Hum Decis Process* 1996;65:232–48.
- 105 Fischhoff B. What forecasts (seem to) mean. *Int J Forecast* 1994;10:387–403.
- 106 Ronis DL. Conditional health threats: health beliefs, decisions, and behaviors among adults. *Health Psychol* 1992;11:127–34.
- 107 van der Velde FW, Hooykaas C, van der Pligt J. Conditional versus unconditional risk estimates in models of aids-related risk behaviour. *Psychol Health* 1996;12:87–100.
- 108 Janssen E, van Osch L, de Vries H, *et al.* Measuring risk perceptions of skin cancer: reliability and validity of different operationalizations. *Br J Health Psychol* 2011;16:92–112.
- 109 Dillman DA, Smyth JD, Internet CLM. *phone, mail, and mixed-mode surveys: the tailored design method*. Hoboken, NJ: John Wiley & Sons, 2014.
- 110 Sudman S, Bradburn NM, Schwarz N. *Thinking about answers: the application of cognitive processes to survey methodology*. San Francisco, CA: Jossey-Bass, 1996.
- 111 Ajzen I, Fishbein M. *Understanding attitudes and predicting social behavior*. Englewood Cliffs, NJ: Prentice-Hall, 1980.
- 112 Kip Viscusi W. Comment: the perils of qualitative smoking risk measures. *J Behav Decis Mak* 2000;13:267–71.
- 113 Windschitl PD, Wells GL. Measuring psychological uncertainty: verbal versus numeric methods. *J Exp Psychol* 1996;2:343–64.
- 114 Persoskie A, Downs JS. Experimental tests of risk ladders in the elicitation of perceived likelihood. *J Behav Decis Mak* 2015;28:424–36.
- 115 Slovic P. Rejoinder: the perils of Viscusi's analyses of smoking risk perceptions. *J Behav Decis Mak* 2000;13:273–6.
- 116 Windschitl PD. Judging the accuracy of a likelihood judgment: the case of smoking risk. *J Behav Decis Mak* 2002;15:19–35.
- 117 Ranby KW, Aiken LS, Gerend MA, *et al.* Perceived susceptibility measures are not interchangeable: absolute, direct comparative, and indirect comparative risk. *Health Psychol* 2010;29:20–8.
- 118 Zajac LE, Klein WM, McCaul KD. Absolute and comparative risk perceptions as predictors of cancer worry: moderating effects of gender and psychological distress. *J Health Commun* 2006;11(Suppl 1):37–49.
- 119 Czoli CD, Fong GT, Mays D, *et al.* How do consumers perceive differences in risk across nicotine products? A review of relative risk perceptions across smokeless tobacco, e-cigarettes, nicotine replacement therapy and combustible cigarettes. *Tob Control* 2017;26:e49–58.