Tobacco outlet density and social disadvantage in New South Wales, Australia

There is evidence to suggest a possible association between ease of access to tobacco and uptake of smoking,^{1 2} and the likelihood of cessation.³ A recent analysis of tobacco outlet density (TOD) in the USA found that TOD was higher in areas where a higher proportion of Hispanics and African-Americans live, and in areas where a higher proportion of families live in poverty.⁴ The authors concluded that higher TOD may contribute to disparities in smoking prevalence. This sort of evidence has led to proposals that control of TOD be considered in tobacco control efforts.^{5 6}

In the Australian state of New South Wales (NSW), a requirement to notify the state government of intention to sell tobacco products became mandatory under the Public Health (Tobacco) Act 2008. The notification data was then acquired by the Cancer Council NSW under a Government Information (Public Access) request in 2011. Using these data, we aimed to examine possible associations between TOD and socioeconomic status and remoteness across NSW.

Of the 12 451 current tobacco retailers registered, 811 were excluded because addresses were incorrectly entered, or were unable to be matched to a local government area (LGA). The outcome of interest was TOD, defined as the number of tobacco retailers per 100 000 people. Socioeconomic disadvantage was measured for each LGA (n=138) using Socio-Economic Indexes for Areas (SEIFA).⁷ SEIFA is calculated by the Australian Bureau of Statistics using census data, including level of education, employment status and household income. LGAs vary significantly in size (from less than 10 km² to more than 50 000 km²) and population (from less than 20 000 to more than 300 000), so a measure of geographical remoteness was included using the Accessibility/Remoteness Index of Australia (ARIA) in order to account for this.⁸ Data on the LGA smoking prevalence was taken from the NSW Population Health Survey.⁹ The TOD was log-transformed, and the proportion of smokers in each LGA, SEIFA Score and ARIA mean were all standardised to adjust for differences in scale and mean. TOD was then regressed on the other variables. The median TOD was 21.72 per 100 000 people (table 1).

Table 1 Tobacco outlet density in NSW and associations with smoking prevalence, social disadvantage and remoteness

	Tobacco outlet density per 100 000 population†			
	Coefficient	Lower 95% CI	Upper 95% CI	p Value
Intercept	-5.67	-5.78	-5.55	<0.0001
Smoking prevalence ⁹	-0.05	-0.18	0.08	0.47
Social disadvantage (SEIFA Score) ⁷	-0.17	-0.32	-0.02	0.025
Remoteness (ARIA mean ⁸)	0.50	0.36	0.65	<0.0001
R squared	0.43			
Group p value	<0.0001			

†1st quartile: 6.503; Median: 21.720; Mean: 64.090; 3rd Quartile: 71.680.

ARIA, Accessibility/Remoteness Index of Australia; SEIFA, Socio-Economic Indexes for Areas.

A statistically significant relationship was found between TOD, social disadvantage (SEIFA; p=0.025) and remoteness (ARIA scores; p<0.0001), independent of smoking prevalence (table 1).

Our results strongly suggest that tobacco outlets are concentrated in areas of higher disadvantage, and that are at greater risk of poor health outcomes. That the association was evident even after controlling for smoking prevalence may reflect a deliberate strategy by the tobacco industry, rather than being a response to higher demand. Further, they are consistent with the recent US study.⁴

The strengths of this study lie in the comprehensive coverage of all types of tobacco retailers and in our ability to control for smoking prevalence, both identified as limitations in earlier studies.⁴ Further, to our knowledge, no other studies of this kind have been conducted in Australia.

More research is required to determine if different types of retail outlets are more concentrated in disadvantaged areas, as there is evidence that particular types of retail outlets decrease the likelihood of cessation.¹⁰ Additionally, longitudinal studies are needed to assess associations between TOD and smoking uptake and cessation.

This study suggests that greater attention needs to be devoted to monitoring TOD, so that tobacco control efforts can be appropriately targeted.

James Kite,¹ Chris Rissel,² Mark Greenaway,² Kelly Williams³

¹Cancer Prevention Division, Cancer Institute NSW, Alexandria, New South Wales, Australia

²Prevention Research Collaboration, School of Public Health, University of Sydney, Camperdown, New South Wales, Australia

³Cancer Council NSW, Woolloomooloo, New South Wales Australia

Correspondence to James Kite, Cancer Prevention Division, Cancer Institute NSW, Postal address: PO Box 41, Alexandria, NSW 1435, Australia; James.kite@cancerinstitute.org.au

Acknowledgements We would like to acknowledge Anne Jones for her role in facilitating the partnership between the Cancer Institute NSW and Cancer Council NSW. We would also like to thank Simon Chapman, Sarah Durkin and Kathy Chapman for providing comments on the draft manuscript.

Contributors JK and CR conceived the study. MG designed and carried out the analysis. KW provided the data. All authors contributed to interpretation of the analysis and writing the manuscript. JK wrote the first draft. All authors reviewed and approved the final draft.

Funding None.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

OPEN ACCESS

Open Access
Scan to access more
free content

What this paper adds

- Our findings indicate that there is an association between tobacco outlet density and social disadvantage and remoteness, after controlling for smoking prevalence, in New South Wales, Australia.
- This suggests that outlet density may reflect a deliberate strategy by the tobacco industry, and that further research should be conducted to explore the effects of outlet density on smoking behaviours.

Open Access: This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

To cite Kite J, Rissel C, Greenaway M, *et al.* *Tob Control* 2014;**23**:181–182.

Received 20 June 2011

Revised 14 November 2012

Accepted 20 November 2012

Published Online First 14 December 2012

Tob Control 2014;**23**:181–182.

doi:10.1136/tobaccocontrol-2012-050648

REFERENCES

- 1 Novak SP, Reardon SF, Raudenbush SW, *et al.* Retail tobacco outlet density and youth cigarette smoking: a propensity-modeling approach. *Am J Public Health* 2006;96:670–6.
- 2 Henriksen L, Feighery EC, Schleicher NC, *et al.* Is adolescent smoking related to the density and proximity of tobacco outlets and retail cigarette advertising near schools? *Prev Med* 2008;47:210–14.
- 3 Reitzel LR, Cromley EK, Li Y, *et al.* The effect of tobacco outlet density and proximity on smoking cessation. *Am J Public Health* 2011;101: 315–20.
- 4 Rodriguez D, Carlos HA, Adachi-Mejia AM, *et al.* Predictors of tobacco outlet density nationwide: a geographic analysis. *Tob Control* 2013;22: 349–355.
- 5 Cohen JE, Anglin L. Outlet density: a new frontier for tobacco control. *Addiction* 2009;104:2–3.
- 6 Paul CL, Mee KJ, Judd TM, *et al.* Anywhere, anytime: retail access to tobacco in New South Wales and its potential impact on consumption and quitting. *Soc Sci Med* 2010;71:799–806.
- 7 Australian Bureau of Statistics. *2006 census of population and housing: socio-economic indexes for area's (SEIFA)*. Canberra, Australia: Commonwealth of Australia, 2008.
- 8 Australian Bureau of Statistics. *Remoteness structure*. Canberra: Commonwealth of Australia, 2012.
- 9 Centre for Epidemiology and Research. *2010 Report on adult health from the New South Wales population health survey*. Sydney: NSW Department of Health, 2011.
- 10 Burton S, Clark L, Heuler S, *et al.* Retail tobacco distribution in Australia: Evidence for policy development. *Australas Mark J (AMJ)* 2011;19:168–73.