

Changes in tar yields and cigarette design in samples of Chinese cigarettes, 2009 and 2012

Liane M Schneller,¹ Benjamin A Zwierchowski,¹ Rosalie V Caruso,¹ Qiang Li,^{2,3} Jiang Yuan,³ Geoffrey T Fong,^{2,4} Richard J O'Connor¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/tobaccocontrol-2014-051803>)

¹Department of Health Behavior, Roswell Park Cancer Institute, Buffalo, New York, USA

²Department of Psychology, University of Waterloo, Ontario, Canada

³Chinese Centers for Disease Control and Prevention, Beijing, People's Republic of China

⁴Ontario Institute for Cancer Research, Toronto, Ontario, Canada

Correspondence to

Dr Richard J O'Connor, Department of Health Behavior, Roswell Park Cancer Institute, Elm and Carlton Streets, Buffalo, NY 14263, USA; Richard.Oconnor@roswellpark.org

Received 23 May 2014

Accepted 10 September 2014

ABSTRACT

Background China is home to the greatest number of smokers as well as the greatest number of smoking-related deaths. An active and growing market of cigarettes marketed as 'light' or 'low tar' may keep health-concerned smokers from quitting, wrongly believing that such brands are less harmful.

Objective This study sought to observe changes in cigarette design characteristics and reported tar, nicotine and carbon monoxide (TNCO) levels in a sample of cigarette brands obtained in seven Chinese cities from 2009 to 2012.

Methods Cigarettes were purchased and shipped to Roswell Park Cancer Institute, where 91 pairs of packs were selected for physical cigarette design characteristic testing and recording of TNCO values. Data analysis was conducted using SPSS, and was initially characterised using descriptive statistics, correlations and generalised estimating equations to observe changes in brand varieties over time.

Findings Reported TNCO values on packs saw mean tar, nicotine and CO levels decrease from 2009 to 2012 by 7.9%, 4.5% and 6.0%, respectively. Ventilation was the only cigarette design feature that significantly changed over time ($p < 0.001$), with an increase of 31.7%. Significant predictors of tar and CO yield overall were ventilation and per-cigarette tobacco weight, while for nicotine tobacco moisture was also an independent predictor of yield.

Conclusions The use of ventilation to decrease TNCO emissions is misleading smokers to believe that they are smoking a 'light/low' tar cigarette that is healthier, and is potentially forestalling the quitting behaviours that would begin to reduce the health burden of tobacco in China, and so should be prohibited.

INTRODUCTION

Low-tar cigarettes are widely promoted throughout China via advertising and cigarette packs with prominently displayed tar values.¹ The State Tobacco Monopoly Administration (STMA) has also reduced maximum limits on International Organisation for Standardization (ISO) tar ratings over time, from 17 mg in 2001 to 15 mg in 2004, to 12 mg in 2011.¹ This has helped to promote 'low-tar' cigarettes as less dangerous.² Indeed, China lags in broad public awareness that cigarettes marketed as 'low tar/light' are not necessarily less hazardous.^{3 4} The Global Adult Tobacco Survey (GATS) 2010 found that 86% of respondents thought that light/low-tar cigarettes were better for one's health compared to regular cigarettes.⁵ Even those with more education, such as healthcare

professionals and teachers, had high levels of misperception.⁵

Reduced tar and 'safer' cigarettes remain a priority for STMA. In 2009, over 3 billion Yuan (~US \$490 million) was expended on 'low-tar' cigarette research and development.¹ However, independent research shows that smoking cigarettes with lower machine yields of tar (according to ISO machine measures) is not associated with lower levels of nicotine metabolites or total polycyclic aromatic hydrocarbons (PAHs) in Chinese smokers' urine.² This is consistent with a large body of literature in Western countries pointing to little benefit from low-tar cigarettes.⁶ Reducing tar levels is typically achieved through cigarette engineering, changing features such as filter density, tipping paper length, filter ventilation levels and tobacco density.⁶ Our research group has shown that UK manufacturers complied with a 2 mg reduction in the tar ceiling enacted in 2004 (from 12 to 10 mg) by increasing filter ventilation, while other facets of engineering remained essentially unchanged.⁷ Typically, smokers of low-yield cigarettes can compensate for lowered yields of tar and nicotine by smoking more intensively (eg, more puffs, larger puffs, blocking filter vents, smoking more cigarettes).⁸ Thus a disconnect exists between machine-measured yields and exposures among smokers.

Following on our earlier report on Chinese cigarette design features,⁹ this paper presents data on the physical design characteristics of a variety of popular cigarette brands manufactured and sold in China in 2009 and 2012, coincident with Framework Convention on Tobacco Control (FCTC) implementation in China and a reduction in the tar limit from 15 to 12 mg.¹⁰ Thus, our study design was capable of assessing which changes in cigarette design were associated with this lowering of the limits on tar ratings, comparable to our earlier work on UK cigarettes.⁷ We sought to explore on which brands tar, nicotine and carbon monoxide yields (as reported on packages) declined and what changes in cigarette design might best explain those changes.

METHODS

The cigarettes analysed for this study were purchased at two time points (2009 and 2012) from seven cities: Beijing, Changsha, Kunming, Shanghai, Shenyang, Guangzhou and Yinchuan.¹¹ Field workers visited three large retail stores in each city and bought packs of cigarettes until every brand family and brand variety available in each year was purchased (total $n = 2,052$; 2009, $n = 907$; 2012, $n = 1145$). The packs were shipped to the

To cite: Schneller LM, Zwierchowski BA, Caruso RV, et al. *Tob Control* Published Online First: [please include Day Month Year] doi:10.1136/tobaccocontrol-2014-051803

Tobacco Research Laboratory at Roswell Park Cancer Institute (RPCI), where they were catalogued and stored unopened at -20°C until analysis. From this larger set we randomly selected 91 pairs of cigarettes based on Universal Product Code matching to measure tar and ventilation at 2009 and 2012. Before testing, in accordance with ISO 3402:1999, the packs were conditioned for a minimum of 48 h at $22\pm 2.0^{\circ}\text{C}$ and $60\pm 2.0\%$ relative humidity in an environmental chamber. Cigarette physical and design characteristics were assessed following previously published procedures by the same laboratory.^{12–14} For a given brand, five cigarettes were selected randomly from each pack after conditioning for physical analysis and the data averaged. Digital callipers were used to measure the entire cigarette length, the tobacco rod length and diameter and the filter length and diameter. Filter weight measurements were made gravimetrically using a Mettler-Toledo analytical balance. The tipping paper was removed from each filter, and its length and the presence of any vent holes were performed using a light box and a transparent ruler. The tobacco moisture and tobacco weight were analysed using a halogen moisture analyser (HR83 or HB43-S, Mettler-Toledo, Columbus, Ohio, USA). The moisture content was determined as the per cent change in weight after heating the tobacco from five cigarettes with a halogen bulb at 125°C . Filter ventilation and pressure drop were measured using a KC-3 apparatus (Borgwaldt-KC, Richmond, Virginia, USA). The level of the paper porosity was measured using the vacuum method on a PPM1000M device (Cerulean, Milton Keynes, UK). Tar, nicotine and carbon monoxide yield values were obtained from the product packages. Analyses were conducted from January to March 2013.

Data analysis was conducted using Statistical Package for the Social Sciences V21.0 (IBM, Armonk, New York, USA). The sample was initially characterised using descriptive statistics, and correlations of design characteristics were also analysed. Changes in brand varieties over time were assessed within-participants using generalised estimating equations (GEE). Models used a normal distribution with identity link function and exchangeable working correlation matrix unless otherwise noted.

RESULTS

Tar and ventilation levels

We identified eight brands as having major design changes in the 2-year study period (see online supplementary table S1). These included the addition of a two-part filter or a charcoal filter (examples in see online supplementary figure S1A), or implementing filter patterning (see online supplementary figure S1B). Huanghelou implemented a channelled filter design reminiscent of Barclay cigarettes, introduced in 1980 in the USA which has been shown to be particularly susceptible to blockage by the smoker in the normal course of puffing, yielding much greater tar delivery than its ISO tar rating.¹⁵

Tar, nicotine and carbon monoxide (TNCO) and cigarette design characteristics

Table 1 presents the design characteristics of Chinese cigarettes from 2009 and 2012. Between 2009 and 2012, mean ISO tar levels decreased by 7.9% (from 11.9 to 10.9 mg), nicotine levels decreased by 4.5% (from 1.1 to 1.0 mg), and CO levels decreased by 5.7% (from 12.7 to 12.0 mg). Mean ventilation levels increased substantially, by 31.7% (from 8.3% to 10.9%). Of the unventilated cigarettes in 2009, 4.4% became ventilated in 2012, while 2.2% of ventilated cigarettes in 2009 lost ventilation in 2012. Furthermore, 1.1% of charcoal-containing filtered cigarettes did not have charcoal filters in 2012, and 2.2% of

Table 1 Average Chinese TNCO and cigarette design characteristics by year

	Year	N	Mean	SD	Minimum	Maximum
Tar	2009	99	11.87	2.09	5.00	15.00
	2012	98	10.93	2.09	5.00	15.00
Nicotine	2009	99	1.05	0.21	0.00	1.40
	2012	98	1.01	0.19	0.40	1.30
CO	2009	96	12.70	1.94	6.00	15.00
	2012	93	11.98	2.00	6.00	15.00
Cigarette length (mm)	2009	99	83.80	2.07	73.49	93.84
	2012	98	83.94	2.44	73.79	96.85
Per cigarette weight (g)	2009	99	0.66	0.07	0.45	0.98
	2012	98	0.65	0.05	0.49	0.76
Tobacco rod length (mm)	2009	99	59.48	3.64	49.68	68.59
	2012	98	59.57	3.17	53.76	68.48
Filter length (mm)	2009	91	23.69	3.55	15.70	29.84
	2012	85	24.12	3.06	18.67	29.96
Tipping paper (mm)	2009	99	30.58	2.78	23.95	36.49
	2012	98	30.90	2.86	23.84	36.76
Cigarette pressure drop	2009	94	110.94	12.76	72.50	146.30
	2012	93	109.08	13.34	59.80	144.90
Ventilation (%)	2009	94	8.28	10.43	0.00	48.00
	2012	93	10.91	11.35	0.00	46.90
Cigarette paper vacuum porosity	2009	93	54.97	10.17	28.47	81.49
	2012	93	54.96	9.17	33.58	78.32
Tobacco moisture (%)	2009	99	15.55	2.42	10.59	18.08
	2012	97	15.97	1.46	12.04	18.81
Rod density	2009	99	252.75	23.62	157.67	343.26
	2012	98	253.40	20.09	192.47	316.10
Filter density	2009	91	121.09	23.45	42.80	343.26
	2012	84	119.11	10.93	97.01	198.13
Overwrap (mm)	2009	91	6.81	2.22	2.55	13.34
	2012	85	6.57	1.86	3.61	11.42

N=99 for 2009 and N=98 for 2012. Deviations due to unlabelled TNCO values, multipart filters that do not conform with measurement protocols, products over or undersized for measurement equipment and equipment failures. TNCO, tar, nicotine and carbon monoxide.

non-charcoal containing filtered cigarettes became charcoal-filtered in 2012.

Correlations of cigarette design characteristics and TNCO

Correlations among cigarette design characteristics and tar, nicotine and CO can be found in online supplementary table S2. Tar, nicotine and CO were significantly intercorrelated (all higher than 0.68). Additionally, ventilation had by far the strongest correlations with tar (-0.66), nicotine (-0.54) and CO (-0.74).

Generalised estimating equations of cigarette design characteristics by year

We next examined whether ventilation, tobacco weight, rod and filter density, tobacco moisture and pressure drop changed as a function of time (table 2). Significant changes with time were found for only ventilation ($B=2.33$, $\text{S.E.}=0.77$, $p=0.002$), which increased between 2009 and 2012. If the analysis is limited to only those cigarette brands with a tar yield above 12 mg in 2009 (ie, those needing to reduce tar to comply), mean ventilation shows an increase, although not statistically significant ($B=0.355$; $p=0.086$).

Relationship of cigarette design features to reported TNCO yields

Stepwise GEE models with unstandardised β estimates were conducted for each of tar, nicotine and carbon monoxide. Year was

Table 2 Changes in various cigarette design features from 2009 to 2012

Variable	B	SE	Hypothesis test		
			Wald χ^2	df	Significance
Ventilation*	2.33	0.77	9.178	1	0.002
Pressure drop	-1.52	1.37	1.223	1	0.269
Percent moisture	0.247	0.193	1.628	1	0.202
Filter density	-1.93	2.57	0.564	1	0.452
Rod density	0.27	2.61	0.010	1	0.918
Tobacco weight per cigarette	-0.01	0.006	2.556	1	0.110

*GEE with tweedie distribution (due to significant number of zero values) and log link function.

GEE, generalised estimating equations.

the primary predictor. Ventilation was the first design feature forced into the model because of its known association with TNCO, followed by tobacco weight, filter density, overwrap and tobacco moisture, based on their bivariate associations with TNCO (table 3). For tar as well as nicotine, predictors were ventilation ($p<0.001$) and per cigarette tobacco weight ($p=0.020$ and $p=0.011$, respectively). Predictors of CO were also ventilation and per cigarette tobacco weight, as well as filter density. Other measured design features did not contribute significantly to the model.

DISCUSSION

O'Connor and colleagues demonstrated that the UK tobacco industry met the 2001 European Union (EU) regulatory standard of lowering yields to tar (10 mg), nicotine (1 mg) and CO (10 mg) by increasing filter ventilation.⁸ The results of the present study, which examined changes in cigarette design characteristics and reported tar, nicotine and carbon monoxide levels in selected Chinese cigarette brands from 2009 and 2012, demonstrate that the Chinese tobacco industry met the 2011 requirement to reduce tar limits from 15 to 12 mg in a similar fashion. Cigarette ventilation, as expected, was the most important predictor of yields, and was the only design parameter that

significantly changed (a 31.7% increase) from 2009 to 2012, replicating the results found in the UK in 2004.^{16 17}

In addition, a subset of brands appear to be innovating in terms of filter design, employing filters with pictograms carved in the ends, channelled filters, dual filters and carbon filters. Future research will need to monitor such innovation as the Chinese market develops to determine if such products obtain significant market share. However, given the substantial evidence that increasing filter ventilation leads to no benefits in actual exposure and uptake, the public health consequences of the lowering of tar ratings in China are trivial if present at all. But in fact the publicity and marketing associated with China's increasing focus on lowering tar limits, coupled with the lack of real health benefits, portends a significant negative impact on public health because Chinese smokers are much more likely than smokers in other countries to believe that light/low-tar cigarettes are less harmful: 71% of Chinese smokers believe that light/low-tar cigarettes are less harmful,⁴ which suggests that the Chinese industry's campaign of promoting such cigarettes will continue to appeal to the 300 million Chinese smokers who are likely to be increasing in their awareness of the harms of smoking and will wrongly see light/low-tar cigarettes as a way to reduce their risk.¹ As a result, Chinese smokers may well respond to increasing awareness of health harms of cigarettes by switching brands with lower tar levels rather than by quitting.

As in Western countries from the 1950s through the 1980s, when cigarette smoking was definitively linked to increased disease risks, the Chinese tobacco industry has responded to increasing health concerns around smoking by promoting cigarettes that delivered less tar in measurements made by machine smoking of cigarettes using a fixed pattern of smoking.^{6 18–20} In the US, Canada and EU, research and court cases have demonstrated the fallaciousness of these products in terms of health benefit, and efforts have been made to counter market the products.^{6 21–23} However, in the Chinese case, there are structural reasons for the continued marketing of low-tar cigarettes despite a large body of evidence that they will not reduce disease burden. Importantly, there are broader political considerations intertwined in the domestic tobacco trade, such as tax revenues, which have important impacts on tobacco control efforts.²⁴ The decision-making structure of the implementation of the FCTC remains in the hands of the Ministry of Industry, Innovation and Technology (MIIT), of which STMA/CNTC is a part. STMA has invested heavily in 'low-tar' cigarette research, one of their important research and development goals.

The current study is limited in several ways. While significant differences were seen between TNCO yields and cigarette ventilation over time, the ISO yields reported on the packs were not tested directly by our laboratory. So it is possible for discrepancies to exist between labelled and measured values. ISO yields are problematic themselves, as they are not representative of smoking behaviour or exposure.^{6 19} Nonetheless, the consistency of findings with the existing literature is encouraging.

The promotion of light/low-tar cigarettes in China represents a significant barrier to reducing smoking in China, the most important preventable cause of death and disease in the world's largest country. China should not only ban misleading descriptors such as low tar, light as they already have, but also ban other misleading claims equating lower tar numbers with low harm or high technology. While removing misleading descriptors is helpful, it is not likely to lead to lasting reductions in misperceptions,²⁵ which appear to be driven in large part by filter ventilation.¹³ As others have recommended, China should also remove TNCO numbers from the pack, as they are irrelevant

Table 3 Primary predictors of TNCO yields

Dependent variable	Parameter	B	SE	Hypothesis test		
				Wald χ^2	df	Significance
Tar*	Year	-0.046	0.0113	16.986	1	<0.001
	Ventilation	-0.010	0.0015	46.037	1	<0.001
	Per cigarette tobacco weight	0.625	0.2694	5.383	1	0.020
Nicotine	Year	-0.029	0.0090	10.704	1	0.001
	Ventilation	-0.008	0.0012	42.441	1	<0.001
	Per cigarette tobacco weight	0.399	0.1572	6.459	1	0.011
CO*	Year	-0.029	0.0099	8.649	1	0.003
	Ventilation	-0.009	0.0011	66.476	1	<0.001
	Per cigarette tobacco weight	0.529	0.1900	7.742	1	0.005
	Filter density	-0.001	0.0003	5.690	1	0.017

*GEE with gamma distribution and log link function.

GEE, generalised estimating equations.

TNCO, tar, nicotine and carbon monoxide.

indicators of human health risk. Otherwise, continued promotion of misperceptions about low-tar cigarettes, coupled with China's limited implementation of other FCTC recommended policies on taxation, smoke-free spaces and health warning labels, will likely prolong the tobacco epidemic.

What this paper adds

The marketing of 'light' and 'low tar' cigarettes in China has resulted in consumer misperceptions of their harmful effects. These lower International Organization for Standardisation (ISO) tar yields can be attributed to design features, such as filter ventilation. We find that across 4 years, to comply with reduced ISO tar ceilings, Chinese cigarette manufacturers increased ventilation to reduce tar levels. Misleading descriptors and ISO yields should be removed from the pack to help address consumer misperceptions.

Acknowledgements The authors thank the China CDC staff and fieldworkers in each city for their assistance with data collection.

Contributors RJO and GTF conceived the study. LMS, BAZ and RVC led data analysis. QL and JY contributed to data collection. All authors contributed to data interpretation and manuscript preparation.

Funding The ITC China Project was supported by grants from the US National Cancer Institute (R01 CA100362, R01 CA125116 and P01 CA138389), the Roswell Park Transdisciplinary Tobacco Use Research Center (P50 CA111236), and the Chinese Center for Disease Control and Prevention.

Competing interests GTF was supported by a Senior Investigator Award from the Ontario Institute for Cancer Research and by a Prevention Scientist Award from the Canadian Cancer Society Research Institute.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Yang G. Marketing 'less harmful, low-tar' cigarettes is a key strategy of the industry to counter tobacco control in China. *Tob Control* 2014;23:167–72.
- Gan Quan WL. Chinese 'low-tar' cigarettes do not deliver lower levels of nicotine and carcinogens. *Tob Control* 2010;374–9.
- Yang J, Hammond D, Driezen P, et al. Health knowledge and perception of risks among Chinese smokers and non-smokers: findings from the Wave 1 ITC China Survey. *Tob Control* 2010;19(Suppl 2):i18–23.
- Elton-Marshall T, Fong GT, Zanna MP, et al. Beliefs about the relative harm of "light" and "low tar" cigarettes: findings from the International Tobacco Control (ITC) China Survey. *Tob Control* 2010;19(Suppl 2):i54–62.
- Global Adult Tobacco Survey: China. Global Tobacco Surveillance System. 2010. http://www.who.int/tobacco/surveillance/en_tfi_china_gats_factsheet_2010.pdf (accessed 9 Jan 2014).
- National Cancer Institute. *Monograph 13: risks associated with smoking cigarettes with low tar machine-measured yields of tar and nicotine*. Rockville: U.S. Department of Health and Human Services; Public Health Service; National Institutes of Health, 2001. http://cancercontrol.cancer.gov/brp/tcrb/monographs/13/m13_complete.pdf (accessed 26 Mar 2013).
- O'Connor RJ, Cummings KM, Giovino GA, et al. How did UK cigarette makers reduce tar to 10 mg or less? *BMJ* 2006;332:302.
- Kozlowski LT, O'Connor RJ. Filter ventilation is a defective design because of lighter taste, bigger puffs, and blocked vents. *Tob Control* 2002;1(Suppl 1):i40–50.
- O'Connor RJ, Li Q, Stephens WE, et al. Cigarettes sold in China: design, emissions and metals. *Tob Control* 2010;19(Suppl 2):i47–53.
- World Health Organization. *WHO Framework Convention on Tobacco Control*. Geneva: WHO Document Production Services, 2005:1–36.
- ITC China Project Report: Findings from the wave 1 to 3 surveys (2006–2009), The International Tobacco Control Policy Evaluation Project. Waterloo: University of Waterloo; December 2012. <http://www.itcproject.org/files/ITC-China-NR-English-web-Dec142012-FINAL.pdf> (accessed 9 Jan 2014).
- Caruso RV, O'Connor RJ. Cigarette design features in low-, middle-, and high-income countries. *J Environ Public Health*, 2012.
- O'Connor RJ, Caruso RV, Borland R, et al. Relationship of cigarette-related perceptions to cigarette design features: findings from the 2009 ITC United States Survey. *Nicotine Tob Res* 2013;15:1943–7.
- O'Connor RJ, Wilkins KJ, Caruso RV, et al. Cigarette characteristics and emission variations across high-, middle-, and low-income countries. *Public Health* 2010;124:667–74.
- Kozlowski LT, Dreschel NA, Stellan SD, et al. An extremely compensatable cigarette by design: documentary evidence on industry awareness and reactions to the Barclay filter design cheating the tar testing system. *Tob Control* 2005;14:64–70.
- Kozlowski LT, Mehta NY, Sweeney CT, et al. Filter ventilation and nicotine content of tobacco in cigarettes from Canada, the United Kingdom, and the United States. *Tob Control* 1998;7:369–75.
- O'Connor RJ, Hammond D, McNeill A, et al. How do different cigarette design features influence the standard tar yields of popular cigarette brands sold in different countries? *Tob Control* 2008;17(Suppl 1):i1–5.
- U.S. Department of Health and Human Services. *The health consequences of smoking: the changing cigarette*. A Report of the Surgeon General. U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health. DHHS Publication No. 81-50156, 1981.
- National Cancer Institute. *The FTC cigarette test method for determining tar, nicotine, and carbon monoxide yields of U.S. cigarettes*. Report of the NCI Expert Committee. Smoking and Tobacco Control Monograph No. 7. U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, NIH Publication No. 96-4028, 1996.
- Warner KE. Tobacco industry response to public health concern: a content analysis of cigarette ads. *Health Educ Q* 1985;12:115–27.
- DOJ final opinion. <http://www.justice.gov/civil/cases/tobacco2/amended%20opinion.pdf>
- Supreme Court of the United States. *Altria Group, INC, et al. v. Good et al.* 2008. <http://www.supremecourt.gov/opinions/08pdf/07-562.pdf> (accessed 23 Jan 2014).
- Kozlowski LT, Yost B, Stine MM, et al. Massachusetts' advertising against light cigarettes appears to change beliefs and behavior. *Am J Prev Med* 2000;18:339–42.
- Li C. *The political mapping of China's tobacco industry and anti-smoking campaign*. China Center Monographs. Washington DC: The John L. Thornton China Center at Brookings, 2012;5.
- Yong HH, Borland R, Cummings KM, et al. Impact of the removal of misleading terms on cigarette pack on smokers' beliefs about 'light/mild' cigarettes: cross-country comparisons. *Addiction* 2011;106:2204–13.