LETTERS TO THE EDITOR

Letters intended for publication should be a maximum of 500 words, 10 references, and one table or figure, and should be sent to the editor at the address given on the inside front cover. Those responding to articles or correspondence published in the journal should be received within six weeks of publication.

Whose standard is it, anyway?

EDITOR—In their recent article in *Tobacco Control* Bialous and Yach¹ create the impression that international standards for the machine smoking of cigarettes were foisted on the smoking public unilaterally by the tobacco industry by its influence on the International Organization for Standardization (ISO) through control of CORESTA (Centre de Co-operation pour les Recherches Scientifiques au Tabac). They also allege, inter alia, that the tobacco industry (1) has, through CORESTA, changed the methodology in order to produce lower smoke yield values to get round the European "tar" ceiling directives, and (2) misleads the public by developing low "tar" cigarettes to cheat the smoking machine, and then makes unjustified health claims about them. ("Standards" are documented agreements containing technical specifications or concise criteria to be used consistently as rules guidelines.)

In their article, Bialous and Yach¹ concentrated predominantly on a few highly selective quotes from internal tobacco company documents. They appear not to have consulted much of the very large volume of scientific literature published on the subject. When this information is taken into account it becomes obvious that the very narrow and restricted literature base of Bialous and Yach's analysis has resulted in them making factual errors, drawing wrong conclusions and writing inaccurate statements on many aspects of the subject.

A review of the published literature on the subject shows clearly that the broad facts are as follows:

(1) Techniques relevant to the machine smoking of cigarettes were developed and refined throughout the 20th century.^{2,3} The first standard was specified by the Federal Trade Commission (FTC), a US federal government agency, in 1966 and first used to test cigarettes in 1967.⁴ The CORESTA recommended method, similar in many respects to that of the FTC, was developed after the FTC standard and was published in 1969.⁵

(2) There were small differences in the details of the smoking machine procedures in the various standard methods developed by the FTC and subsequently CORESTA, ISO and authorities in the UK, Germany, Canada and elsewhere between 1969 and the late 1970s.⁶ These differences resulted in about a 10% difference in the "tar" yield of the same cigarette measured by authorities in Britain and Germany,⁷ for example. By the late 1980s it was recognised that this situation was unacceptable in view of pending European directives which specified "tar" ceilings for all cigarettes sold in member states across Europe from 1993. Consequently, the differences in methodology were

harmonised in a common ISO standard method in 1991, developed following a considerable amount of inter laboratory comparisons of the developing methodology undertaken within CORESTA across 29 laboratories from 15 countries.⁷ This revised standard method is now used in all countries except the USA where the slightly different FTC method still continues to be used, and in Japan where some minor differences are used in their national standard. Changing to the ISO standard in the early 1990s, "tar" yields determined in the UK, for example, decreased by up to 0.5 mg while "tar" yields in Canada, for example, increased by up to 3 mg for some brands.

(3) The purpose of the smoking machine standards is to determine the "tar", nicotine, and carbon monoxide content of cigarette smoke when the cigarette is smoked under precisely defined conditions, and hence to allow a comparison of the yields from different cigarettes. Such yields are not predictive of the yields humans obtain when smoking, nor were they ever expected to be so, since no two smokers smoke exactly the same nor does a smoker smoke a cigarette the same way on every occasion. This purpose has been stated consistently many times, originally by the FTC in 1967,⁴ and subsequently in the scientific literature, published by the tobacco industry and health/regulatory authorities, over the last 35 years, e.g.^{8,9}

(4) Compensation by smokers when switching to a low "tar" cigarette has been discussed in the scientific literature for 40 years. The phenomenon was first published by the tobacco industry and tobacco industry scientists have published books and papers on the subject, e.g.^{10,11} The available evidence, albeit limited, indicates that compensation is partial in the short term (up to a few weeks), and that smokers switching from a higher to a lower "tar" yield cigarette do generally obtain a reduction in smoke delivery.¹¹

(5) Since the 1950s numerous health scientists have advocated that lower "tar" cigarettes should be developed on the grounds that they may represent a less hazardous form of smoking, e.g.^{2,12-15} Health authorities have consistently advised smokers to quit, but for those who choose to continue to smoke that they should smoke lower "tar" cigarettes, e.g.^{8,16,17} The tobacco industry has responded to these health authorities by developing cigarettes with lower "tar" but has also followed public health advice by not advertising lower "tar" cigarettes as safe cigarettes.

RICHARD R BAKER
Research & Development Centre,
British American Tobacco,
Southampton SO15 8TL, UK

- 1 Bialous SA, Yach D. Whose standard is it, anyway? How the tobacco industry determines the International Organization for Standardization (ISO) standards for tobacco and tobacco products. *Tobacco Control* 2001;10:96-104.
- 2 Wynder EL, Hoffmann D. *Tobacco and tobacco smoke: studies in experimental carcinogenesis*. New York: Academic Press, 1967:94-105; 503-516.
- 3 DeBardeleben MZ, Wickham JE, Kuhn WF. The determination of tar and nicotine in cigarette smoke from an historical perspective. *Rec Adv Tob Sci* 1991;17:115-48.
- 4 Federal Trade Commission. Cigarettes: testing for tar and nicotine content. *Federal Register* 1967;32:11178.
- 5 CORESTA Recommended Method No. 10: Machine smoking of cigarettes, determination of crude and dry smoke condensate. *CORESTA Inf Bull* No 1 1969:24-33.

- 6 Rodgman A. FTC "tar" and nicotine in mainstream smoke: a retrospective. *Rec Adv Tob Sci* 1997;23:5-74.
- 7 Thomsen HV. International reference method for the smoking of cigarettes. *Rec Adv Tob Sci* 1992;18:69-94.
- 8 Froggatt P (chairman). *Fourth Report of the Independent Scientific Committee on Smoking and Health*. London: HMSO, 1988.
- 9 CORESTA. Standard smoking methods: a CORESTA perspective. *CORESTA Inf Bull* No. 2 1994:54-7.
- 10 Thornton RE (ed). *Smoking behaviour: physiological and psychological influences*. London: Churchill Livingstone, 1978.
- 11 Scherer G. Smoking behaviour and compensation: a review of the literature. *Psychopharm* 1999;145:1-20.
- 12 Wynder EL, Kopf P, Ziegler H. A study of tobacco carcinogenesis. II. Dose-response studies. *Cancer* 1957;10:1193-200.
- 13 Gori GB. Low-risk cigarettes: a prescription. *Science* 1976;194:1243-6.
- 14 Peto R, Doll R. The control of lung cancer. *New Scientist* 1985;26-30.
- 15 Poswillo D (chairman). *Report of the Scientific Committee on Tobacco and Health*. London: Department of Health, 1998:14.
- 16 World Health Organization. Controlling the smoking epidemic. *Report of the WHO expert committee on smoking control. Technical Report Series No 636*. Geneva: WHO 1979:72.
- 17 World Health Organization. Smoking and its effects on health. *Report of a WHO expert committee. Technical Report Series No 568*. Geneva: WHO, 1975:29-34.

EDITOR—In a recent article, Bialous and Yach¹ attempt "to describe the extent of the tobacco industry involvement in establishing international standards for tobacco and tobacco products." They assert that "it is clear that the tobacco industry, through [CORESTA], play a major role in determining the scientific evidence and suggesting the standards that are eventually developed as international standards." Finally, they conclude that "ISO's tobacco and tobacco products standards are not adequate to guide tobacco products regulatory policies, and no health claims can be made based on [these] standards." Moreover, along the way, these authors seem to suggest some impropriety in CORESTA's involvement in the standards setting process and offer some examples that, they believe, support a contention that is in fact untrue.

CORESTA (Cooperation Centre for Scientific Research Relative to Tobacco) is an organisation devoted entirely to issues related to tobacco science. Those issues range from plant breeding and good agricultural practices to technological aspects of manufacturing and analytical determination of smoke yields. That the majority of worldwide tobacco science expertise resides within the tobacco industry should come as surprise to no one (as would be true for most industries facing technical challenges). That many of these same experts find themselves involved in CORESTA and International Organization for Standardization (ISO) TC-126 activities should, likewise, be of no surprise. Development of technical standards, whether within CORESTA or ISO or elsewhere, without relying on the best available technical expertise would, of course, be irrational.

Regarding the suggestion of impropriety, Bialous and Yach outline three areas they believe support their case. First, they claim "[ISO] standards are approved as recommended by CORESTA, with limited opportunity for significant amendments". Offered as an example is an excerpt of CORESTA minutes regarding ISO/DIS 11454, stating that the DIS (Draft

International Standard) will be published as an international standard with no changes other than editorial. What Bialous and Yach apparently fail to appreciate is that the ISO approval process leading up to the DIS stage involved a development period of at least four years and multiple balloting stages (opportunities) for significant amendments. For example, balloting at the previous committee draft stage (ISO/TC 126 N 537, July 1995) generated nine pages of comments from 12 countries. The 1997 CORESTA minutes reflect only that at the DIS stage in the overall ISO process, no changes had been requested by ISO members other than those of an editorial nature.

Secondly, Bialous and Yach made a compound assertion, that "CORESTA works with ISO directly or that CORESTA works through one of ISO's member bodies". CORESTA does have a liaison member status with ISO, but does not work with any of ISO's member bodies.

Lastly, Bialous and Yach assert "CORESTA resists any interference with its proposed standards, and make efforts to keep overall control of the situation and the outcomes of ISO meetings". Offered as support is a matter concerning updates to the ISO smoking methods. Again, an egregious misinterpretation has resulted. A CORESTA working group and the British Standards Institution (BSI) independently prepared editorial commentary on similar issues within the text of the ISO smoking methods. Wishing to defer to the ISO process, CORESTA postponed an update to the existing CORESTA methods, instead wishing to wait for ISO to finish their deliberations.

On these claims of impropriety, Bialous and Yach offer examples that do not support their contention. Rather to the contrary, these examples serve as testament to the propriety of the CORESTA-ISO relationship.

We look forward to a continued relationship with all parties with an interest in the business of technical standardisation related to tobacco and tobacco products.

FRANÇOIS JACOB
Secretary General, CORESTA
11 rue du Quatre Septembre,
75002 Paris,
France
coresta.foj@wanadoo.fr

- 1 Bialous SA, Yach D. Whose standard is it anyway. How the tobacco industry determines the International Organization for Standardization (ISO) standards for tobacco and tobacco products. *Tobacco Control* 2001; 10:96-104.

Author's reply

EDITOR—Mr Jacob and Dr Baker's criticisms of our paper¹ mostly indicate an incomplete reading of it. We believe our paper reached its stated objective and showed that "the tobacco industry, through [CORESTA], play a major role in determining the scientific evidence and suggesting the [international] standards." Our conclusion that "ISO's tobacco and tobacco products standards are not adequate to guide tobacco products regulatory policies, and no health claims can be made based on [these] standards" derives both from the evidence in the paper and evidence from studies and international reviews.²⁻³ Such standards are useful to rank cigarettes' nicotine and tar yields according to machine measurements, they do not reflect human smoking, and measurement methods can be modified to provide different readings

without the need for product modification, a point also made by Dr Baker.

Recent literature and health authorities' recommendations regarding "lower" tar and nicotine cigarettes show that switching to lower yields is no longer perceived as an effective public health measure to address tobacco related morbidity and mortality.³⁻⁵ Several countries and the European Union have already banned or are moving to ban the utilisation of labels such as "mild" and "lights" based on scientific and tobacco industry document evidence that these mislead the public into thinking these cigarettes are safer.⁶⁻¹⁰

Indeed, a 1977 British America Tobacco (BAT) document stated: "All work in this area should be directed towards providing consumer reassurance about cigarettes and the smoking habit. This can be provided in different ways, for example, by claiming low deliveries, by the perception of low deliveries, and by the perception of "mildness". Furthermore, advertising for low delivery or traditional brands should be constructed in ways so as not to provoke anxiety about health, but to alleviate it, and enable the smoker to feel assured about the habit and confident in maintaining it over time."¹¹

We do not suggest "impropriety in CORESTA's involvement" since, as per the ISO definition, the tobacco industry is entitled to participate in the process. However, we question the adequacy of having tobacco interests being the sole provider of scientific evidence in the area of tobacco products standardisation methods. We question CORESTA's involvement as much as the lack of involvement from other interested parties (for example, health and consumer groups) as well as whether or not the ISO is fulfilling its aim of protecting the health and safety of consumers of tobacco products.

Mr Jacob's assertion that the "majority of worldwide tobacco science expertise resides within the tobacco industry should come as surprise to no one" is correct. However, recent litigation has shown that this expertise has not been used to benefit the health and safety of the consumers of tobacco products.¹²⁻¹⁴ It is also no surprise that many of the tobacco industry experts "find themselves involved in CORESTA and ISO TC-126 activities", but it is unacceptable that these experts be the only ones participating in these activities.

As for the three specific areas of criticism in Mr Jacob's letter:

(1) From the description of the ISO standards approval process,¹⁵ the majority of work is done at the Technical Committee (TC) level, and final approval of a DIS (Draft International Standard) is by the TC as well. In the case of TC 126, with a majority of members representing the tobacco industry, and CORESTA being the organisation conducting the work on the proposed standards, amendments are referred back to the TC and to CORESTA. In the example offered, ISO/DIS 11454, reference 34 in our paper¹⁶ describes some of these comments and how they represent the tobacco industry's perspective.

(2) The assertion "CORESTA works with ISO either directly or through one of ISO's members bodies . . ." emphasises the great overlap among the people (and interests) who are members of CORESTA, TC 126 and those involved in tobacco issues at member bodies such as British Standards Institution

(BSI) and American National Standards Institution (ANSI). Although no official relation between CORESTA and ISO's member bodies exists, it is clear that efforts are often agreed upon. In the example provided, at a CORESTA Scientific Commission meeting¹⁷ a method for environmental tobacco smoke (ETS) determination was sent to ISO via ANSI. (An unquoted example, from reference 45,¹⁸ states that CORESTA had prepared a draft standard on ambient air to be circulated at BSI.)

(3) We saw no evidence that CORESTA accepts outsiders' input in preparing standards forwarded to ISO, but it is clear that it wants to maintain "CORESTA methods and ISO standards . . . close or identical."¹⁹ The following quote from the same document¹⁹ addresses how CORESTA planned to deal outside participation (by participation in the validation stage, not in the development, of a measurement method).

We appreciate the opportunity to address these comments, and Mr Jacob's offer for a continued dialogue in the area of standardisation of tobacco products.

STELLA AGUINAGA BIALOUS
DEREK YACH
Tobacco Free Initiative,
World Health Organization,
Geneva, Switzerland
stella@bialous.com

- 1 Bialous SA, Yach D. Whose standard is it, anyway? How the tobacco industry determines the International Organization for Standardization (ISO) standards for tobacco and tobacco products. *Tobacco Control* 2001;10:96-104.
- 2 Stratton K, Shetty P, Wallace R, et al, eds. *Clearing the smoke: the science base for tobacco harm reduction*. Committee to Assess the Science Base for Tobacco Harm Reduction, Board on Health Promotion and Disease Prevention. Washington DC: Institute of Medicine, February 2001:514 pages.
- 3 Tobacco Free Initiative/World Health Organization. *Monograph: Advancing knowledge on regulating tobacco products*. Geneva: WHO, May 2001:24 pages. URL: <http://tobacco.who.int/en/fctc/Oslo/OsloMonograph.pdf>
- 4 Jarvis MJ, Boreham R, Primatesta P, et al. Nicotine yield from machine-smoked cigarettes and nicotine intakes in smokers: evidence from a representative population survey. *J Natl Cancer Inst* 2001;93:134-8.
- 5 Bates C, McNeill A, Jarvis M, et al. The future of tobacco product regulation and labelling in Europe: implications for the forthcoming European Union directive. *Tobacco Control* 1999;8:225-35.
- 6 Jarvis M, Bates C. *Why low tar cigarettes don't work and how the tobacco industry has fooled the smoking public*. 18 March 1999 ed. London: Action on Smoking and Health UK. URL: <http://www.ash.org.uk/html/regulation/html/big-one.html>
- 7 Commission of the European Communities. *Directive of the European Parliament and of the Council on the approximation of the laws, regulation and administrative provision of the Member States concerning the manufacture, presentation and sale of tobacco products*. Brussels: Commission of the European Communities, 1999:26.
- 8 Health and Consumer Protection Directorate General/The European Commission. *Tough EU rules on manufacture, presentation and sale of tobacco products agreed*. Press Release, Brussels, 15 May 2001. URL: http://europa.eu.int/comm/dgs/health_consumer/library/press/press137_en.html
- 9 Physicians for a Smoke-Free Canada & Non-Smokers' Rights Association. *Misleading cigarette marketing: the 'light' and 'mild' deception*. URL: <http://www.nsr-aadnf.ca/english/lights/index.html> Accessed 19 August 2001.
- 10 Rock A. Speaking notes for Allan Rock, Minister of Health, Health Canada. Canadian Medical Association's 134th Annual General Meeting. Quebec City, Quebec. Monday, 13 August 2001. URL: <http://www.hcsc.gc.ca/english/archives/speeches/13aug2001mine.htm>
- 11 Short PL. Smoking and health item 7: the effect on marketing. 14 April 1977. BAT Co Minnesota Trial Exhibit 10,585.

- 12 Committee of Experts on Tobacco Industry Documents. *Tobacco company strategies to undermine tobacco control activities at the World Health Organisation. Report of the Committee of Experts on Tobacco Industry Documents*. July 2000, Geneva: WHO:260 pages. URL: <http://tobacco.who.int/en/fctc/Oslo/Monograph.pdf>
- 13 Glantz SA, Slade J, Bero LA, et al. *The cigarette papers*. Berkeley, California: University of California Press, 1996. URL: <http://www.library.ucsf.edu/tobacco/cigpapers/book/contents.html>
- 14 Kessler DA. *A question of intent: a great American battle with a deadly industry*. New York: Public Affairs, 2001.
- 15 <http://www.iso.ch/iso/en/stdsdevelopment/whowhenhow/proc/proc.html>
- 16 ISO/TC 126 N 535. 1995 Draft Report of the Nineteenth Plenary Meeting held in Cape Town (South Africa) on 22 and 23 February 1995. April 1995, Philip Morris: Report. Accessed June 2000, Bates 51749 6783/6810. URL: www.rjrtdocs.com
- 17 Reif H. 1998 Memorandum to Richard Carchman: monthly activities 1st through 31st May, 1998. Philip Morris. Accessed February 2000, Bates 2060565811/5814. URL: www.pmdocs.com
- 18 Scientific Liaison Working Group of the EIG. 1994 Draft minutes of the 28th meeting. 20 June 1994. London: EIG. Minutes. Accessed March 2000, Bates 2026219657/9661, URL: www.pmdocs.com
- 19 CORESTA Outgoing Scientific Commission. 1996 Minutes of the Outgoing Scientific Commission Meeting, 3 November 1996, Japan. CORESTA: Accessed February 2000, Bates 51880 5267/5273. URL: www.rjrtdocs.com

Desire to stop smoking among intermittent and daily smokers: a population-based study

EDITOR—An important fraction of all smokers are intermittent, non-daily smokers,^{1,3} and the proportion of intermittent smokers may even be rising.^{1,4} Intermittent smokers are younger and have a higher educational and occupational status than daily smokers.^{1,4} Some intermittent smokers are either in the uptake phase of smoking, or are preparing for smoking cessation. However, intermittent smoking can also be a long term behaviour.^{1,2} Intermittent smokers are more likely than daily smokers to have a strong intention to quit

smoking. They are also more likely to actively start the process of smoking cessation.^{1,5} Intermittent smokers probably also suffer less severe withdrawal symptoms during cessation attempts than do daily smokers and, therefore, have a greater potential for success.¹ Intermittent smokers perceive quitting as not being very difficult.⁶ However, there are no studies concerning the prevalence of the desire to stop smoking among intermittent compared to daily smokers.

The public health survey in Malmö 1994 is a cross sectional study. A total of 5600 individuals born in 1913, 1923, 1933, 1943, 1953, 1963, 1968, and 1973 were randomly selected from the general Malmö population and interviewed by a postal questionnaire in the spring of 1994. In each age group, 700 participants (350 men and 350 women) were interviewed. The participation rate was 71%. The desire to stop smoking item, "Do you want to stop smoking?", had two alternative answers, "yes" and "no", and the item was dichotomised accordingly. The *smoking* item contained four alternatives: never smoked, stopped smoking, daily smoker, and intermittent (non-daily) smoker. The sex differences in daily smoking, intermittent smoking, never smoked, and stopped smoking were calculated using *t* tests (results only presented in text). The differences in proportions of daily and intermittent smokers that report a desire to stop smoking were also calculated with *t* tests (results only presented in text). The proportions of daily and intermittent smokers that express desire to stop smoking were calculated separately using logistic regression in order to analyse associations between sociodemographic variables and desire to stop smoking (table 1). The SPSS software package was used.

A total of 56.4% of all male and 59.9% of all female smokers (both daily and intermittent) had expressed a desire to stop smoking ($p < 0.001$). The proportion of daily smokers was 24.5% among men and 23.7% among women ($p < 0.001$). Men were

intermittent smokers to a slightly higher extent (8.7%) than women (6.3%) ($p < 0.001$). Only 34.9% of the men had never smoked, while 50.3% of the women had never smoked ($p < 0.001$). In contrast, 31.9% of the men had stopped smoking compared to only 19.8% of the women ($p < 0.001$). A higher proportion (67.7%) of all daily smokers expressed a desire to stop smoking, compared to only 32.3% of all intermittent smokers ($p < 0.001$). Table 1 shows that the desire to stop smoking was highest among daily smokers born in 1963, odds ratio 1.9 (95% confidence interval (CI) 1.04 to 3.6), compared to daily smokers born in 1973. The desire to stop smoking then decreased significantly with age. No significant differences according to age in the desire to stop smoking were seen among the intermittent smokers. Daily smokers born in Sweden and abroad had expressed a desire to stop smoking to the same extent. In contrast, intermittent smokers born abroad had expressed a desire to stop smoking to a much higher extent than the group born in Sweden. No significant differences between the educational status groups in desire to stop smoking were observed for either daily or intermittent smokers.

Non-participation is not likely to have produced serious selection bias. There was a good concordance between the data and register census data concerning foreign origin and the general educational level of the Malmö population.⁷ There is also the possibility of bias from self reported data. Although self reporting of desire to stop smoking is the only conceivable way to assess desire to stop smoking, it is subjective and may even reflect a person's general perception of health. However, this fact would probably not bias the systematic differences between intermittent and daily smokers reported in this study. The proportion of all smokers that express a desire to stop smoking is very similar to the results of other studies.⁸⁻¹⁰

The results may seem surprising, given the fact that other studies have reported that some intermittent smokers may be former regular smokers in the process of smoking cessation, and that these studies also have reported a stronger intention to quit and a greater likelihood of having recently attempted to quit among intermittent smokers.^{1,5} However, the prevalence of pre-intentional desire to stop smoking seems to be much lower among intermittent than among daily smokers, and this finding strengthens the conclusion of other studies that a substantial fraction of all intermittent smokers are long term intermittent smokers and others are in the uptake phase of smoking. The conclusion that a higher proportion of daily smokers than intermittent smokers report a desire to stop smoking has yet another dimension. No less than 63.7% of all male and 65.4% of all female daily smokers that reported a desire to stop smoking also reported a desire to get help to stop smoking (yes/no). In contrast, the proportion that expressed a desire to get help to stop smoking was only 20.4% of male and 14.3% of female intermittent smokers that expressed a desire to stop smoking ($p < 0.001$ when comparing daily and intermittent smokers among both men and women).

Pomerleau and Pomerleau have stressed that nicotine has a variety of effects that may be directly reinforcing, even in the absence of dependence.¹¹ Specific benefits of smoking

Table 1 Crude odds ratios (OR) and 95% confidence intervals of desire to stop smoking among daily smokers and intermittent smokers according to sociodemographic and snuff consumption characteristics. The public health survey in Malmö 1994.

	Daily smoker			Intermittent smokers		
	n	%	OR (95% CI)	n	%	OR (95% CI)
<i>Year of birth</i>						
1973	85	62.4	1.0	55	43.6	1.0
1968	107	65.4	1.1 (0.6 to 2.1)	59	42.4	1.0 (0.5 to 2.0)
1963	113	76.1	1.9 (1.04 to 3.6)	51	37.2	0.8 (0.4 to 1.7)
1953	152	73.7	1.7 (0.96 to 3.0)	32	43.8	1.0 (0.4 to 2.4)
1943	150	62.7	1.0 (0.6 to 1.8)	21	19.0	0.3 (0.1 to 1.02)
1933	110	64.5	1.1 (0.6 to 2.0)	13	30.8	0.6 (0.2 to 2.1)
1923	101	50.5	0.6 (0.3 to 1.1)	9	33.3	0.6 (0.2 to 2.9)
1913	49	28.6	0.2 (0.1 to 0.5)	7	57.1	1.7 (0.4 to 8.4)
Missing	13			25		
<i>Country of origin</i>						
Sweden	590	62.5	1.0	186	34.4	1.0
Other	277	65.7	1.1 (0.9 to 1.5)	61	52.5	2.1 (1.2 to 3.8)
Missing	13			25		
<i>Education</i>						
> 12 years	201	65.2	1.0	101	32.7	1.0
10–12 years	168	70.2	1.3 (0.8 to 2.0)	76	43.4	1.6 (0.9 to 2.9)
< 9 years	428	61.0	0.8 (0.6 to 1.2)	60	40.0	1.4 (0.7 to 2.7)
Others	64	56.2	0.7 (0.4 to 1.2)	7	42.9	1.5 (0.3 to 7.3)
Missing	19			28		
<i>Snuff user</i>						
No	701	63.1	1.0	175	40.6	1.0
Yes	35	74.3	1.7 (0.8 to 3.7)	53	30.2	0.6 (0.3 to 1.2)
Missing	144			44		
<i>Total</i>	880			272		

have been documented in cognitive and psychomotor performance.¹² The smoking of intermittent smokers may be motivated by these effects.

The results further support the notion that intermittent smokers are a specific group of smokers with smoking cessation characteristics that differ from the characteristics of daily smokers.

MARTIN LINDSTRÖM

Department of Community Health,
University Hospital MAS,
Lund University,
S 205 02 Malmö,
Sweden
martin.lindstrom@smi.mas.lu.se

- 1 Hennrikus DJ, Jeffrey RW, Lando HA. Occasional smoking in a Minnesota working population. *Am J Public Health* 1996;86:1260-6.
- 2 Evans NJ, Gilpin E, Pierce JP, et al. Occasional smoking among adults: evidence from the California tobacco survey. *Tobacco Control* 1992;1:169-75.
- 3 Cigarette smoking among adults—United States, 1992, and changes in the definition of current cigarette smoking. *MMWR Morb Mortal Wkly Rep* 1994;43:342-6.
- 4 Husten CG, McCarty MC, Giovino GA, et al. Intermittent smokers: a descriptive analysis of persons who have never smoked daily. *Am J Public Health* 1998;88:86-9.
- 5 Owen N, Kent P, Wakefield M, et al. Low-rate-smokers. *Prev Med* 1995;24:80-4.
- 6 Shiffman S. Refining models of dependence: variation across persons and situations. *Br J Addict* 1991;86:611-15.
- 7 City of Malmö. *Malmö statistical yearbook 1994*. Malmö: City of Malmö, 1994.
- 8 National Opinion Poll Omnibus Services. *Smoking habits 1991*. London: NOP, 1992.
- 9 Prochaska JO, Di Clemente CC. Self change process, self efficacy and decisional balance across five stages of smoking cessation. *Prog Clin Biol Res* 1984;156:131-40.
- 10 Viscusi WK. *Smoking: making the risky decision*. New York: Oxford University Press, 1992.
- 11 Pomerleau OF, Pomerleau CS. Neuroregulators and the reinforcement of smoking: towards a behavioral explanation. *Neurosci Biobehav Rev* 1984;8:503-13.
- 12 Wesnes K, Warburton DM. Smoking, nicotine, and human performance. *Pharmacol Ther* 1983;21:189-208.

Smoking among Japanese nursing students: a nationwide survey

EDITOR—In some developed countries including Japan, smoking prevalence among nursing students tends to be the same or higher than that of the general female population of the same age group.¹⁻⁶ In Japan, an increase in the prevalence of smoking among women in their 20s was recently reported,⁷ and this trend is assumed to be reflected in the smoking prevalence of nursing students, where young women are over represented.

To obtain data on smoking prevalence of nursing students, a nationwide survey was conducted among the students of nursing (three year programme), public health nursing (PHN), and midwifery schools. Students of PHN and midwifery are qualified as nurses, and involved in one year training to acquire qualification as PHN and midwives, respectively. The survey was conducted in October 2000 using self reported questionnaires.

In 2000, there were 465 three year nursing schools (total number of students: 66 430,) 66 PHN schools (1697 students), and 73 midwifery schools (1420 students) in Japan. Among these, 27 nursing schools, 17 PHN schools, and 16 midwifery schools were selected at random, and the survey was carried out on all students in the selected schools. Between selected schools and non-selected schools, little difference was

observed with respect to their geographical distribution and student volume size.

Each subject from the selected schools filled in the questionnaire, put it into an envelope, sealed and handed it to the person in charge. The questionnaire included the items of a previous survey on smoking behaviour among nurses,⁸ and eight items related to the nicotine dependency scale of Fagerstrom.⁹

The return rates were 93% (3866/4169) for the nursing schools, 91% (539/592) for the PHN schools, and 95% (325/343) for the midwifery schools. After excluding incompletely answered questionnaires, 3762, 530, and 303 responses were analysed, respectively.

The prevalence of smoking among women was 25% in the nursing schools, 13% in the PHN schools, and 22% in the midwifery schools. In the nursing schools, the prevalence of smoking increased as the grade advanced. In the third year, the prevalence of smoking was 31%, higher than that among the general population in their 20s (23%).⁷ As to male students in third year, the prevalence of smoking was nearly the same as that of the general male population in their 20s (60%).⁷ Furthermore, the nicotine dependency among female daily smokers in the nursing schools was higher than that in the PHN schools or midwifery schools. Therefore, anti-smoking education in nursing schools is urgently needed. In this survey, smoking prevalence was lower among students in the PHN and midwifery schools. The difference occurs because those who had already qualified as nurses and wished to continue studying to acquire another qualification were less likely to smoke than those who were not in the same career level. It is therefore suggested that the prevalence of smoking among less educationally motivated students is lower. Adriaanse and colleagues¹⁰ reported that nurses who were motivated in their jobs had a tendency not to smoke, which is consistent with our results although our subjects were nursing students.

TAKASHI OHIDA
AIKO SAKURAI
AMM KAMAL
TOMOFUMI SONE
SHINJI TAKEMURA
FUJIKO FUKUSHIMA*

Department of Public Health Administration,
National Institute of Public Health
Department of Public Health Nursing*,
Tokyo, Japan
tohida@niph.go.jp

This survey was supported in part by a Health Science Grant from the Ministry of Health and Welfare, Japan (currently called Ministry of Health, Labor, and Welfare, Japan).

- 1 Casey FS, Haughey BP, Dittmar SS, et al. Smoking practices among nursing students: a comparison of two studies. *J Nurs Educ* 1989; 28:397-401.
- 2 Carmichael A, Cockcroft A. Survey of student nurses' smoking habits in a London teaching hospital. *Res Med* 1990;84:277-82.
- 3 O'Connor AM, Harrison M. Survey of smoking prevalence among Canadian nursing students and registered nurses. *Can J Public Health* 1992;83:417-21.
- 4 Boccoli E, Federici A, Melani AS, et al. Results of a questionnaire about nurse students' smoking habits and knowledge in an Italian teaching school of nursing. *Eur J Epidemiol* 1996;12:1-3.
- 5 Ohida T, Osaki Y, Okada O, et al. Factors related to smoking habits of students and newly employed nurses [in Japanese with English abstract]. *Jpn J School Health* 1998; 40:332-40.
- 6 Okada K. Smoking behavior among student nurses in Japan. The Japan Academy of Nursing Science, Kobe: *Second International Nursing Research Conference in Kobe*, 1995:300.

- 7 Anon. Nationwide cigarette smoking survey in 1999 [in Japanese]. Tokyo: Japan Tobacco Industry Inc, 1999.
- 8 Ohida T, Osaki Y, Kobayashi U, et al. Smoking prevalence of female nurses in the national hospital of Japan. *Tobacco Control* 1999;8:192-5.
- 9 Fagerstrom KO. Measuring degree of physical dependence on tobacco smoking with reference to individualization of treatment. *Addictive Behaviors* 1978;3:235-41.
- 10 Adriaanse H, Van Reek J, Zandbelt L, et al. Nurses' smoking worldwide: a review of 73 survey of nurses' tobacco consumption in 21 countries in the period 1959-1988. *Int J Nurs Stud* 1991;28:361-75.

On-line teen smoking cessation: what's porn got to do with it?

EDITOR—As part of our research on adolescent smoking, we conducted a search of the internet for on-line support for teen smoking cessation. We searched google.com using the words *teen quit smoking* (without quotation marks) which resulted in hundreds of potential links. In order to narrow the search to more specific tobacco-related sites, we used an advanced exact-phrase search of the key words "*teen quit smoking*" (with quotation marks) on the same search engine. To our amazement, seven out of the top 20 sites (35%) were teen pornography sites. The phrase "*teen quit smoking*" was deliberately placed among the descriptors for each of these seven pornography links. On further review of several of these sites, we found no smoking cessation material or links to actual cessation sites. Although we are unsure why this phrase would be placed among the descriptors for pornography sites, it raises concern about a teenager's ability to find legitimate on-line cessation support. This unexpected placement of "*teen quit smoking*" potentially encourages teenagers to access on-line pornography, an activity that certainly would be discouraged by many proponents of teen smoking cessation. Fortunately, the same search strategies did not yield the same results with other popular internet search engines. Health educators need to be aware of this potential problem, as more and more teenagers are encouraged to access the internet for smoking cessation support and other health related information.

SEAN P ELLIOTT
CHRISTINE C EDWARDS
SUSAN I WOODRUFF
TERRY L CONWAY
Graduate School of Public Health,
San Diego State University,
9245 Sky Park Ct, Suite 120,
San Diego, CA 92123,
USA
swoodruff@mail.sdsu.edu

Smoking in movies in 2000 exceeded rates in the 1960s

EDITOR—Smoking in movies has been linked to increased smoking among teens.¹⁻⁵ We have previously published data from 1960 through 1997 that shows that smoking fell from the 1960s through the 1980s, then increased during the 1990s.^{6,7} We have used similar methods (analysis of a random sample of five of the top 20 grossing US films each year) to extend the data set through 2000 (fig 1).

We conducted a regression analysis of these data by filling a quadratic equation in time to the amount of tobacco use per hour. The equation, smoke/hour = 801 - 0.405 (± 0.19, p = 0.04) year + 0.0124 (± 0.0044, p = 0.006) year², confirms that, after falling during the early part of this period, smoking

Figure 1 Frequency of tobacco use (events per hour) in a random sample of top grossing films from 1960 through 2000. The films were watched in five minute intervals and each use of tobacco in a given interval was counted as a single event. The total number of events was then divided by the duration of the film.^{7,8}

is now increasing significantly. Based on this regression equation, on average there were 7.3 instances of tobacco use per hour in films in 1960 compared with 10.9 in 2000.

The messages continue to reflect tobacco industry marketing themes of glamour, rebelliousness, and independence, rather than the realities of addiction, suffering, and death.

KAREN KACIRK
STANTON A GLANTZ

University of California, San Francisco
San Francisco, CA 94143-0130,
USA

kacirkk@cvti.ucsf.edu, glantz@medicine.ucsf.edu

This work was supported by National Cancer Institute Grant CA-61021.

- 1 Chapman S, Davis R. Tobacco use in films: is it a problem? *Tobacco Control* 1997;6:269-71.
- 2 Pechmann C, Shih C. Smoking in movies and antismoking advertisements before movies: effects on youth. *J Marketing* 1999;63:1-13.
- 3 Distefan J, Gilpin E, Sargent J, et al. Do movie stars encourage adolescents to start smoking? Evidence from California. *Prev Med* 1999;28:1-11.
- 4 Sargent JD, Beech ML, Daton MA, et al. Is viewing tobacco use in motion pictures associated with trying smoking? A cross-sectional study. *BMJ* (in press).
- 5 Sargent J, Tickle J, Beach M, et al. Brand appearances in contemporary cinema films and contribution to global marketing of cigarettes. *Lancet* 2001;357:29-32.
- 6 Hazan A, Lipton H, Glantz S. Popular films do not reflect current tobacco use. *Am J Public Health* 1994;84:998-1000.
- 7 Teti T, Glantz S. Smoking in movies remained high in 1997. *Tobacco Control* 1998;7:441-2.

BOOK

Book reviews and books of interest to "Tobacco Control" should be sent to the editor at the address given on the inside front cover.

Smoke in their eyes

Smoke in their eyes: lessons in movement leadership from the tobacco wars. Michael Pertschuk. Nashville, Tennessee: Vanderbilt University Press, 2001.

There have been memorable dates, both glorious and infamous, that have defined aims and charted progress and setbacks for

the US tobacco control movement. The most notable of these events have had global repercussions.

On 15 December 1953, the heads of four major US tobacco firms met in New York City's Park Plaza Hotel, where they launched the Tobacco Industry Research Council and hammered out the seminal text of a nationwide newspaper ad, "A frank statement to cigarette smokers". On 11 January 1964, US Surgeon General Luther Terry released the *Report of the Advisory Committee on Smoking and Health*, concluding "it is the judgment of the Committee that cigarette smoking contributes substantially to mortality from certain specific diseases and to the overall death rate". At the Waxman Hearings on 14 April 1994, the "seven dwarfs", chief executive officers (CEOs) of the top seven US tobacco companies, were photographed for posterity as they prepared to declare that "nicotine is not addictive".

At least one more date marks the US tobacco control calendar, but it evokes no signal image, conjures no immortal quote. And yet, on 3 April 1997, at the Sheraton Hotel in Crystal City, Virginia, an extraordinary meeting did take place. That Thursday afternoon, Geoffrey Bible and Steve Goldstone, CEOs of Philip Morris and RJ Reynolds, respectively, met in secret with trial lawyers and state attorneys general, hoping to hammer out a settlement of litigation pending against the tobacco industry. Along with the CEOs, the lawyers, the attorneys general, and their minions, there was one more participant at that meeting: Matthew Myers, vice president and general counsel of the National Center for Tobacco-Free Kids.

In *Smoke in their eyes: lessons in movement leadership from the tobacco wars*, Michael Pertschuk describes the political path that led Myers to that Virginia hotel and chronicles what happened in the meeting's wake. Pertschuk, former head of the US Federal Trade Commission, founder and co-director of the Advocacy Institute, and longtime combatant in the tobacco wars, interviews many key players and makes innovative use of email records to "set the record straight" on the

role of various advocates during the tumultuous debate over the 1997 "global" settlement and the 1998 McCain bill. However, as the author repeatedly makes clear, the book is as much an impassioned defence of Myers as it is an analysis of the colossal, scarring failures of tobacco control advocates during that time.

At its core, *Smoke in their eyes* pits Myers against Stanton Glantz, University of California professor of medicine and lead author of *The cigarette papers*. For Pertschuk, the plausible, if arguable benefits of the McCain bill could have been realised if not for the schism cleaving former allies into hostile camps. According to Pertschuk, great public health gains could have been realised had Glantz and his zealous followers not framed the debate to suit their purposes.

As a behind-the-scenes look at the personalities and polemics of both advocacy groups and political agencies, the book is a rousing success. Though there are few felicitous literary passages, Pertschuk has obtained detailed accounts from former Surgeon General C Everett Koop, former head of the Food and Drug Administration David Kessler, and other principal players, with the glaring exception of Glantz. The book's central failing, however, is Pertschuk's unwillingness or inability to focus on Myers's secret, unilateral decision to attend that first Virginia meeting.

Myers was like Caesar crossing the Rubicon, with just a slight difference or two. Firstly, the general neglected to tell the troops he'd crowned himself emperor. Then, he realised he didn't know the way to the river's edge. Those failings are paramount. Myers's good intentions should not be doubted, but he paved the path to acrimonious, rancorous debate. The Center for Tobacco-Free Kids was not a well established entity in 1997 and many former allies felt betrayed by Myers' "lone ranger" tactics. Once turned off, they could not easily be convinced to follow Myers anywhere, as demonstrated by the caustic, pitched battles between the rival ENACT and Save Lives, Not Tobacco coalitions.

The what-could-have-beens of the McCain bill are still being debated. The USA is again playing a negative role on the global tobacco stage, this time with respect to the Framework Convention on Tobacco Control. What is certain is that the tobacco industry knew what it wanted back in 1997 and still knows what it wants today.

Philosopher Isaiah Berlin famously borrowed the dictum of the Greek poet Archilochus, who wrote: "The fox knows many things, but the hedgehog knows one big thing." The tobacco industry is a huge, knowing hedgehog. Michael Pertschuk's insight and intellect help explain how the tobacco control movement has outfoxed itself lately, but his Manichean dichotomy of Myers-good, Glantz-bad does the movement a disservice. The hedgehog rolls along. The fox needs a new game plan.

STAN SHATENSTEIN
Editor, *GLOBALink Tobacco News*

5492-B Trans Island,
Montreal, Quebec,
Canada H3W 3A8
shatensteins@sympatico.ca