RESEARCH PAPER

Health consequences of reduced daily cigarette consumption

Aage Tverdal, Kjell Bjartveit

Tobacco Control 2006;15:472-480. doi: 10.1136/tc.2006.016246

Objective: To determine the risk of dying from specified smoking-related diseases and from any cause in heavy smoking men and women (≥ 15 cigarettes/day), who reduced their daily cigarette consumption by $>50\%$.

Design: A prospective cohort study.

Setting: Three counties in Norway.

Participants: 24 959 men and 26 251 women, aged 20–49 years, screened for risk factors of cardiovascular disease in the mid-1970s, screened again after 3–13 years, and followed up throughout 2003.

Outcomes: Absolute mortality and relative risks adjusted for confounding variables, of dying from all causes, cardiovascular disease, ischaemic heart disease, all smoking-related cancer and lung cancer.

Results: With sustained heavy smokers as reference, the smokers of both sexes who reduced their daily consumption (reducers) had the following adjusted relative risks (95% confidence interval (CI)): of dying from any cause, 1.02 (0.84 to 1.22); cardiovascular disease, 1.02 (0.75 to 1.39); ischaemic heart disease, 0.96 (0.65 to 1.41); smoking-related cancer, 0.86 (0.57 to 1.29); and lung cancer, 0.66 (0.36 to 1.21). The difference in cigarette consumption between two examinations was not a significant predictor of death from any of the causes. A follow-up from a third screening of the subgroup who were reducers at both second and third examinations (sustained reducers) did not have a lower risk than those who were heavy smokers at all three examinations.

Conclusions: Long-term follow-up provides no evidence that heavy smokers who cut down their daily cigarette consumption by $>50\%$ reduce their risk of premature death significantly. In health education and patient counselling, it may give people false expectations to advise that reduction in consumption is associated with reduction in harm.

See end of article for authors' affiliations

Correspondence to:
K Bjartveit, Fridtjof
Nansens vei 24 B, N-0369
Oslo, Norway; kjell.
bjartveit@chello.no

Received 14 February 2006
Accepted 16 August 2006

Does reduced daily cigarette consumption lead to lower mortality from the serious health consequences of cigarette smoking? In other words, does a reduction in consumption bring about reduction in harm?

Numerous population studies have given ample evidence that quitting smoking entirely results in a marked reduction in the ill effects of smoking. Up to now, however, only one large prospective study has explored the long-term effects of unassisted reduced smoking. Godtfredsen *et al*^{1–4} have pioneered in this field by following up a population of nearly 20 000 men and women living in Copenhagen, Denmark (the Copenhagen Centre for Prospective Population Studies). After mean observation periods ranging from 13.8 to 18 years, they published their results in a series of articles.

The aim of this paper is to determine the risk in heavy smokers who reduced their cigarette consumption by at least 50%, named "reducers", compared with those who continued as heavy smokers. We report on a Norwegian population of 51 210 men and women, aged 20–49 years, who were examined in the mid-1970s for cardiovascular disease risk factors and were examined again during the next 3–13 years at least once. These people were followed up throughout 2003 for deaths from serious smoking-related diseases and from all causes. The mean observation period is 21.2 years, maximum 27 years. A subgroup attended three screenings. This gave us the possibility to compare those who were reducers at the two last screenings (sustained reducers) with those who were heavy smokers at all three screenings (sustained heavy smokers).

METHODS

Initial screening

Between 1974 and 1978, screening examinations for cardiovascular disease were undertaken in three Norwegian

counties with a mainly rural settlement (Finnmark, Sogn og Fjordane, and Oppland). All male and female residents aged 35–49 years were invited, and so was a 10% random sample of residents aged 20–34 years. The attendance was 88%.⁵

An identical protocol was applied for the screening programme in the three counties concerned. It included a questionnaire related to well-known risk factors of cardiovascular disease. Height, weight and blood pressure were measured, and a non-fasting blood sample was drawn and serum analysed at the same laboratory for total cholesterol, triglycerides and glucose. Details on results from the first screening have been published previously.⁶

A more extensive report on this study population after 13 years of observation with relevance to smoking and mortality has been published previously,⁷ and also another report compiled after 25 years of observation, concentrating on the risk in people smoking 1–4 cigarettes.⁸

Second and third screenings

In one county, Finnmark, two new screenings were conducted after 3 and 13 years. In the other two counties (75% of the total study population), the re-screenings were carried out around 5 and 10 years after the initial screening.^{9–11}

At the second screening, all those invited to the initial screening were re-invited and, in addition, a 9% random sample of people aged 20–34 years who were not invited to the first examination. The attendance was 88%.¹²

At the third screening, minor modifications were made in the eligibility to the examinations; the main core of participants in the previous rounds, however, was re-invited. Owing to capacity constraints, only a 10% random sample of the cohort aged 45–49 years at the first screening was invited. The attendance rate was 84%.^{10 13 14}

Table 1 Background characteristics at the last examination of participants*, by category of smoking

	Never smokers	Ex-smokers	Quitters	Moderate smokers	Reducers	Heavy smokers	p Value†
Men (n = 23 978)							
Age (years)	40.6 (7.1)	42.9 (5.9)	41.5 (6.9)	41.4 (7.3)	41.5 (7.4)	41.7 (6.9)	0.576
Cigarette consumption (no/day)	—	13.7 (8.4)	14.7 (8.4)	10.6 (4.6)	9.7 (4.1)	18.3 (6.6)	<0.001
Duration of smoking (years)	—	14.4 (6.9)	20.7 (8.7)	24.0 (7.8)	23.6 (7.7)	24.8 (7.3)	0.006
Systolic BP (mm Hg)	134.0 (20.7)	135.5 (20.2)	135.6 (21.4)	134.9 (19.3)	133.2 (23.6)	134.2 (19.6)	0.359
Diastolic BP (mm Hg)	84.3 (13.8)	86.2 (13.4)	85.3 (14.1)	83.8 (12.7)	83.3 (15.5)	84.1 (13.1)	0.260
Total serum cholesterol (mmol/l)	5.90 (1.28)	6.12 (1.27)	6.14 (1.35)	6.20 (1.29)	6.26 (1.56)	6.27 (1.33)	0.865
Serum triglycerides (mmol/l)	1.94 (1.17)	2.07 (1.28)	2.10 (1.28)	2.05 (1.27)	1.95 (1.16)	2.02 (1.29)	0.342
Physical activity leisure‡	2.32 (0.76)	2.27 (0.73)	2.19 (0.72)	2.15 (0.70)	2.03 (0.68)	2.03 (0.71)	0.982
BMI (kg/m ²)	25.3 (2.9)	25.9 (2.9)	26.0 (3.0)	24.7 (3.0)	24.7 (3.1)	24.9 (3.1)	0.296
Height (cm)	176.3 (6.8)	176.0 (6.5)	175.6 (6.8)	175.0 (6.8)	174.5 (7.0)	175.4 (6.7)	0.016
Disability pension (%)	3.4 (0.2)	2.8 (0.2)	2.9 (0.2)	4.5 (0.2)	6.4 (0.4)	3.8 (0.2)	0.019
Sick leave (%)	2.6 (0.2)	3.1 (0.2)	4.4 (0.2)	4.3 (0.2)	7.2 (0.3)	4.9 (0.2)	0.053
Family history of CHD (%)§	25.6 (4.4)	30.6 (4.6)	26.4 (4.4)	26.6 (4.4)	32.9 (4.7)	29.5 (4.6)	0.386
Women (n = 25 698)							
Age (years)	42.2 (6.5)	41.3 (6.7)	40.2 (7.6)	40.8 (7.3)	41.6 (6.7)	40.9 (6.9)	0.256
Cigarette consumption (no/day)	—	8.5 (5.2)	9.8 (5.6)	9.2 (3.8)	8.6 (2.5)	16.4 (5.2)	<0.001
Duration of smoking (years)	—	10.4 (6.2)	15.5 (7.5)	19.4 (7.3)	20.6 (7.3)	21.6 (6.9)	0.173
Systolic BP (mm Hg)	131.3 (21.0)	128.3 (19.9)	127.7 (22.3)	128.0 (20.0)	129.8 (16.6)	127.1 (20.7)	0.142
Diastolic BP (mm Hg)	82.1 (12.3)	80.7 (12.3)	79.8 (14.0)	80.0 (12.3)	80.0 (11.4)	80.0 (12.9)	0.978
Total serum cholesterol (mmol/l)	6.05 (1.27)	5.97 (1.31)	6.00 (1.43)	6.20 (1.38)	6.11 (1.23)	6.22 (1.37)	0.394
Serum triglycerides (mmol/l)	1.33 (0.74)	1.32 (0.85)	1.42 (0.84)	1.50 (0.90)	1.41 (0.81)	1.52 (0.90)	0.171
Physical activity leisure‡	2.01 (0.57)	2.02 (0.56)	1.98 (0.56)	1.95 (0.57)	1.87 (0.60)	1.84 (0.59)	0.588
BMI (kg/m ²)	25.1 (4.0)	24.6 (3.7)	25.0 (4.1)	23.7 (3.6)	24.0 (3.8)	23.7 (3.7)	0.448
Height (cm)	162.8 (6.1)	163.6 (6.1)	162.7 (6.2)	162.7 (5.9)	161.6 (6.0)	163.4 (6.0)	0.001
Disability pension (%)	4.2 (0.2)	2.5 (0.2)	4.0 (0.2)	4.6 (0.2)	8.7 (0.3)	4.7 (0.2)	0.044
Sick leave (%)	3.0 (0.2)	3.7 (0.2)	4.2 (0.2)	5.0 (0.2)	10.2 (0.3)	5.6 (0.2)	0.034
Family history of CHD (%)§	29.9 (4.6)	32.7 (4.7)	31.1 (4.6)	31.7 (4.7)	33.1 (4.7)	36.7 (4.8)	0.333

BMI, body mass index; BP, blood pressure; CHD, coronary heart disease.

Values are mean (SD) or % (SD).

Table 4 shows the number of participants in each category.

*Participants not reporting cardiovascular disease, diabetes or treatment for hypertension, or symptoms of angina pectoris or atherosclerosis obliterans.

†p Values represent differences between the reducers and the continuous heavy smokers.

‡Physical activity during leisure was graded 1–4, with 4 denoting the heaviest activity.

§One or more of parents or siblings reported to have had a heart infarction or angina pectoris.

The most important elements of the screening programme were kept unchanged throughout all screenings—in particular, that part of the questionnaire covering smoking habits.

The study population

For each individual who underwent two or more examinations, paired results were selected according to the following priority, on the basis of attendance to the

- first and second screenings (n = 53 078);
- first and third screenings (n = 1339); and
- second and third screenings (n = 7075).

Consequently, there was a total of 61 492 people (30 956 men and 30 536 women) who had results from two examinations.

The follow-up period started from the last examination.

Table 2 Mean number of cigarettes per day at first and last examinations in moderate smokers, reducers and heavy smokers, by sex

	Moderate smokers	Reducers	Heavy smokers
Men			
First examination	9.2	22.9	18.8
Last examination	10.6	9.7	18.3
Women			
First examination	8.1	19.5	17.0
Last examination	9.2	8.6	16.4

Table 4 shows the number of participants in each category.

Exclusions

We excluded the following people: men and women who at the last examination reported a history of myocardial infarction, angina pectoris, stroke, diabetes, atherosclerosis of legs, treatment for hypertension, use of glyceryl trinitrate and symptoms indicative of angina pectoris or atherosclerosis obliterans, totalling 6506 people. They were excluded to avoid the problem of reverse causality that might occur if people reduced smoking owing to illness. Also, 1594 men smoking a pipe at the last examination were excluded. The few women who smoked a pipe and the few men and women who smoked cigars were disregarded. Also, 1884 people who started smoking between the first and second examination were excluded, as were 298 people for unspecified reasons. Consequently, the total number of people excluded were 10 282 (5997 men and 4285 women).

Thus, 24 959 men and 26 251 women remained as participants for analysis. All of them underwent two examinations, and at the time of the last investigation they reported neither a history related to cardiovascular disease or diabetes nor symptoms indicative of angina pectoris or atherosclerosis obliterans. They were present or previously daily smokers of only cigarettes, or had never smoked cigarettes daily. For a negligible number of participants, information was missing on some of the confounding variables.

Quantity of daily cigarette consumption

The attending participants reported the actual number of cigarettes smoked per day in a special box in the questionnaire. Here, they were allowed to cite a range, such as 10–15 cigarettes.

Table 3 Determination of serum thiocyanate at the last examination, by category of smoking

	Never smokers	Ex-smokers	Quitters	Moderate smokers	Reducers	Heavy smokers	p Value†
Men (n = 5851)*							
Participants (n)	1168	1038	666	1347	139	1493	
Mean (SD) serum thiocyanate (μmol/l)	30.5 (15.7)	28.7 (17.2)	29.6 (18.0)	65.2 (27.6)	66.8 (28.1)	75.8 (30.7)	0.001
Women (n = 5630)*							
Participants (n)	2213	643	403	1747	45	579	
Mean (SD) serum thiocyanate (μmol/l)	28.5 (13.4)	31.8 (16.8)	34.9 (19.5)	76.2 (31.0)	81.3 (33.4)	91.0 (35.4)	0.074

*Participants not reporting cardiovascular disease, diabetes or treatment for hypertension, or symptoms of angina pectoris or atherosclerosis obliterans.

†p Values represent differences between the reducers and the continuous heavy smokers.

At the examination site, the nurses carefully checked through the questionnaire with the participants. Factory-made and hand-rolled cigarettes had to be counted together. The nurses were instructed that one pack of tobacco for hand rolling (50 g) equalled 50 cigarettes. For those who gave a range of consumption, we used the highest figure.

Categories of participants

To compare our results with those of the Danish study,¹ we found it suitable to apply the categories used by Godtfredsen *et al.*,¹ with a slight modification as follows:

- *Never smokers*: People who at both examinations stated that they did not smoke cigarettes daily, and at the first examination said that they had never done so previously.
- *Ex-smokers*: People who at both examinations stated that they did not smoke cigarettes daily, and at the first examination said that they had done so previously.
- *Quitters*: People who at the first examination stated that they smoked cigarettes daily, but had quit smoking at the time of the last examination.
- *Moderate smokers*: People who at the first examination stated that they smoked 1–14 cigarettes per day, and at the last examination said that they smoked cigarettes daily.

Table 4 Number of participants* and person-years; and deaths from all causes, cardiovascular disease, ischaemic heart disease, smoking-related cancer or lung cancer, number and per 100 000 person-years, age adjusted, by sex and smoking category

	Never smokers	Ex-smokers	Quitters	Moderate smokers	Reducers	Heavy smokers
Men (n = 24 959)						
Participants (n)	6 716	5 140	2 676	5 340	348	4 739
Number of person-years	141 145	109 381	56 512	109 562	6 925	94 452
All causes						
Deaths	663	683	409	1 264	103	1 386
Deaths per 100 000 person-years	507	575	727	1 153	1 498	1 462
Cardiovascular disease						
Deaths	233	238	168	542	41	534
Deaths per 100 000 person-years	184	199	295	494	599	563
Ischaemic heart disease						
Deaths	151	160	114	394	26	381
Deaths per 100 000 person-years	120	133	200	361	393	401
Smoking-related cancer						
Deaths	111	120	79	269	19	354
Deaths per 100 000 person-years	85	99	142	247	279	371
Lung cancer						
Deaths	7	20	16	128	10	189
Deaths per 100 000 person-years	5	16	29	116	134	199
Women (n = 26 251)						
Participants (n)	12 657	2 954	1 584	7 098	127	1 831
Number of person-years	275 859	63 718	34 265	151 963	2 558	38 306
All causes						
Deaths	954	205	154	1 002	27	318
Deaths per 100 000 person-years	332	329	492	690	1 090	873
Cardiovascular disease						
Deaths	186	36	36	291	5	84
Deaths per 100 000 person-years	63	59	118	203	207	236
Ischaemic heart disease						
Deaths	98	14	25	152	3	43
Deaths per 100 000 person-years	34	22	80	106	72	120
Smoking-related cancer						
Deaths	112	27	19	222	8	82
Deaths per 100 000 person-years	40	44	59	153	328	222
Lung cancer						
Deaths	11	7	6	119	2	57
Deaths per 100 000 person-years	4	10	20	82	48	153

*Participants not reporting cardiovascular disease, diabetes or treatment for hypertension, or symptoms of angina pectoris or atherosclerosis obliterans at the last examination.

- **Reducers:** People who at the first examination stated a consumption of ≥ 15 cigarettes per day, and at the last examination reported a consumption of at least 50% less cigarettes per day.
- **Heavy smokers:** People who at the first examination stated a consumption of ≥ 15 cigarettes per day, and at the last examinations did not fall into the categories of reducers or quitters.

Determination of serum thiocyanate

In 1976, determination of serum thiocyanate was introduced at the initial screening in one of the counties. Methods and quality assurance have been described by Foss and Lund-Larsen,¹⁵ who have also reported the results from the initial survey. Tobacco smoke contains hydrogen cyanide, which is absorbed in the lungs and then metabolised to thiocyanate. Owing to its long half life, serum thiocyanate has the advantage that it better reflects the average exposure to tobacco smoke during the previous couple of weeks than the exposure on a given day. The authors underline, however, that diet also contributes to the level of thiocyanate in serum; therefore, this level is not a specific indicator for tobacco smoking. A single determination can neither confirm nor weaken a statement on current individual tobacco consumption.

Analyses of serum thiocyanate in large population groups, however, will give valuable information on the group's tobacco consumption. This is shown by results from the initial screening, which show a remarkable dose-response relationship between mean serum thiocyanate concentration and mean cigarette consumption of the group, based on questionnaire answers of the participants.¹⁵

At the second screening, serum thiocyanate was determined only in Finnmark county; therefore, we have information on serum thiocyanate for not more than 5851 men and 5630 women—that is, 22.4% of the total number of the participants.

End points

We carried out a mortality follow-up by linking our records with the National Register of Causes of Death, using the 11-digit personal identification number as record linkage. Each person accrued person-years from the day of the last examination until the date of death, date of emigration or 31 December 2003.

In addition to deaths from all causes, we studied deaths from the following:

- **Cardiovascular disease:** International classification of diseases (ICD), 8th edition: 390–444.1, 444.3–458; ICD9: 390–459; ICD10: I00–99
- **Ischaemic heart disease:** ICD8 and ICD9: 410–414; ICD10: I20–25
- **Smoking-related cancer:** Types of cancer that the International Agency for Research on Cancer has classed as liable to be caused by smoking, and later applied by Doll *et al.*^{16–17} Thus, the following types of cancer were included:
 - Lip, oral cavity and pharynx: ICD8 and ICD9: 140–149; ICD10: C00–14
 - Nose and nasal sinuses: ICD8 and ICD9: 160; ICD10: C30–31
 - Larynx, trachea, bronchus and lung: ICD8 and ICD9: 161–162; ICD10: C32–34
 - Oesophagus: ICD8 and ICD9: 150; ICD10: C15
 - Stomach: ICD8 and ICD9: 151; ICD10: C16
 - Pancreas: ICD8 and ICD9: 157; ICD10: C25

Figure 1 Deaths from (A) all causes, (B) cardiovascular disease and (C) smoking-related cancer, as cumulative proportions of male and female participants in each category, by time of death. Nelson-Aalen cumulative hazard estimates. The x axis shows the observation years and the y axis shows the cumulative proportions.

- Liver: ICD8: 155, 197.8; ICD9: 155; ICD10: C22
- Kidney and renal pelvis: ICD8 and ICD9: 189.0–1; ICD10: C64–65

- Bladder and ureter: ICD8 and ICD9: 188; ICD10: C66–67
- Myeloid leukaemia: ICD8 and ICD9: 205; ICD10: C92
- Lung cancer: ICD8 and ICD9: 161–162; ICD10: C32–34.

Statistical methods

Comparisons of baseline factors between heavy cigarette smokers and the reducers were made by the *t* test or the χ^2 test. Adjustments for age were carried out by the direct method, by applying the rates in 1-year age groups in the smoking categories during the observation years in the total study population. This was carried out separately for men and women. Relative risks adjusted for confounders were estimated using the Cox's proportional hazards model. Age attained was used as the time variable. The proportional hazards assumption was evaluated by visual inspection of the plots of log(person-years) against log-log(survival probability). The lines in the smoking categories were fairly parallel. In the Nelson–Aalen plots, person-years was the time variable. Values of $p < 0.05$ were considered significant.

RESULTS

Table 1 shows background characteristics of the participants at the start of follow-up.

In both sexes, never smokers, ex-smokers and quitters had a lower mean cholesterol level and a higher degree of leisure physical activity than smokers from any category. Ex-smokers had a shorter duration of smoking than present smokers; duration in quitters was between that in smokers and that in ex-smokers. In all smoking categories, the duration of smoking was 3–5 years less in women than in men.

Comparing reducers with heavy smokers, reducers had a considerably lower stature and a higher frequency of disability pension (both sexes). Male reducers had a shorter duration of smoking, and female reducers had a higher frequency of sick leave.

Table 2 shows the mean number of cigarettes at the first and last examinations in heavy smokers, reducers and moderate smokers.

At the last examination, reducers had a daily cigarette consumption that was 58% lower in men and 56% lower in women compared with consumption at the first examination. In both sexes, moderate smokers had a slightly higher and heavy smokers had a slightly lower daily consumption at the last examination than at the first examination.

Table 3 shows the mean serum thiocyanate values at the time of the last examination in participants in whom thiocyanate was determined.

In both sexes, there are clearly higher mean serum thiocyanate levels in the smokers than in never smokers and previous smokers. Both male and female reducers have mean values that are lower than in heavy smokers, the difference being statistically significant for men ($p = 0.001$), but not for women ($p = 0.074$). Male and female reducers have mean thiocyanate values that are slightly above those of the moderate smokers. They report a cigarette consumption, however, which on average is slightly below those of the moderate smokers (table 2).

Table 4 shows the age-adjusted mortality from any cause, cardiovascular disease, ischaemic heart disease, smoking-related cancer or lung cancer, by sex and smoking category.

In men, the mortality among reducers was lower than that among heavy smokers only for smoking-related cancer and lung cancer. These differences were not beyond chance. Only a few cause-specific deaths were seen in women. Mortality from any cause, however, was non-significantly higher in reducers than in heavy smokers.

Figure 1 shows the cumulative proportions of deaths from all causes, cardiovascular disease and smoking-related cancer, by time of death. For all causes in men, the reducers have slightly lower death rates than heavy smokers during the first 15 years, whereas afterwards the reducers catch up with the heavy smokers. For women, reducers have higher total mortality than heavy smokers during the whole period. For cardiovascular disease in both sexes, there is on the whole no distinct difference between reducers and heavy

Table 5 Adjusted relative risk* (95% CI) of death from all causes, cardiovascular disease, ischaemic heart disease, smoking-related cancer or lung cancer, by smoking category, sex individually and both sexes together, with heavy smokers as reference (23 798 male and 25 698 female participants aged 20–49 at initial screening†)

Sex category, deaths from	Never smokers	Ex-smokers	Quitters	Moderate smokers	Reducers	Heavy smokers
Men (n = 23 798 aged 20–49 years)†						
All causes	0.34 (0.31 to 0.38)	0.36 (0.33 to 0.40)	0.45 (0.40 to 0.51)	0.77 (0.71 to 0.83)	0.99 (0.80 to 1.22)	1.00
Cardiovascular disease	0.31 (0.26 to 0.36)	0.31 (0.26 to 0.36)	0.46 (0.39 to 0.55)	0.85 (0.75 to 0.96)	1.04 (0.75 to 1.43)	1.00
Ischaemic heart disease	0.29 (0.24 to 0.35)	0.30 (0.25 to 0.37)	0.45 (0.36 to 0.56)	0.89 (0.77 to 1.03)	0.94 (0.63 to 1.41)	1.00
Smoking-related cancer	0.23 (0.19 to 0.29)	0.26 (0.21 to 0.33)	0.33 (0.26 to 0.43)	0.65 (0.55 to 0.76)	0.72 (0.45 to 1.18)	1.00
Lung cancer	0.03 (0.01 to 0.06)	0.09 (0.05 to 0.14)	0.14 (0.08 to 0.23)	0.57 (0.45 to 0.72)	0.71 (0.36 to 1.39)	1.00
Women (n = 25 698 aged 20–49 years)†						
All causes	0.35 (0.31 to 0.40)	0.37 (0.31 to 0.44)	0.53 (0.43 to 0.65)	0.77 (0.68 to 0.88)	1.11 (0.74 to 1.67)	1.00
Cardiovascular disease	0.24 (0.19 to 0.32)	0.25 (0.16 to 0.37)	0.47 (0.31 to 0.69)	0.85 (0.66 to 1.10)	0.86 (0.35 to 2.13)	1.00
Ischaemic heart disease	0.28 (0.19 to 0.41)	0.20 (0.11 to 0.37)	0.68 (0.41 to 1.12)	0.92 (0.65 to 1.32)	1.13 (0.35 to 3.67)	1.00
Smoking-related cancer	0.16 (0.12 to 0.22)	0.18 (0.12 to 0.29)	0.24 (0.14 to 0.40)	0.64 (0.49 to 0.83)	1.40 (0.68 to 2.90)	1.00
Lung cancer	0.02 (0.01 to 0.04)	0.07 (0.03 to 0.16)	0.13 (0.05 to 0.29)	0.50 (0.36 to 0.69)	0.51 (0.12 to 2.08)	1.00
Both sexes						
All causes	0.34 (0.32 to 0.37)	0.37 (0.34 to 0.40)	0.47 (0.43 to 0.52)	0.76 (0.72 to 0.82)	1.02 (0.84 to 1.22)	1.00
Cardiovascular disease	0.28 (0.25 to 0.33)	0.30 (0.26 to 0.34)	0.46 (0.39 to 0.54)	0.86 (0.77 to 0.96)	1.02 (0.75 to 1.39)	1.00
Ischaemic heart disease	0.28 (0.24 to 0.33)	0.29 (0.24 to 0.34)	0.48 (0.39 to 0.58)	0.89 (0.78 to 1.02)	0.96 (0.65 to 1.41)	1.00
Smoking-related cancer	0.20 (0.17 to 0.24)	0.25 (0.20 to 0.30)	0.31 (0.25 to 0.39)	0.66 (0.58 to 0.76)	0.86 (0.57 to 1.28)	1.00
Lung cancer	0.03 (0.02 to 0.04)	0.08 (0.06 to 0.12)	0.14 (0.09 to 0.21)	0.55 (0.45 to 0.66)	0.66 (0.36 to 1.21)	1.00

*Adjusted for age, systolic and diastolic blood pressure, total serum cholesterol, serum triglycerides, physical activity during leisure, body mass index (kg/m^2), height, disability pension, sickness leave and family history of coronary heart disease recorded at the last examination. In the section covering both sexes, adjustments are made also for sex.

†Participants not reporting cardiovascular disease, diabetes or treatment for hypertension, or symptoms of angina pectoris or atherosclerosis obliterans at the last examination.

Table 6 Relative risk* (95% CI) of death from all causes in reducers at the second examination who attended the third examination, by group at the third examination, with sustained heavy smokers as reference, in both sexes

	Reducers at second examination, group at third examination			Sustained heavy smokers from second to the third examination
	New quitters	Sustained reducers	Increasesers	
Relative risk	0.47 (0.21 to 1.04)	1.16 (0.73 to 1.85)	1.23 (0.90 to 1.69)	1.00
Participants (n = 271)†	37	69	165	3574
Number of deaths	6	18	42	818
Men (%)	86	75	70	72
Mean (SD) daily cigarette consumption				
First examination	21.8 (6.4)	23.6 (8.8)	20.6 (6.7)	18.0 (4.6)
Second examination	9.1 (3.3)	10.0 (4.0)	9.2 (3.6)	17.5 (6.0)
Third examination	0	10.4 (4.8)	17.2 (6.4)	18.3 (6.6)

*Adjusted for sex.

†Participants not reporting cardiovascular disease, diabetes or treatment for hypertension, or symptoms of angina pectoris or atherosclerosis obliterans. n = 271 aged 20–49 years at initial screening.

smokers. For smoking-related cancer, a clear difference can be seen between the sexes. In men, the reducers have lower death rates than the heavy smokers, whereas the reverse picture is seen in women. For the remaining smoking categories, the mortality levels are as expected. For female ex-smokers, however, the curve falls slightly below that for never smokers. Mortality due to cardiovascular disease among female heavy smokers is comparable with that in men who have quit smoking.

Table 5 shows the adjusted relative risks of death with heavy smokers as reference. Adjustments are made for age, systolic and diastolic blood pressure, serum total cholesterol, serum triglycerides, physical activity during leisure, body mass index (kg/m^2), height, disability pension, sickness leave and recorded coronary heart disease in the family. Table 5 also gives the relative risk for men and women together, to make comparisons with the reports by Godtfredsen *et al.*^{1,3,4} Here adjustments are also made for sex.

For both sexes, reducers have almost the same adjusted relative risk for death from any cause as heavy smokers. For cardiovascular disease and ischaemic heart disease, the adjusted risk in reducers is not significantly different from that in heavy smokers (see table 5). For smoking-related cancer, the adjusted relative risk is lower for male reducers but higher for female reducers compared with the heavy smokers, but neither difference is statistically significant. For both men and women, the adjusted relative risk for lung cancer is clearly lower than in heavy smokers (29% lower in men, 49% in women), but does not reach statistical significance. In both men and women, and in all cause groups, relative risks decrease regularly: from reducers,

through moderate smokers, quitters and ex-smokers, to never smokers. An exception is cardiovascular deaths in women, where we find a somewhat higher relative risk in never smokers than in ex-smokers.

As mentioned earlier, a third screening was carried out in all three counties. In Finnmark (25% of the total study population), this screening took place 10 years after the second screening; in the other two counties, 5 years after the second. Of the 475 reducers at the second examination, 271 attended the third examination. The main reason for the lower number of participants was that only a 10% random sample of the oldest 5-year age group was invited to the third examination.

We grouped these 271 participants by their status at the third examination:

- *New quitters*: Those who quit smoking entirely between the second and third examinations
- *Sustained reducers*: Those who were daily smokers at the third examination, and still reported a consumption of at least 50% less cigarettes per day compared with the first examination
- *Increasesers*: Those who were daily smokers at the third examination, but reported to have a daily cigarette consumption that had increased so much from the second examination that they no longer fulfilled the criteria of reducers.

As a reference group, we introduced sustained heavy smokers—that is, smokers with a daily cigarette consumption of ≥ 15 cigarettes at all three examinations.

Table 7 Relative risk (95% CI)* of dying from all causes and from cardiovascular disease, ischaemic heart disease, smoking-related cancer or lung cancer in daily smokers at both examinations, per 10 decrease in cigarette consumption from the first to the last examination (n = 24 959 men and 26 251 women)†

Deaths from	Heavy smokers		All smokers	
	Number of deaths	Relative risks (95% CI) per 10 cigarette consumption decrease	Number of deaths	Relative risks (95% CI) per 10 cigarette consumption decrease
All causes	1809	0.97 (0.90 to 1.04)	4042	1.00 (0.94 to 1.05)
Cardiovascular disease	650	0.90 (0.79 to 1.03)	1479	0.98 (0.89 to 1.08)
Ischaemic heart disease	447	0.85 (0.73 to 1.01)	989	0.97 (0.87 to 1.10)
Smoking-related cancer	453	0.91 (0.79 to 1.06)	935	0.99 (0.89 to 1.11)
Lung cancer	253	0.97 (0.80 to 1.18)	497	1.01 (0.87 to 1.17)

*Adjusted for sex and consumption level.

†Participants not reporting cardiovascular disease, diabetes or treatment for hypertension, or symptoms of angina pectoris or atherosclerosis obliterans.

Table 8 Comparison of the relative risks (95% CI) in reducers v heavy smokers from the study of the three Norwegian counties with those presented from the Copenhagen Centre for Prospective Population Studies, both sexes*

End point	Copenhagen, Denmark	Three counties, Norway
All deaths	1.02 (0.89 to 1.17)	1.02 (0.84 to 1.22)
Cardiovascular disease	1.01 (0.76 to 1.35)	1.02 (0.75 to 1.39)
Ischaemic heart disease	1.15 (0.94 to 1.40)	0.96 (0.65 to 1.41)
Smoking-related cancer	0.91 (0.63 to 1.31)	0.86 (0.57 to 1.28)
Lung cancer	0.73 (0.54 to 0.98)	0.66 (0.36 to 1.21)

*In the Norwegian counties, the risks are based on deaths only; in Copenhagen the same applies for cardiovascular disease and smoking-related cancer. In Copenhagen, ischaemic heart disease refers to fatal and non-fatal endpoints and lung cancer to incidence.

Table 6 shows the relative risk of death from all causes in the three groups mentioned, with sustained heavy smokers as reference (both sexes together, but adjusted for sex).

Relative risks in none of the three groups differ significantly from the reference group. The tendency of new quitters to have the same relative risk as quitters at the second examination is clear. Sustained reducers have a relative risk in line with the sustained heavy smokers, whereas increasers have an even higher relative risk.

Table 6 gives also the mean daily cigarette consumption in all groups at all three examinations. At the second examination, this mean differed only slightly between the three groups. At the third examination, the mean remained at the same level in sustained reducers, nearly doubled in those who increased smoking, and went down to zero in new quitters. In sustained heavy smokers, the mean consumption was almost constant at all three examinations.

So far, we have concentrated on participants who were heavy smokers, who at the second examination reported a consumption of at least 50% less cigarettes per day. As a last approach, we present relative risk by degree of change in daily cigarette consumption between the first and the last examinations. This is done by running Cox's proportional hazards analyses among the daily smokers at both the first and last examinations, with sex, consumption level and consumption change as covariates. Consumption level was defined as the mean number of cigarettes at the first and last examinations; consumption change was defined as the difference between number of cigarettes at the first and last examinations.

Table 7 shows the relative risks of dying from any cause and from specified smoking-related diseases associated with a per 10 cigarette decrease in cigarette consumption, adjusted for sex and consumption level, for heavy smokers and for all smokers at the first examination. None of the relative risks is significantly different from 1. The largest decrease in risk is somewhat unexpectedly seen for ischaemic heart disease.

In all smokers, the maximum decrease was 80 cigarettes, the maximum percentage decrease being 90%. The maximum increase was 62 cigarettes and the maximum relative increase was 10-fold.

DISCUSSION

Principal findings

In heavy smoking men and women who have reduced their daily cigarette consumption by >50%, mortality of all causes combined is not reduced compared with mortality in those who continued as heavy smokers or those who did not reduce their consumption by as much as 50%. Also for the cause-specific mortality presented, we found no statistical difference between reducers and sustained heavy smokers.

It would be interesting to compare the results from the study of the three Norwegian counties with those presented from the Copenhagen Centre for Prospective Population Studies^{1 3 4}

(table 8). Both studies are based on large populations that have been observed for long periods; both present relative risks for reducers with continuous heavy smokers as reference; and in both studies the relative risks have been adjusted for a series of confounding variables, which are partly the same for the two populations. For the reducers, the consumption decrease is about the same in the two countries.

The results are remarkably similar to each other. For mortality of all causes, and for ischaemic heart disease, cardiovascular disease and smoking-related cancer, there is no significant decrease in risk in reducers compared with sustained heavy smokers. The risk for lung cancer in reducers may differ slightly from this overall picture. For the Copenhagen population, the relative risk in reducers decreased by 27%, with a confidence interval not covering zero effect. For the Norwegian population, the corresponding decrease in risk was 34%, but this decline did not reach statistical significance. Moreover, for the Norwegian population, the analyses based on degree of consumption change showed that the reduction in cigarette consumption had only a modest and non-significant effect on mortality in all cause groups (table 7).

Certainly, the question remains as to whether the lower cigarette consumption at the second examination is due to under-reporting. Also, to what degree compensatory smoking (deeper inhalation and shorter butts) occurs in the reducers is yet unknown. Godtfredsen *et al*¹⁻⁴ have discussed this question thoroughly. For our part, we find that the mean serum thiocyanate level is slightly higher in reducers than in the moderate smokers, although their mean cigarette consumption is slightly lower. This should point to the direction of some under-reporting in the reducers. We emphasise, however, that, compared with the heavy smokers, the reducers have a lower level of serum thiocyanate, although their all-cause mortality is exactly the same as that in the heavy smokers. This indicates that a substantial proportion of their reduction in consumption has been real, although its health consequences are largely unchanged.

One important point that we cannot fully tackle is the extent changes in smoking habits have taken place during the follow-up period. For the subgroup that underwent three examinations, however, it was possible to take into consideration the smoking status of the reducers 5 or 10 years later (table 6). There was no mortality reduction among those who remained as reducers compared with those who remained as heavy smokers. Those who had changed from reducers to quitters had, on the other hand, 50% lower mortality than the sustained heavy smokers.

The sustained reducers had a mean daily cigarette consumption that was almost the same at the second and third examinations—that is, they had maintained their lower level of consumption between 5 and 10 years (in one county for at least 13 years). It is reasonable that their reduced daily consumption then had stabilised and had lasted for the remaining observation period. Some uncertainty still remains

on this point, however, and it would have strengthened the study to elucidate this question further by a new screening later in the observation period.

The consumption increase in most of the subgroup that underwent three examinations may explain the observation made in fig 1, that during the first 15 years male reducers have slightly lower total mortality than heavy smokers, and that they then catch up with the heavy smokers.

It should be added that background characteristics such as serum lipids, blood pressure, physical activity and body mass index (kg/m^2) did not differ significantly between reducers and sustained heavy smokers. A change in cigarette consumption does not go together with a change in lifestyle factors beneficial to health.

A higher prevalence was seen, however, of disability pension and sick leave among the reducers. This could indicate that for some reducers, their lower cigarette consumption accompany a consequence of some disorder. To clarify this question, we conducted separate analyses excluding people who reported disability or sick leave (data not shown). This exercise, however, gave essentially the same risk estimates as those in table 4.

Female reducers have clearly higher death rates than female heavy smokers, both from any cause and from smoking-related cancer. We have no explanation for this phenomenon, beyond the fact that this could be ascribed to chance.

Strengths and weaknesses of the study

The strength of the study is that it includes >51 000 people who were examined according to standardised procedures and observed for a mean of more than two decades. The number of person-years is 566 670 for men and 517 977 for women. We have also information on a series of relevant confounding variables and of a biochemical marker in one of the counties. Furthermore, the follow-up is complete.

Although we have results from a third examination for 57% of the reducers, it would have strengthened the study if we could have made an even better assessment as to what extent some of our findings are due to changes in smoking habits later during the follow-up period.

Possible implications for policymakers

In health education and patient counselling, it is widespread to offer smokers a last resort: "If you are unable to quit, cut down".

Undoubtedly, reduction in consumption may have a place as a temporary measure in systematic smoking cessation. Nevertheless, the results of this study, and those of the Copenhagen Study, make it imperative to reassess this recommendation as a permanent solution, and raise the question whether it offers people false expectations.

The study proves quite clearly the only safe way out of the risk caused by smoking: people who quit smoking have achieved a risk level that is remarkably lower than in those who continued to smoke.

CONCLUSION

In both sexes, a reduction in cigarette consumption by >50% was not associated with a markedly lower risk of all-cause mortality and, specifically, of dying from cardiovascular disease or smoking-related cancer. Accordingly, a reduction in consumption does not seem to bring about harm reduction.

Authors' affiliations

A Tverdal, Norwegian Institute of Public Health, Oslo, Norway
K Bjartveit, National Health Screening Service, Oslo, Norway

What this paper adds

- Only one large prospective study from Copenhagen, Denmark, has so far explored the long-term effects of unassisted reduced smoking.
- The long-term effects of a substantial reduction in smoking did not show any benefits in comparison with persistent heavy smoking, with respect to all-cause mortality and premature deaths from cardiovascular disease, smoking-related cancer and respiratory disease.
- No notable reduction was found in risk of hospital admission for chronic obstructive pulmonary disease, and in incidence of fatal and non-fatal myocardial infarction. Incidence of lung cancer was reduced by about 25%.
- This study, which includes over 51 000 people who were examined twice at an interval of 3–5 years and observed over a mean of more than two decades, confirms in principle the results from the Copenhagen study.
- Compared with sustained heavy smokers, people with reduced cigarette consumption showed no reduced risk of all-cause mortality, or of premature deaths from ischaemic heart disease, cardiovascular disease and smoking-related cancer. There was a modest but insignificant reduction in risk of death from lung cancer.
- This study gives more detailed risk assessment by degree of change in daily cigarette consumption and gives results from a mortality follow-up of people who underwent a third examination some years after the second one, supporting the findings.

Competing interests: KB was involved in national and international tobacco control.

Contributors: KB and AT participated in designing the cardiovascular disease study in Norwegian counties. KB was responsible for the administration of the screening part. AT carried out the data extract and analyses. KB and AT drafted the paper.

REFERENCES

- 1 Godtfredsen NS, Holst C, Prescott E, *et al*. Smoking reduction, smoking cessation, and mortality: a 16-year follow-up of 19,732 men and women from the Copenhagen Centre for Prospective Population Studies. *Am J Epidemiol* 2002;**156**:994–1001.
- 2 Godtfredsen NS, Vestbo J, Osler M, *et al*. Risk of hospital admission for COPD following smoking cessation and reduction: a Danish population study. *Thorax* 2002;**57**:967–72.
- 3 Godtfredsen NS, Osler M, Vestbo J, *et al*. Smoking reduction, smoking cessation, and incidence of fatal and non-fatal myocardial infarction in Denmark 1976–1998: a pooled cohort study. *J Epidemiol Community Health* 2003;**57**:412–16.
- 4 Godtfredsen NS, Prescott E, Osler M. Effect of smoking reduction on lung cancer risk. *JAMA* 2005;**294**:1505–10.
- 5 Bjartveit K, Foss OP, Gjervig T, *et al*. The cardiovascular disease study in Norwegian counties. Background and organization. *Acta Med Scand*, 1979;(Suppl 634):1–70.
- 6 Bjartveit K, Foss OP, Gjervig T. The cardiovascular disease study in Norwegian counties. Results from first screening. *Acta Med Scand*, 1983;(Suppl 675):1–184.
- 7 Tverdal A, Thelle D, Stensvold I, *et al*. Mortality in relation to smoking history: 13 years' follow-up of 68,000 Norwegian men and women 35–49 years. *J Clin Epidemiol* 1993;**46**:475–87.
- 8 Bjartveit K, Tverdal A. Health consequences of smoking 1–4 cigarettes per day. *Tob Control* 2005;**14**:315–20.
- 9 Bjartveit K, ed. *Håndbok for hjerte-karundersøkelsen*. Oslo: Statens Skjermbildefotofering, 1980.

- 10 **Bjartveit K**, ed. *Håndbok for hjerte-karundersøkelsen*. Oslo: Statens Skjermbildefoto-grafering, 1985.
- 11 **Bjartveit K**, Wøien G. *Cardiovascular disease risk factors in Norway. Results from surveys in 18 counties*. Oslo: Statens Helseundersøkelser (National Health Screening Service), 1997.
- 12 **National Health Screening Service, Health Services of Finnmark, Sogn og Fjordane, and Oppland counties, Ullevål Hospital, Central Laboratory, Oslo**. *The cardiovascular disease study in Norwegian counties. Results from second screening*. Oslo: National Health Screening Service, 1988.
- 13 **Westlund K**, Sjøgaard AJ. *Helse, livsstil og levekår i Finnmark. Finnmark III*, ISM skriftserie nr 28. Tromsø: Institute of Social Medicine, 1993.
- 14 **Stensvold I**. *Coffee and health. Is coffee consumption a risk factor for cardiovascular disease or cancer?*. Oslo: National Health Screening Service, 1996.
- 15 **Foss OP**, Lund-Larsen PG. Serum thiocyanate and smoking: interpretation of serum thiocyanate levels observed in a large health study. *Scand J Clin Lab Invest* 1986;**46**:245–51.
- 16 **International Agency for Research on Cancer**. *Tobacco smoke and involuntary smoking*, IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Vol 83. Lyon: International Agency for Research on Cancer, 2004.
- 17 **Doll R**, Peto R, Boreham J, *et al*. Mortality from cancer in relation to smoking: 50 years observation on British doctors. *Br J Cancer* 2005;**92**:426–9.