study area (5.6) is much greater than that in developed countries such as USA (2.6 in 2006, US Census Bureau) and Japan (2.6 in 2005, Ministry of Internal Affairs and Communications). Thus, many non-smoking household members may share indoor environments with smokers.

The WHO FCTC is the first global public health treaty that the 166 WHO member states have already ratified.³ Article 8 of the WHO FCTC addresses the issue of protection from exposure to tobacco smoke in public places, however, there's no statement on protection from domestic SHS. Despite the lack of biological data, our results clearly indicate that SHS exposure occurs in public places and also in households. For women and children in particular, the household likely represents the primary location of exposure.⁶ Public educational campaigns for smoke-free homes⁷⁻⁸ are warranted in Southeast Asian countries, to protect women and children who remain exposed to SHS at home.

Acknowledgements We thank all of the project members (Hideki Yanai, Toru Matsubayashi, Konosuke Morimoto, Le Huu Tho and Truong Tan Minh) for contribution to the original study.

Motoi Suzuki,¹ Vu Dinh Thiem,² Lay-Myint Yoshida,¹ Dang Duc Anh,² Paul E Kilgore,³ Koya Ariyoshi¹

¹Department of Clinical Medicine, Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan; ²National Institute of Hygiene and Epidemiology, Hanoi, Vietnam; ³International Vaccine Institute, Seoul, Republic of Korea, Seoul, Vietnam

Correspondence to Dr Koya Ariyoshi, Department of Clinical Medicine, Institute of Tropical Medicine, Nagasaki University, Sakamoto 1-12-4, Nagasaki, Japan; kari@nagasaki-u.ac.jp

► Supplementary material (supplementary table) is available online only. To view this files please visit the journal online (<http://tobaccocontrol.bmj.com>).

Funding This study was funded by the Program of Founding Research Centers for Emerging and Reemerging Infectious Diseases, Ministry of Education, Culture, Sports, Science and Technology, Japan. The funding source did not have any role in the study design, execution, analysis, writing of the manuscript or conclusions.

Competing interests None.

Ethics approval This study was approved by the Institutional Review Board (IRB) of the National Institute of Hygiene and Epidemiology, Vietnam and the IRB of the Institute of Tropical Medicine, Nagasaki University, Japan.

Contributors KA initiated the study. KA, PEK, DDA and LY were responsible for study conception and design. VDT and DDA collected the baseline data. MS analysed and interpreted the data. MS, LY and KA drafted the manuscript and PEK revised it. All authors had full access to all of the data in the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Accepted 17 December 2009
Published Online First 27 June 2010

Tobacco Control 2010;19:344–345.
doi:10.1136/tc.2009.032227

This paper is freely available online under the BMJ Journals unlocked scheme, see <http://tobaccocontrol.bmj.com/site/about/unlocked.xhtml>

REFERENCES

1. **World Health Organization.** *Protection from exposure to second-hand tobacco smoke: policy recommendations.* Geneva: World Health Organization, 2007.
2. **Samet JM, Yoon S-Y.** WHO Tobacco Free Initiative. *Women and the tobacco epidemic: challenges for the 21st century.* Geneva: World Health Organization, 2001.
3. **World Health Organization.** *WHO framework convention on tobacco control.* Geneva: World Health Organization, 2003.
4. **Suzuki M, Thiem VD, Yanai H, et al.** Association of environmental tobacco smoking exposure with an increased risk of hospital admissions for pneumonia in children under 5 years of age in Vietnam. *Thorax* 2009;64:484–9.
5. **World Health Organization.** *WHO report on the global tobacco epidemic, 2008: the MPOWER package.* Geneva: World Health Organization, 2008.
6. **Wipfli H, Avila-Tang E, Navas-Acien A, et al.** Famri Homes Study Investigators. Secondhand smoke exposure among women and children: evidence from 31 countries. *Am J Public Health* 2008;98:672–9.
7. **Thomson G, Wilson N, Howden-Chapman P.** Population level policy options for increasing the prevalence of smokefree homes. *J Epidemiol Community Health* 2006;60:298–304.
8. **Priest N, Roseby R, Waters E, et al.** Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke. *Cochrane Database Syst Rev* 2008 Oct 8;(4):CD001746.

Newspaper coverage about smoking in leading Chinese newspapers in past nine years

Recent research has clearly identified the importance of media advocacy in advancing tobacco control objectives.^{1–4} Generating news coverage of tobacco control-related issues is a low cost activity that can generate extensive and ongoing coverage to which

Figure 1 The number of articles focused on passive smoking, secondhand smoking or adolescent smoking in important Chinese newspapers from 2000 to 2008.

Figure 2 The number of articles about the Olympics among smoking-related articles in important Chinese newspapers from 2000 to 2008.

millions of citizens, including politicians and decision makers, are exposed. Media advocacy has thus become an important component of comprehensive tobacco control programs.

China has the world's largest population, with 52.4% of adult men and 3.4% of adult women smoking.⁵⁻⁶ In China, news coverage of health issues has increased dramatically in recent years. In our previous study of cancer coverage in Chinese newspapers, there was a sharp increase in cancer-related reports from 578 articles in 2000 to 1403 articles in 2006.⁷ However, there has been little analysis or evaluation of news coverage on smoking in China.

In this study we aim to provide a systematic overview of all tobacco news coverage in a database of major Chinese newspapers over 9 years from 2000 to 2008.

METHODS

Reports about smoking-related matters were obtained from the Database of Important Chinese Newspapers (<http://www.cnki.net>) from 2000 to 2008. As of 18 June 2008, the database included 152 national and 362 local newspapers. The search keywords were fixed as “吸烟”, “抽烟” (吸烟, 抽烟 in English: smoking), “戒烟”, “控烟” or “禁烟” (戒烟, 控烟, 禁烟 in English: smoking cessation, or smoking restriction) in the title to calculate the number of smoking-related articles.

Every smoking-related article obtained from the database was carefully read to ensure that tobacco was central to each article, and articles without smoking-related content were excluded. Articles were allocated to a primary topic classification, which included government law/policy/regulation, health consequences, prevention/cessation programs, affiliated organisation/business news, negative social effects or other. Stories about secondhand smoking, adolescent smoking and smoking-related Olympic Games stories were also noted.

We tested whether the number of articles in each year was the same using the Poisson heterogeneity test.⁸

Table 1 Theme of smoking-related articles from 2000 to 2008 in China

Themes	2000	2001	2002	2003	2004	2005	2006	2007	2008	Total
Health	10	23	13	28	31	40	134	185	71	535*
Government law/policy/regulation	7	1	3	4	10	13	17	21	127	203*
Tobacco prevention/cessation program	18	9	4	21	11	26	54	71	72	286*
Affiliated organisation news	2	1	2	4	5	11	24	32	96	177*
Negative social effects	1	1	1	3	4	10	10	21	27	78*
Other	5	1	3	11	7	21	12	11	16	87*
Total	43	36	26	71	68	121	251	341	409	1366*

*Poisson heterogeneity test $p < 0.001$.

RESULTS

A total of 1366 articles about smoking or smoking control were located (table 1). There was a significant difference in the number of articles appearing each year ($p < 0.001$). Except for small decreases in 2001 and 2002, press attention to smoking increased every year, indicating increasing attention paid to smoking-related matters in recent years. The total number of articles was highest in May (234 articles, $p < 0.05$) because of press reports of 'World No Tobacco Day' on 31 May.

The principal focuses of 1366 smoking-related articles, as judged by headline, are shown in table 1. For each theme, there was a significant difference in the number of articles appearing each year ($p < 0.001$). Almost 2 in 5 of all articles (535/1366) focused on the effects of smoking on health with tobacco prevention/cessation programs being the next most frequent category. Examples included 'Smoking increases the incidence of lung cancer', 'Smoking addiction is a risk factor for coronary heart diseases' and 'Smoking is harmful to eye health'. However, in 2008 stories about government law/policy/regulation concerning tobacco appeared most frequently.

In the years prior to 2005 there were less than 10 reports on secondhand smoke, but in 2007 and 2008 this increased to 44 and 37 reports, respectively.

References to smoking and the Olympics numbered just 3 in 2006, and then rose sharply to 30 in 2007, and further to 67 in 2008, suggesting that the then-forthcoming Olympics stimulated this increase.

The total volume of articles (1366) reflects an ongoing presence of tobacco issues in the news, with an average of nearly one article per day in 2007 and 2008 in Chinese newspapers. The growing coverage may reflect increasing attention to smoking and the endeavours being made to promote tobacco control in China. In our previous research, we found 7643 newspaper reports about cancer in newspapers between 2000 and 2007. Smoking-related articles (1366) were far less frequent over the same period. More attention should be paid to promoting newspaper coverage about smoking in newspapers.

The coverage of issues in newspapers is typically highly correlated with the presentation of the same issues in other media such as radio and television.⁹ Indeed, newspaper and

magazines articles are often cited by the public as common sources of health information.¹⁰

In the US and Australia, secondhand smoke was the most prevalent tobacco-related theme.¹¹ Its low ranking in Chinese newspapers suggests there is plenty of scope to raise coverage of this issue in the news media in China.

There are limitations to our study. For example, we only examined newspapers, but did not analyse the slant of articles. The development of strategies aimed at improving health news reporting deserves more focused attention from the media and researchers.

Acknowledgements This work was supported by the Specialised Research Fund for Heilongjiang High Education.

Yu Liu,¹ Lifei Jiang,² Hai Xiao,³ Qiaojing Liu,¹ Bing Li,⁴ Wanhai Xu¹

¹The Fourth Affiliated Hospital of Harbin Medical University, Harbin, P R China; ²Heilongjiang University, Harbin, P R China; ³Harbin Medical University, Harbin, P R China; ⁴Harbin Center for Disease Control, Harbin, P R China

Correspondence to Dr Wanhai Xu, The Fourth Affiliated Hospital of Harbin Medical University, Yiyuan Road 37, Nangang District, Harbin, Heilongjiang Province 150001, P R China; Liuy.doctor@gmail.com

Funding The Specialized Research Fund for Heilongjiang High Education.

Competing interests None.

Patient consent Obtained.

Contributors YL, LJ and WX contributed to the analysis of content of smoking-related papers. QL and BL contributed to obtaining the smoking-related articles from important Chinese newspaper databases. HX contributed to the statistical analysis of results. YL and LJ made equal contribution to this work.

Provenance and peer review Not commissioned; externally peer reviewed.

Accepted 6 January 2010
Published Online First 14 June 2010

Tobacco Control 2010;**19**:345–346.
doi:10.1136/tc.2009.034793

This paper is freely available online under the BMJ Journals unlocked scheme, see <http://tobaccocontrol.bmj.com/site/about/unlocked.xhtml>

REFERENCES

1. Clegg Smith K, Chapman S. *How the news media influence tobacco use*. National Cancer Institute. *The Role of the Media in Promoting and Controlling Tobacco Use*. Tobacco Control Monograph No. 19. Bethesda, MD: US Department of Health and Human Services; 2008: 327–56. National Institutes of Health, National Cancer Institute. NIH Publication No: 07-6242.
2. Champion D, Chapman S. Framing pub smoking bans: an analysis of Australian print news media coverage, March 1996–March 2003. *J Epidemiol Community Health* 2005;**59**:679–84.
3. Durrant R, Wakefield M, McLeod K, et al. Tobacco in the news: an analysis of newspaper coverage of tobacco issues in Australia, 2001. *Tob Control* 2003;**12**(Suppl 2):ii75–81.
4. Chapman S, Lupton D. *The fight for public health: principles and practice of media advocacy*. London: British Medical Journal Books, 1994.
5. Kenkel D, Lillard DR, Liu F. An analysis of life-course smoking behavior in China. *Health Econ* 2009;**18** (Suppl 2):S147–56.
6. Shi J, Liu M, Zhang Q, et al. Male and female adult population health status in China: a cross-sectional national survey. *BMC Public Health* 2008;**8**:277.
7. Cai JP, Yang LD, Liu ZH, et al. The comprehensive analysis of cancer coverage in China important newspapers between 2000 and 2007. *Support Care Cancer* 2009;**17**:329–32.
8. Armitage P, Berry G, Matthews JNS. *Statistical methods in medical research*. Oxford; Malden, MA: Blackwell Science, 2002:234–5.
9. Wanta W. The messenger and the message: differences across news media. In: McCombs M, Shaw DL, Weaver D, eds. *Communication and democracy: exploring the intellectual frontiers of agenda setting theory*. Mahwah, NJ: Lawrence Erlbaum Associates, 1997:137–51.
10. Ellis JA, Gwynn C, Garg RK, et al. Secondhand smoke exposure among nonsmokers nationally and in New York City. *Nicotine Tob Res* 2009;**11**:362–70.
11. Smith KC, Terry-McElrath Y, Wakefield M, et al. Media advocacy and newspaper coverage of tobacco issues: a comparative analysis of 1 year's print news in the United States and Australia. *Nicotine Tob Res* 2005;**7**:289–99.

Misuse of the Official Information Act by the tobacco industry in New Zealand

The purpose of the Official Information Act (OIA) in New Zealand is to facilitate the democratic process by promoting the accountability of government.¹ Tobacco companies exploit freedom of information acts which make official information publicly available. For example, they disrupted and discredited tobacco control work with information gained from public records requests to the California Department of Health Services in the 1990s.²

We examined all OIA requests to the Ministry of Health (MoH) New Zealand, about tobacco control and smoke-free services from 2005 to August 2009 to determine their origin, content and the charges made for information. Internet searches were