

Health knowledge and perception of risks among Chinese smokers and non-smokers: findings from the Wave 1 ITC China Survey

Jilan Yang,¹ David Hammond,^{1,2} Pete Driezen,² Geoffrey T Fong,^{3,4} Yuan Jiang⁵

¹Department of Health Studies and Gerontology, University of Waterloo, Waterloo, Canada

²Propel Centre for Population Health Impact, University of Waterloo, Waterloo, Canada

³Department of Psychology, University of Waterloo, Waterloo, Canada

⁴Ontario Institute for Cancer Research, Canada

⁵National Tobacco Control Office, Chinese Center for Disease Control and Prevention, Beijing, China

Correspondence to

Dr David Hammond, Department of Health Studies and Gerontology, University of Waterloo, 200 University Ave West, Waterloo, Canada N2L 3G1; dhammond@uwaterloo.ca

Received 24 February 2009

Accepted 29 September 2009

ABSTRACT

Background Awareness of health risks of smoking is strongly associated with smoking behaviour. However, there are no population-based studies of smoking-related health knowledge in China.

Objective The aim of current study was to use a population-based sample from the International Tobacco Control China Wave 1 survey to examine variations between current, former and never smokers' health knowledge about smoking and the impact of health knowledge awareness on smokers' intention to quit.

Methods A face-to-face interview was conducted with 5986 adult smokers and non-smokers from six cities in China. Respondents were asked whether they believed smoking causes heart disease, stroke, impotence, lung cancer, emphysema, stained teeth, premature ageing in smokers and lung cancer in non-smokers. Current smokers were also asked additional questions on how smoking affects their current and future health as well as whether they had plans to quit smoking and if they believe they would have health benefit from quitting.

Findings The overall awareness of health risks of smoking in China was low compared to developed countries. Current smokers in China were less likely than non-smokers and former smokers to acknowledge the consequences of smoking. Current smokers who were more aware of the health consequences of smoking were more likely to intend to quit smoking.

Conclusion These findings highlight the need to increase awareness about the health effects of smoking in China, particularly among current smokers to increase quitting.

INTRODUCTION

At present, approximately 1.3 billion people smoke and more than 5 million people die globally from smoking each year.¹ In contrast to the decline in smoking prevalence among high income countries, tobacco use continues to rise in low and middle-income countries.¹ As a consequence, global tobacco consumption is still increasing and smoking attributable death is projected to rise beyond eight million deaths per year by 2030.

Approximately one-third of the world's smokers—350 million—live in China. Recent estimates suggest that more than a third of Chinese over the age of 15 are smokers, with dramatic differences between genders: approximately 66% of Chinese males smoke compared to only 3% of Chinese females.² In addition, more than 70% of all Chinese, or over 600 million Chinese people, are

regularly exposed to secondhand smoke.³ Given that smoking is a major contributor to chronic obstructive pulmonary disease and lung cancer, both of which are leading causes of death in China, the health burden from tobacco use in China is daunting: tobacco use is estimated to have caused approximately one million premature deaths in 2000 and the death toll is projected to double by 2020 if the present smoking trends continue.^{4 5} Overall, up to 100 million Chinese smokers currently under 30 years old are projected to die from smoking.¹ Thus, the success of global tobacco control relies to a large extent upon reductions of smoking in China.

Tobacco control policies have emerged more slowly in China than in many other countries in the region. Until October 2008, Chinese cigarette packages carried obscure health warnings printed on the side of the pack with the vague text-only message that, 'Smoking may harm your health'. Although tobacco advertisements have been banned from mass media, such as TV, radio and newspapers, tobacco companies have successfully used sponsorships and promotions to maintain a visible marketing presence.⁶ To date, China does not have a national law to restrict smoking in workplaces and other indoor public venues, while smoking remains common in healthcare facilities and educational facilities.⁷ In short, both the state of existing tobacco control regulations in China and their enforcement remain at an early stage.

A primary focus of tobacco control is to raise awareness of the health risks of smoking, and this may be particularly true in low and middle-income countries, most of which have not engaged in campaigns to educate the public on the hazards of smoking. Although increases in perceptions of risk are not always sufficient to reduce smoking on their own, increases in health knowledge are strongly associated with reductions in smoking initiation, increases in cessation behaviour and long-term abstinence from smoking.^{1 8 9} Increases in health knowledge also provide support for other tobacco control measures, such as increases in taxation and more comprehensive workplace smoking restrictions.

In China, as in many other low and middle-income countries, there is little published information about the level of health knowledge about smoking.¹ A study conducted among industrial workers in Shanghai two decades ago found that only 53% of smokers and 76% of non-smokers were generally aware that smoking is harmful to health, and only 51% of smokers reported that smoking



This paper is freely available online under the BMJ Journals unlocked scheme, see <http://tobaccocontrol.bmj.com/site/about/unlocked.xhtml>

causes lung cancer.¹⁰ A more recent study among Chinese physicians found that a majority knew that smoking causes lung cancer and other chronic obstructive pulmonary disease, but only two-thirds (67%) knew that smoking could cause heart disease.¹¹ These findings are particularly troubling given that smoking is a primary risk factor for heart disease, another leading cause of premature death in China.¹² Similar gaps in knowledge have been identified in other studies with healthcare providers.^{13 14} However, to our knowledge, there are no population-based studies of smoking-related health knowledge in China.

The purpose of this study was to examine beliefs about the health risks of smoking in China among never, former and current smokers. The study also examined sociodemographic differences in the levels of health beliefs, as well as associations with intentions to quit smoking among current smokers.

METHODS

The ITC China Survey is a prospective cohort survey of 800 adult smokers and 200 adult non-smokers in each of six cities in China: Beijing, Shanghai, Guangzhou, Shenyang, Changsha and Yinchuan. The six cities in the ITC China Survey were selected based on geographical representativeness and levels of economic development. In each city, the ITC China Survey employed a multistage cluster sampling design. Ten street districts (Jie Dao) were randomly selected from each city with probability of selection proportional to the population size of the Jie Dao. Within each of these Jie Dao, two residential blocks (Ju Wei Hui) were selected, again with probability proportional to the population size of the Ju Wei Hui. Within each selected Ju Wei Hui, a complete list of addresses of the dwelling units (households) was first compiled, and then a sample of 300 households were drawn from the list by simple random sampling without replacement in each city.

In the enumeration process, information on age, gender and smoking status for all adults living in these 300 households was collected. The enumerated 300 households were then randomly ordered, and adult smokers and non-smokers were subsequently approached following the randomised order until 40 adult smokers and 10 adult non-smokers were surveyed. Because of low smoking prevalence among women, one male smoker and one female smoker from every selected household were surveyed whenever possible to increase the sample size for female smokers. At most, one non-smoker was interviewed per household. Where there was more than one person in a sampling category to choose from in a household, the next birthday method was used to select the individual to be interviewed. Up to four visits to a household were made in order to interview the target person(s) within that household. All surveys were conducted 'face-to-face'.

The findings reported here are from Wave 1 of the ITC China Survey, which was conducted between April and August 2006. The Wave 1 cooperation rates, calculated as ratio of completed interviews among the total number of successfully contacts in the initial sample recruitment plan, ranged from 80.0% in Beijing and Guangzhou to 95.0% in Changsha. The response rate, calculated as the ratio of completed interviews among the total number of respondents selected in the initial sample plan, ranged from 39.4% in Yinchuan to 66.0% in Guangzhou. Additional information on survey rates is available in the ITC China technical report.¹⁵

The enumerators and survey interviewers were organised and trained by China Center for Disease Control (CDC) staff in each city, with supervision from the ITC China Project team at the

China National CDC. Several quality control procedures were put in place, including making MP3 recordings for each of the 800 smoker interviews in each of the six cities, with subsequent random monitoring. All materials and procedures used in the ITC China Survey were reviewed and cleared for ethics by the research ethics board at the University of Waterloo and by the institutional review board at China National Center for Disease Control and Prevention. Additional detail on the ITC China survey is described elsewhere.¹⁵

Measures

Demographic variables

Age, gender, education and income were recorded in the survey. Age was categorised into four groups as '18–24; 25–39; 40–54; 55+'. Education level was categorised into 'low (no education and elementary school); middle (junior high school and high school); high (college and higher)'. Household monthly income was classified as '1000 yuan and under; 1001–5000 yuan; 5001 yuan and above'.

Smoking status classification

Respondents who had smoked at least 100 cigarettes in their lifetime and who were smoking at the time of the survey were classified as current smokers. Smokers who smoked at least 100 cigarettes in their lifetime but who were not smoking at the time of the survey were classified as former smokers. Those respondents who had smoked fewer than 100 cigarettes were classified as never smokers.

For current smokers, intention to quit was measured by 'Are you planning to quit in the next month, 6 months, beyond 6 months, or not at all?' and categorised into two groups; 1='In the next month/6 months/beyond 6 month' and 0='not at all'.

Health knowledge of smoking

Respondents were asked whether they believed smoking causes heart disease, stroke, impotence, lung cancer, emphysema, stained teeth, premature ageing in smokers and lung cancer in non-smokers. Responses were coded as 0='no/don't know' versus 1='yes.' A *health knowledge scale* was created by summing the number of 'yes' responses across the 8 diseases/health effects (range=0 to 8).

Other health beliefs

For current smokers, other health beliefs were measured by asking if respondents agreed with the following statements: (1) 'Every cigarette you take damages your health'; (2) 'Tobacco is addictive'; and (3) 'Your cigarette smoke is dangerous to non-smokers'. Responses were given on a five-point Likert scale where 1='strongly disagree' and 5='strongly agree'. Current smokers were also asked three additional questions: (1) 'How much do you think you would benefit from health and other gains if you were to quit smoking permanently in the next 6 months'; (2) 'To what extent, if at all, has smoking damaged your health'; and (3) 'How worried are you, if at all, that smoking will damage your health in the future?' Response categories were 'not at all', 'a little', 'very much' and 'don't know'.

ANALYSIS

All statistical analyses were performed using SAS version 9.1 (SAS Institute Inc). Analyses were weighted on sex and age within each city to account for the sampling design of the ITC China Survey. χ^2 Tests were conducted to examine bivariate differences with smoking status. All odds ratios presented for

Table 1 Sample characteristics (n=5986)

		Never smokers		Former smokers		Current smokers		Overall	
		%	n	%	n	%	n	%	n
Sex	Male	32.4	348	90.8	168	95.1	4494	83.7	5010
	Female	67.6	726	9.2	17	4.9	233	16.3	976
Age	18–24	5.0	54	0.0	0	1.4	66	2.0	120
	25–39	22.4	240	8.7	16	17.6	831	18.2	1087
	40–54	42.3	454	27.6	51	49.4	2335	47.4	2840
	55+	30.4	326	63.8	118	31.6	1495	32.4	1939
Income	Low	18.5	199	18.9	35	19.6	925	19.4	1159
	Moderate	71.1	764	71.9	133	69.6	3291	70.0	4188
	High	2.9	31	2.2	4	3.7	173	3.5	208
	No answer	7.4	80	7.0	13	7.2	338	7.2	431
Education	Low	10.3	111	20.5	38	13.1	620	12.8	769
	Moderate	60.1	645	55.1	102	65.5	3098	64.2	3845
	High	29.6	318	24.3	45	21.3	1009	22.9	1372
Cigarette smoked/day	0–10					34.9	1639		
	11–20					49.0	2304		
	21–30					8.6	404		
	31+					7.5	354		

the logistic regression model were adjusted for gender, age, income and education.

RESULTS

Sample characteristics

As shown in table 1, more than 90% of current and former smokers surveyed were male. The majority of respondents (65.6%) were 25–54 years old and 83.9% of current smokers smoked fewer than 20 cigarettes per day.

Health knowledge overall

Table 2 shows the extent to which respondents agreed that smoking was indeed a cause of each of the eight health effects. Overall, respondents were most likely to agree that smoking causes stained teeth and lung cancer, followed by emphysema and lung cancer from secondhand smoke. Only about 40% of respondents agreed that smoking causes coronary heart disease (CHD), while only one-fifth agreed that smoking causes stroke and impotence. Approximately 7% of respondents agreed that smoking caused all eight of the health effects.

Significant differences were observed in health knowledge between current, former and never smokers. Overall, current smokers agreed with fewer health effects (mean=3.82 out of 8) compared to former smokers (mean=5.5; p<0.001) and never smokers (mean=5.5; p<0.001). For individual health effects, only 68% of current smokers agreed that smoking causes lung

cancer in smokers, compared to more than 90% of former and never smokers. In addition, only 36% of current smokers agreed that smoking causes CHD, compared to more than half of former and never smokers.

Logistic regression models were conducted to examine differences between current, former and never smokers for each health effect, adjusting for sex, age, income and education. Current smokers were significantly less likely than never and former smokers to agree with each of the eight health effects (p<0.001 in all cases). No significant differences were observed between former and never smokers for any of the eight health effects. Linear regression models were also conducted to examine potential differences in the overall health knowledge scale, adjusting for age, sex, income and education. The pattern was the same as for the individual health effects: current smokers had significantly lower health knowledge scores than never smokers (mean difference=-1.5; p<0.001) and former smokers (mean difference=-1.8; p<0.001), with no difference between never and former smokers.

Additional analyses were conducted to examine the association between sociodemographic factors and measures of health knowledge among current smokers. Higher levels of health knowledge were observed among: older smokers (40–54 years vs 25–39 years, OR=1.78 p=0.018); more educated smokers (high vs. low, OR= 2.56 p=0.007); and smokers with lower daily cigarette consumption (21–30 vs 0–10 CPD, OR=0.42 p=0.008; 11–20 vs 0–10 CPD, OR=0.72 p=0.02). No other significant

Table 2 Agreement of each knowledge of health effect by smoking status (n=5986)

Knowledge of health effect	Never smokers (n=1074)		Former smokers (n=185)		Current smokers (n=4732)		Overall	
	%	n	%	n	%	n	%	n
Smoking causes stained teeth	95.0	1026	96.1	178	85.0	4024	87.2*	5288
Smoking causes lung cancer in smokers	91.8	996	90.2	162	68.1	3,244	73.0*	4402
Smoking causes emphysema	86.9	935	87.2	158	59.4	2858	65.2*	3951
Secondhand smoke causes lung cancer in non-smokers	83.2	889	76.9	135	53.2	2,531	59.2*	3555
Smoking causes premature ageing	76.0	821	74.2	131	47.3	2,268	53.3*	3220
Smoking causes CHD	54.2	581	59.2	101	36.3	1,737	40.2*	2419
Smoking causes stroke	35.4	401	37.5	60	16.0	791	20.1*	1252
Smoking causes impotence	29.2	320	29.3	46	16.6	814	19.2*	1180
Knowledge of all health effects	13.2	151	16.3	26	5.2	261	7.0*	438

*Differences between never, former and current smokers significant at p<0.01.

Table 3 Association between covariates/selected health beliefs and intention to quit among current smokers (n=4673)

Health beliefs		%	OR (95% CI)
Smoke is dangerous to non-smokers	Disagree	4.6	1.00
	Neither	4.1	0.41 (0.16 to 1.07)
	Agree	88.5	1.82 (1.24 to 3.21)
Tobacco is addictive	Don't know	2.9	0.37 (0.16, 0.83)
	Disagree	9.1	1.00
	Neither	5.2	0.63(0.39 to 1.00)
Every cigarette damages health	Agree	83.4	1.26 (0.90 to 1.76)
	Don't know	2.4	0.91(0.38, 2.19)
	Disagree	10.6	1.00
Worried about future damage from smoking	Neither	10.3	1.14 (0.70 to 1.84)
	Agree	75.4	5.11 (3.32 to 7.86)
	Don't know	3.7	1.64 (0.67 to 4.00)
Smoking has damaged health	Not at all	33.7	1.00
	A little	41.7	4.01 (3.02 to 5.33)
	Very much	18.9	10.44 (7.59 to 14.38)
How much benefit if quit smoking	Don't know	5.7	1.75 (0.99 to 3.09)
	Not at all	33.2	1.00
	A little	40	2.71 (2.19 to 3.36)
Knowledge of health effects scale (0–8)	Very much	17.5	5.70 (4.35 to 7.47)
	Don't know	9.4	1.13 (0.69 to 1.85)
	Not at all	19.3	1.00
Mean among those NOT intending to quit	A little	32.1	3.50 (2.36 to 5.20)
	Very much	35.2	9.96 (6.64 to 14.94)
	Don't know	13.4	1.11 (0.70 to 1.77)
Mean among those intending to quit	Mean among those NOT intending to quit	3.53	
	Mean among those intending to quit	4.71	
	OR for 1 unit increase in score		1.27 (1.21 to 1.34)

OR, 'unadjusted' odds ratio of intending to quit smoking (0, not planning to quit/don't know; 1, within the next month/within the next 6 months/sometime in the future, beyond 6 months).

differences were observed among sex, income and health knowledge levels.

Other health beliefs

Current smokers were asked to report additional health beliefs (see table 3). The vast majority of current smokers agreed that smoke is dangerous to non-smokers (88.5%) and tobacco is addictive (83.4%). About three-quarters of current smokers agreed that every cigarette damages health. In contrast, less than one-fifth of smokers agreed that smoking has damaged his/her health, one-third of current smokers (33.7%) reported no worry about the future damage from smoking and approximately 35% of current smokers agreed that they would experience future health benefits very much if they quit smoking within the next 6 months.

Predictors of health belief perception among current smokers

Analyses were conducted to examine the extent to which health knowledge and other health beliefs among current smokers predicted intention to quit. Overall, 31.9% of current smokers intended to quit at some point in the future. Most of the selected health beliefs were significantly associated with intentions to quit in bivariate analyses, except the belief that 'tobacco is addictive' (see table 3).

We conducted multivariate analyses to examine the predictive relation of health knowledge and health beliefs on intention to quit in greater detail. Measures of health beliefs and the health

knowledge scale were entered in a logistic regression model predicting intention to quit, adjusting for sex, age, income and education (see table 4). The health knowledge scale, and the health beliefs of 'smoke is dangerous to non-smokers; tobacco is addictive' were not significantly associated with intentions to quit; however, all other health beliefs significantly predicted intentions to quit among current smokers.

DISCUSSION

The findings of this study shed light onto beliefs about smoking in China—one of the leading public health threats to the country. The results indicate reasonable levels of health knowledge for health effects such as lung cancer, comparable with levels in Western countries.¹⁶ However, the findings reveal major gaps in the knowledge of other health effects. For example, approximately 40% of respondents endorsed smoking as a risk factor for heart disease and only one-fifth acknowledged smoking causes stroke—both leading causes of death in China.¹² In addition, less than two-thirds of respondents agreed that secondhand smoke can cause lung cancer.

In terms of health knowledge towards smoking among different smoking status groups, the findings indicate that Chinese smokers were less likely to agree to health effects than either never smokers or former smokers. For example, over 90% of never and former smokers agreed that smoking causes lung cancer, compared to only two-thirds of smokers. Similarly, about 83% of never smokers and 77% of former smokers agreed that secondhand smoke causes lung cancer compared to only half of smokers. These differences persisted even after adjusting for sociodemographic factors, which suggests that the lower levels of health beliefs among Chinese smokers may reflect either an optimistic bias, lower exposure to health information or both. This finding is similar to patterns observed in Western countries, where smokers systematically underestimate their personal risk from smoking, presumably in attempt to minimise cognitive dissonance from smoking and shield themselves from worry.^{17–19}

Compared to historical estimates, the findings suggest that levels of health knowledge about the dangers of smoking may be increasing in China. In particular, data from two decades ago shows that only 51% of smokers know smoking causes lung cancer,¹⁰ although differences in the sample profiles complicate comparisons between studies. In addition, the current study only included respondents living in highly developed, urban areas. One might expect levels of health knowledge to be lower in rural areas of China, where access to health information is typically lower. More generally, levels of health knowledge among Chinese smokers were considerably lower than levels previously reported in Western countries.¹⁶ For example, in Canada and Australia—countries widely acknowledged to be leaders in tobacco control—approximately 90% of smokers agreed that smoking causes heart disease and 80% agreed that smoking causes stroke.¹⁶ In contrast, only 36% and 16% of Chinese smokers in the current study agreed that smoking causes heart disease and stroke, respectively. The knowledge awareness gap also exists in secondhand smoke. In Canada and Australia, over 70% of smokers agreed cigarette smoking causes lung cancer in non-smokers, while only about half of Chinese smokers agree with this statement in current study.

The findings of this study indicate that relatively few Chinese smokers intend to quit smoking. Less than a third of smokers indicated that they intend to quit at any point in the future—levels far below Western countries such as Canada and Australia, where approximately three-quarters of all smokers

Table 4 Adjusted odds ratios† from logistic regression of selected health beliefs on intention to quit (n=4673)

Covariate		OR (95% CI)
Sex	Male	1.00
	Female	1.07 (0.58 to 1.95)
Age	18–24	1.00
	25–39	1.02(0.45 to 2.32)
	40–54	1.28 (0.50 to 3.27)
	55+	1.23 (0.51 to 2.97)
Income	Low	1.00
	Moderate	1.09 (0.87 to 1.37)
	High	1.36 (0.77 to 2.41)
	No answer	0.91 (0.57 to 1.46)
Education	Low	1.00
	Moderate	0.91 (0.67 to 1.24)
	High	0.92 (0.66 to 1.28)
Cigarettes smoked/day	0–10	1.00
	11–20	0.68 (0.57 to 0.81)
	21–30	0.49 (0.35 to 0.68)
	31+	0.48 (0.33 to 0.70)
Knowledge of health effects scale (OR for 1 unit increase)		1.03 (0.98 to 1.09)
Smoke is dangerous to non-smokers	Disagree	1.00
	Neither	0.62 (0.29 to 1.32)
	Agree	0.97 (0.59 to 1.60)
	Don't know	0.52 (0.21, 1.28)
Tobacco is addictive	Disagree	1.00
	Neither	1.02 (0.57 to 1.84)
	Agree	1.05 (0.67 to 1.65)
	Don't know	1.50 (0.54, 4.20)
Every cigarette damages health**	Disagree	1.00
	Neither	0.98 (0.61 to 1.57)
	Agree	1.65 (1.15 to 2.38)
	Don't know	2.05 (1.07,3.93)
Worried about future damage from smoking***	Not at all	1.00
	A little	2.13 (1.63 to 2.78)
	Very much	3.70 (2.35 to 5.83)
	Don't know	1.90 (1.18, 3.08)
	Not at all	1.00
Smoking has damaged health*	A little	1.18 (0.87 to 1.60)
	Very much	1.48 (1.13 to 1.94)
	Don't know	0.91 (0.56, 1.47)
	Not at all	1.00
How much benefit if quit smoking***	A little	2.00 (1.40 to 2.86)
	Very much	3.85 (2.52 to 5.88)
	Don't know	0.91 (0.57 to 1.45)
	Not at all	1.00

Significant levels are indicated as follows:

*p<0.05; **p<0.01; ***p<0.001.

†Odds of having intention to quit (0, not planning to quit/don't know, 1, within the next month/within the next 6 months/sometime in the future, beyond 6 months).

intend to quit smoking.¹⁴ As in other studies, intention to quit was positively associated with health beliefs about smoking.^{16 20} In particular, smokers who reported greater worry about the future health effects of smoking and smokers who reported health benefits from quitting were most likely to intend to quit. The direction of this association is likely to be a reciprocal one: greater health beliefs are likely to increase intentions to quit, just as greater intentions are also likely to increase perceived risk and agreement with health effects.

Surveys capable of monitoring perceptions of risk and health knowledge among smokers are not well established globally and they are particular rare among low and middle income countries.¹ What few data exist from low and middle-income countries, suggest that health knowledge is considerably lower than in high-income countries, consistent with the current

findings from China.^{21–24} However, more direct comparisons across studies are precluded by different sample profiles and the use different measures to assess health knowledge.

LIMITATIONS

The limitations of this study are common to population-based surveys, including potential biases from non-response. For example, in this study, respondents were usually interviewed at night which resulted in lower proportion of young people, especially in the age group 18–24, in the sample than in the general population. The findings should also be interpreted within the context of the sampling frame: only smokers in large urban areas were sampled in this study and one would expect different results in rural areas. It should also be noted that the measures of health knowledge used in this study—agreement with a list of health effects—represents a fairly low threshold for measuring health knowledge. For example, one might expect lower levels with the use of unprompted questions.

IMPLICATIONS

China accounts for approximately one-third of the world's smokers and China is one of the biggest cigarette production countries in the world.^{1 25} As a result, health knowledge about smoking in China represents an important indicator for tobacco control in China and global efforts to reduce the health burden from tobacco use. Whereas research from high-income countries shows increasing public concern about the health effects of smoking, the current findings suggest that China lags in this key outcome. However, China has ratified the Framework Convention on Tobacco Control—the world's first treaty devoted to public health—which includes provisions in key areas of tobacco control, including more prominent health warnings, restrictions on tobacco marketing and more prominent public smoking restrictions. Effective implementation of these policy measures will be critically important to increasing perceptions of risk from smoking and helping China to avert the looming public health crisis from tobacco use.

Funding The ITC China Project was supported by grants from the US National Cancer Institute (R01 CA125116 and the Roswell Park Transdisciplinary Tobacco Use Research Center (P50 CA111236)), Canadian Institutes of Health Research (79551), Chinese Center for Disease Control and Prevention, and the Ontario Institute for Cancer Research. Additional support was provided by the Propel Centre for Population Health Impact with funds from the Canadian Cancer Society of the National Cancer Institute of Canada/Canadian Cancer Society. The funding sources had no role in the study design, in collection, analysis, and interpretation of data, in the writing of the report, and in the decision to submit the paper for publication.

Competing interests None.

Patient consent Obtained.

Ethics approval All materials and procedures used in the ITC China Survey were reviewed and cleared for ethics by the Research Ethics Board at the University of Waterloo and by the Institutional Review Board at China National Center for Disease Control and Prevention.

Contributors All of the authors contributed in each of the following ways: conception and design, acquisition of data or analysis and interpretation of data; drafting the article or revising it critically for important intellectual content; final approval of the version published.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. *WHO Report on the global tobacco epidemic, 2008: the MPOWER package*. Geneva: World Health Organization, 2008.
2. **Yang G**, Ma J, Liu N, et al. Smoking and passive smoking in Chinese, 2002. *Zhonghua Liu Xing Bing Xue Za Zhi* 2005;**26**:77–83.
3. **Yang G**, Fan L, Tan J. Smoking in China: findings of the 1996 National Prevalence Survey. *JAMA* 1999;**282**:1247–53.

4. *Preventing chronic diseases: a vital investment: WHO Global Report*. Geneva: World Health Organization, 2005. http://www.who.int/chp/chronic_disease_report/full_report.pdf.
5. **Lin H**, Murray M, Cohen T, *et al*. Effects of smoking and solid-fuel use on COPD, lung cancer, and tuberculosis in China: time-based, multiple risk factor, modelling study. *Lancet* 2008;**372**:1473–83.
6. **Li L**, Yong HH, Borland R, *et al*. Reported awareness of tobacco advertising and promotion in China compared to Thailand, Australia and the USA. *Tob Control* 2009;**18**:222–7. Published online First: 29 Mar 2009. doi:10.1136/tc.2008.027037.
7. **Li Q**, Hyland A, O'Connor J, *et al*. Support for smoke free policies among smokers and non-smokers in six cities in China: ITC China Survey. *Tob Control* 2010;**19** (Suppl 1):i40–46.
8. **Curry SJ**, Grothaus L, McBride C. Reasons for quitting: intrinsic and extrinsic motivation for smoking cessation in a population-based sample of smokers. *Addict Behav* 1997;**22**:727–39.
9. **Hyland A**, Li Q, Bauer JE, *et al*. Predictors of cessation in a cohort of current and former smokers followed over 13 years. *Nicotine Tob Res* 2004;**6**(suppl 3):S363–69.
10. **Wang W**, Annette JD. Cigarette smoking and sick leave in an industrial population in Shanghai, China. *Int J of Epidemiol* 1992;**21**:293–7.
11. **Jiang Y**, Ong MK, Tong EK, *et al*. Chinese physicians and their smoking knowledge, attitudes, and practices. *AJPM* 2007;**33**:15–22.
12. **He J**, Gu D, Wu X, *et al*. Major causes of death among men and women in China. *N Engl J Med* 2005;**353**:1124–34.
13. **Yan J**, Xiao S, Ouyang D, *et al*. Smoking behavior, knowledge, attitudes and practice among health care providers in Changsha city, China. *Nicotine Tob Res* 2008;**10**:737–44.
14. **Chan SSC**, Sarna L, Wong DCN. Nurses' tobacco-related knowledge, attitudes, and practice in four major cities in China. *JNS* 2007;**39**:46–53.
15. Wave 1(2006) ITC China technical report. <http://www.itcproject.org/library/countries/itcchina/reports/technical/finalitcch>.
16. **Hammond D**, Fong GT, McNeill A, *et al*. Effectiveness of cigarette warning labels in informing smokers about the risks of smoking: findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control* 2006;**15**(Suppl III):iii19–25.
17. **Weinstein ND**. Accuracy of smokers' risk perceptions. *Ann Beh Med* 1998;**20**:135–40.
18. **Slovic P**. *Smoking: risk, perception & policy*. Thousand Oaks, CA: Sage, 2001.
19. **Weinstein N**, Slovic P, Waters E, *et al*. Public understanding of the illnesses caused by smoking. *Nicotine Tob Res* 2004;**6**:349–55.
20. **Romer D**, Jamieson P. The role of perceived risk in starting and stopping smoking. In: Slovic, ed. *Smoking: risk, perception, and policy*. Thousand Oaks California: Sage, 2001.
21. **Unsal M**, Hamzacebi H, Dabak S, *et al*. Smoking status and levels of knowledge regarding cigarettes among primary school teachers. *South Med J* 2008;**101**:1227–31.
22. **Singh V**, Gupta R. Prevalence of tobacco use and awareness of risks among school children in Jaipur. *J Assoc Physicians India* 2006;**54**:609–12.
23. **Al-Haddad NM**, Hamadeh RR, Bahram SA. Public knowledge and attitudes towards passive smoking. *Saudi Med J* 2005;**26**:2004–6.
24. **Khan JA**, Hussain SF, Malik A, *et al*. Knowledge, attitudes and the prevalence of smoking among hospital attendants in a developing country. *Trop Doct* 2003;**33**:231–4.
25. **The World Market for Tobacco**. Euromonitor International 2007. http://www.euromonitor.com/The_World_Market_for_Tobacco (accessed 4 Apr 2008).