

Supplementary Table 2. Smoking-attributable mortality rates with 95% confidence intervals among men aged 35-64 years, 1990-1994 and 2000-2004, by occupational class

	1990-1994				2000-2004			
	Manual		Non-Manual		Manual		Non-Manual	
	SAMR	95% C.I.	SAMR	95% C.I.	SAMR	95% C.I.	SAMR	95% C.I.
Finland	231,2	(220,6 - 243,2)	73,7	(66,5 - 82,2)	150,5	(142,7 - 159,1)	38,8	(34,8 - 43,5)
Denmark	195,6	(184,6 - 206,6)	99,2	(91,1 - 108,8)	102,4	(96,5 - 109,4)	42,4	(36,9 - 48,2)
England & Wales	205,4	(187,3 - 224,2)	99,4	(85,2 - 113,9)	127,4	(111,1 - 144,6)	51,5	(42,3 - 62,0)
France	314,5	(269,4 - 363,9)	129,8	(106,4 - 158,3)	303,0	(262,0 - 346,7)	90,9	(74,0 - 110,8)
Italy (Turin)	198,8	(182,5 - 218,6)	120,7	(105,5 - 138,4)	146,7	(129,5 - 164,7)	68,7	(58,3 - 80,7)

Only those countries for which the indicator occupational class was available in both periods were included in this table.

SAMR: Smoking-Attributable Mortality Rate, calculated with the Preston-Glei-Wilmoth method (for details, see text).

Rates are expressed in deaths per 100.000 person-years.