



OPEN ACCESS

Exploring the gateway hypothesis of e-cigarettes and tobacco: a prospective replication study among adolescents in the Netherlands and Flanders

Thomas Martinelli ^{1,2}, Math J J M Candel,³ Hein de Vries,⁴ Reinskje Talhout,⁵ Vera Knapen,⁴ Constant P van Schayck,⁶ Gera E Nagelhout^{1,4}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/tobaccocontrol-2021-056528>).

¹IVO, The Hague, The Netherlands

²Tranzo, Tilburg School of Social and Behavioral Sciences, Tilburg University, Tilburg, The Netherlands

³Department of Methodology and Statistics, Maastricht University, Maastricht, The Netherlands

⁴Department of Health Promotion, Maastricht University, Maastricht, The Netherlands

⁵Laboratory for Health Protection Research, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

⁶Department of Family Medicine, Maastricht University, Maastricht, The Netherlands

Correspondence to

Thomas Martinelli, IVO, The Hague, The Netherlands; martinelli@ivo.nl

Received 2 February 2021

Accepted 10 June 2021

Published Online First

5 July 2021



Watch Video
tc.bmj.com



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: Martinelli T, Candel MJJM, de Vries H, *et al.* *Tob Control* 2023;**32**:170–178.

ABSTRACT

Background Studies demonstrated that adolescent e-cigarette use is associated with subsequent tobacco smoking, commonly referred to as the *gateway effect*. However, most studies only investigated gateways from e-cigarettes to tobacco smoking. This study replicates a cornerstone study revealing a positive association between both adolescent e-cigarette use and subsequent tobacco use; and tobacco and subsequent e-cigarette use in the Netherlands and Flanders.

Design The longitudinal design included baseline (n=2839) and 6-month (n=1276) and 12-month (n=1025) follow-up surveys among a school-based cohort (mean age: 13.62). Ten high schools were recruited as a convenience sample. The analyses involved (1) associations of baseline e-cigarette use and subsequent tobacco smoking among never smokers; (2) associations of e-cigarette use frequency at baseline and tobacco smoking frequency at follow-up; and (3) the association of baseline tobacco smoking and subsequent e-cigarette use among non-users of e-cigarettes.

Findings Consistent with prior findings, baseline e-cigarette use was associated with higher odds of tobacco smoking at 6-month (OR=1.89; 95% CI 1.05 to 3.37) and 12-month (OR=5.63; 95% CI 3.04 to 10.42) follow-ups. More frequent use of e-cigarettes at baseline was associated with more frequent smoking at follow-ups. Baseline tobacco smoking was associated with subsequent e-cigarette use (OR=3.10; 95% CI 1.58 to 6.06 at both follow-ups).

Conclusion Our study replicated the positive relation between e-cigarette use and tobacco smoking in both directions for adolescents. This may mean that the gateway works in two directions, that e-cigarette and tobacco use share common risk factors, or that both mechanisms apply.

INTRODUCTION

Debates about electronic cigarettes are dividing many of those concerned with tobacco control. Proponents of e-cigarettes point at evidence which suggests that e-cigarettes are an effective smoking cessation aid¹ and that e-cigarette use is substantially less harmful than tobacco smoking.² Therefore, e-cigarettes could dramatically reduce disease and death caused by smoking.^{3,4} However, opponents are concerned that e-cigarettes attract new generations of youth into nicotine addiction,⁵ that most e-cigarette users simultaneously use tobacco^{6,7} and that it acts as a gateway to smoking tobacco.^{8,9}

The *gateway* or *stepping stones* theory originates from the 1970s as a mix of academic and popular explanations of the observed sequence from cannabis use to other illicit drug use.^{10–12} While initially descriptive, the theory was also used to explain causal relationships of substance use.¹³ However, the theory was and remains controversial as it fails to exclude alternative explanations, particularly the notion that all substance use is associated with shared characteristics of individuals, especially a propensity to use drugs.¹⁴ More recently, the gateway theory is applied to e-cigarettes in both policy and research.¹⁵ The European Tobacco Products Directive, for example, states that ‘Electronic cigarettes can develop into a gateway to nicotine addiction and ultimately traditional tobacco consumption’.¹⁶

Various cohort studies show that in the last decade e-cigarette use among adolescents increased while tobacco use decreased.^{17–20} Simultaneously, longitudinal studies show that e-cigarette use is associated with initiation of tobacco smoking. The first study that revealed such association among adolescents was published in 2015.^{21,22} These findings generated ample media attention and led to restrictions for e-cigarettes in the USA, prohibiting the sale to persons under 18 years.²³ Since then, cohort studies^{3,17,20,24–28} have followed finding similar associations, including in the Netherlands.²⁹ Despite the above research into the relation between e-cigarette use and tobacco smoking, only few studies examined the reverse relation between tobacco smoking and initiation of e-cigarette use,^{21,30,31} which has also been found for tobacco and alcohol use, for example.³²

In this paper, we present a replication study with new data using the same protocol as the cornerstone study by Leventhal *et al.*^{21,22} Replication is crucial to the scientific method as it enables one to build on demonstrated and confirmed findings.³³ We collected data among adolescents in the Netherlands and Flanders, the Dutch-speaking region of Belgium. Both regions have similar legislation in which e-cigarettes (with and without nicotine) are treated as tobacco products, meaning that sales to minors (under 18 years) and advertising are prohibited.^{34,35} This is different in the original study setting in the USA, where few federal restrictions on e-cigarette marketing existed at the time of the study and some leading e-cigarette brands have been investigated for particularly targeting youth.^{36–39} In the Netherlands, in 2019, 25% of youth between 12




Figure 1 Ever use of tobacco and e-cigarettes in Dutch youth. Source: Health Behaviour in School-aged Children (HBSC 2001, 2005, 2009, 2013, 2017) and Peilstationsonderzoek (Peil 2003, 2007, 2011, 2015, 2019).

and 16 years old had ever tried an e-cigarette and 17% had ever tried smoking tobacco¹⁹ (see [figure 1](#)). Currently, more youth experiment with e-cigarettes than with tobacco. With this paper, we aim to answer the following research questions:

1. Is e-cigarette use associated with subsequent combustible tobacco use?
2. Is e-cigarette use frequency associated with subsequent tobacco smoking frequency?
3. Is combustible tobacco use associated with subsequent e-cigarette use?

METHODS

Participants and procedure

Data were collected with online surveys between September 2018 and December 2019 throughout the Netherlands and Flanders. Since high schools are very hard to recruit for research, due to the large number of study requests and ongoing studies, we used multiple recruitment strategies. The research was presented as a study on smoking, alcohol, drugs and other risk behaviours. Different national and regional organisations (including addiction services, school health promotion and youth organisations) were approached to help contact schools. A representative selection of 82 schools were approached by telephone, of which 50 received recruitment packages. At 14 schools we reached the right contacts by phone. Another 10 schools were approached through informal networks of the research group. Lastly, a mass recruitment e-mail with reminder was sent to 580 schools in the Netherlands and 1343 in Flanders (Belgium). In total, 10 schools responded with interest in our study, and of those, all agreed to participate, including eight schools in the Netherlands and two in Flanders. The schools were provided with informed consent forms about the study, which they disseminated among parents and students. Students enrolled through passive consent and were excluded if they or their parents refused participation. Data collection involved three waves that took place 6 months apart. We provided participating schools with links to online surveys at each wave and the schools administered the surveys in the classes. Since recruitment of schools was difficult, some schools enrolled later. This meant that some schools only participated 6 months in the study and did not complete all three surveys. This partly explains the attrition rates across follow-ups (see online supplemental figure 1). Other reasons for attrition were that some students left the school (either to change schools or because they graduated), some schools had shop classes (such as

wood or metal shop class, which meant they were at other locations), illness at the day of the survey and, lastly, one school could not be reached to conduct a follow-up despite many reminders.

Measures

All measures are the same as in the replicated study²¹ and possess adequate psychometric properties in youth samples.^{40–44}

E-cigarette and combustible tobacco use

Youth Risk Behavior Surveillance (YRBS)⁴⁰ and Monitoring the Future Surveys (MTF)⁴¹ based items measured lifetime (ever) and past 6-month use (yes/no) of e-cigarettes and combustible tobacco products, which included: combustible cigarettes (including ‘even a few puffs’), full-size cigars, little cigars/cigarillos and hookah waterpipes. Lifetime (ever vs never) e-cigarette use at baseline was the primary exposure variable. Outcome variables were any use in the last 6 months (yes/no) of: (1) any combustible tobacco product; (2) combustible cigarettes; (3) cigars (full-size cigars or little cigars); (4) hookah; and (5) the number of different combustible tobacco products (range: 0–3). The terms ‘ever-smokers’ and ‘never-smokers’ refer to participants who either have ever or never used any of the three combustible tobacco products, respectively. The four-level continuous e-cigarette use frequency variable was categorised as never, prior (ever use with no past 30-day use), infrequent (1–2 days during the past 30 days) or frequent (≥ 3 days during the past 30 days). The cigarette use frequency variable consisted of non-smokers (smoked 0 day in the past 30 days); infrequent smokers (smoked 1–2 days in the past 30 days); and frequent smokers (smoked ≥ 3 days in the past 30 days).

Covariates

Variables that are potentially associated with risk of combustible tobacco use initiation based on previous literature^{45–50} were selected a priori as covariates, as they potentially overlap with both e-cigarette use and tobacco use. Covariate categories are described below.

Sociodemographics

Age, gender, ethnicity and highest parental education represented sociodemographics ([table 1](#) contains response categories). In the main analysis, Belgian participants from Belgian (Flanders) schools were coded as Dutch students, as they belonged to the ethnic majority (native) group of their school.

Environmental variables

Environment indicators included family living situation, assessed by asking: ‘Who do you live with most of the time?’ (both biological parents/other).⁴⁶ Family history of smoking was assessed by asking: ‘Does anyone in your immediate family (brothers, sisters, parents, or grandparents) have a history of smoking cigarettes?’ (yes/no). Peer smoking was measured by asking: ‘In the last 30 days, how many of your five closest friends have smoked cigarettes?’ (range: 0–5).⁵¹

Intrapersonal factors

Personality traits, mental health and psychological processes that are linked to risky behaviour, experimentation and smoking were assessed. Depressive symptoms were assessed through the 20-item Center for Epidemiologic Studies Depression Scale⁴³ composite sum of past week frequency ratings (eg, 0=Rarely or none of the time (0–1 day) to 3=Most or all of the

Table 1 Sample characteristics by baseline ever e-cigarette use status among baseline never smokers

Outcome	Baseline e-cigarette use			Contrast by ever e-cigarette use P value
	Overall (n=2185)	Never use (n=1994)	Ever use (n=191)	
Sociodemographics				
Sex, n (%)				<0.001*
Girls	1160 (53.1)	1097 (55.0)	63 (33.0)	
Boys	1025 (46.9)	897 (45.0)	128 (67.0)	
Age, M (95% CI)	13.62 (13.56 to 13.67)	13.57 (13.51 to 13.62)	14.16 (13.99 to 14.34)	<0.001†
Ethnicity, n (%)				0.063*
Dutch	1978 (90.5)	1816 (91.1)	162 (84.8)	
Belgian	76 (3.5)	65 (3.3)	11 (5.8)	
Polish	8 (0.4)	7 (0.4)	1 (0.5)	
Turkish	30 (1.4)	24 (1.2)	6 (3.1)	
German	5 (0.2)	5 (0.3)	0	
British	4 (0.2)	4 (0.2)	0	
Moroccan	16 (0.7)	13 (0.7)	3 (1.6)	
Chinese	6 (0.3)	4 (0.2)	2 (1.0)	
Italian	5 (0.2)	5 (0.3)	0	
Spanish	5 (0.2)	5 (0.3)	0	
Other, don't want to say or unknown	52 (2.4)	47 (0.2)	1 (0.5)	
Highest parental education, n (%)				0.029‡
Graduate degree	271 (12.4)	248 (12.4)	23 (12.0)	
College graduate	478 (21.9)	426 (21.4)	52 (27.2)	
Secondary higher education	368 (16.8)	347 (17.4)	21 (11.0)	
Vocational education	301 (13.8)	267 (13.4)	34 (17.8)	
Secondary vocational education	211 (9.7)	189 (9.5)	22 (11.5)	
Lower vocational education	42 (1.9)	34 (1.7)	8 (4.2)	
Primary school	14 (0.6)	13 (0.7)	1 (0.5)	
None	23 (1.1)	22 (1.1)	1 (0.5)	
Don't know	477 (21.8)	448 (22.5)	29 (15.2)	
Environmental factors				
Lives with both biological parents, n (%)	1729 (79.1)	1606 (80.5)	123 (64.4)	<0.001*
Family history of smoking, n (%)	1522 (69.7)	1364 (68.4)	158 (82.7)	<0.001*
Peer smoking, M (95% CI)	1.32 (1.29 to 1.36)	1.27 (1.24 to 1.31)	1.84 (1.64 to 2.04)	<0.001†
Intrapersonal factors				
CESD-Depressive symptoms, M (95% CI)	9.91 (9.55 to 10.27)	9.69 (9.32 to 10.07)	12.19 (10.84 to 13.55)	<0.001†
TCI-Impulsivity, M (95% CI)	7.99 (7.87 to 8.10)	8.08 (7.97 to 8.20)	6.97 (6.55 to 7.39)	<0.001†
Ever substance use, n (%)	757 (34.6)	617 (30.9)	140 (73.3)	<0.001*
Delinquent behaviour, M (95% CI)	14.39 (14.23 to 14.55)	14.13 (13.97 to 14.29)	17.08 (16.32 to 17.85)	<0.001†
Smoking susceptibility, M (95% CI)	3.47 (3.43 to 3.51)	3.40 (3.36 to 3.44)	4.20 (4.00 to 4.41)	<0.001†
Smoking expectancies, M (95% CI)	6.90 (6.85 to 6.95)	6.93 (6.88 to 6.98)	6.53 (6.36 to 6.71)	<0.001†

* χ^2 test.

†Independent samples t-test.

‡Spearman's r test.

CESD, Center for Epidemiologic Studies Depression Scale; M, mean; TCI, Temperament and Character Inventory.

time (5–7 days)). Impulsivity was assessed with Temperament and Character Inventory⁵² Impulsivity subscale sum score (eg, 'I often do things based on how I feel at the moment'; range: 0–5). Lifetime use of other (non-nicotine or tobacco) substances was assessed with the MTF and YRBS items on ever use (yes/no) of alcohol and 13 other substances. Delinquent behaviour⁵³ was assessed by calculating a mean of frequency ratings for engaging in 11 different behaviours (eg, stealing, lying to parents; 1=Never to 6=Ten or more times) in the past 6 months. Susceptibility to smoking was measured by using a mean of a three-item index⁴⁴ (eg, 'Would you try smoking a cigarette if one of your best friends offered it to you?' (1=Definitely not, 4=Definitely yes)). Smoking outcome expectancies were assessed by a mean of the two responses for 'I think I might enjoy (...) smoking'

and (reversed) 'I think I might feel bad (...) from smoking' (1=Strongly disagree, 4=Strongly agree).⁵⁴

Data analysis

Following the protocol of Leventhal *et al.*²¹ prevalence and associations of lifetime (ever vs never) e-cigarette use and lifetime (ever vs never) combustible tobacco use in the overall baseline sample were first analysed. Subsequently, study attrition and descriptive statistics in the sample of baseline never smokers were reported. For the primary analyses, separate generalised linear mixed models⁵⁵ for each outcome were used with the follow-up data at two time points (6 and 12 months) as repeated measurements. For research question 1, each binary outcome (eg, any lifetime combustible tobacco product, cigarettes, cigars, hookah)

was analysed with logistic mixed regression, whereas the ‘number of combustible products’ outcome was analysed with ordinal mixed regression. All models included baseline e-cigarette ever use, school and time (6-month vs 12-month follow-up) as fixed effects and were fit with and without adjustment for all covariates. To examine whether the association between baseline e-cigarette use and combustible tobacco use differed across follow-ups, the baseline ‘e-cigarette \times time’ interaction term was added to each model. If this interaction was significant, the association between e-cigarette use and the outcome was examined separately for the 6 and 12-month follow-ups, including differences in outcome for ever and never use of e-cigarettes at baseline. If not significant, the interaction term was removed from the model, and associations between baseline e-cigarette use and the outcome were averaged across both follow-ups. Participants with missing data on baseline e-cigarette use or baseline measurements of the outcome variable were not included. For research question 2, an ordinal logistic mixed regression model was used to determine the association between baseline frequency of e-cigarette use and follow-up frequency of smoking, fit with and without adjustment for covariates, but always adjusting for time, school and baseline smoking frequency.²² Finally, for research question 3, on the association between baseline combustible tobacco ever use and ever use of e-cigarettes at follow-ups among never users of e-cigarettes at baseline, a similar analysis protocol as research question 1 was used: a binary logistic mixed regression model predicting e-cigarette use from baseline ever tobacco use status (with and without adjustment for covariates).

To account for missing data, multiple imputation was done with an imputation model that was identical to the analysis model including the covariates. The fully conditional specification method was used,⁵⁶ which has been shown to yield unbiased parameter estimates and SEs.^{57,58} To obtain unbiased estimates of the relations between the predictor variables and the outcomes, missing data on baseline covariates and on the outcome variables were imputed.⁵⁹ An analysis done on imputed data sets introduces power loss and also some uncertainty concerning the p value. By taking the number of imputations at least as large as the proportion of incomplete cases, 80 imputations in the present analysis, this is estimated to yield an acceptable power loss⁵⁹ and a Monte Carlo SE of the p value 0.05 less than 0.01.⁵⁵ Although analysis with multiply imputed data is valid under the assumption of missingness at random,⁵⁶ this assumption cannot be tested. Therefore, for research question 1, a sensitivity analysis was performed using

imputations according to a scenario violating this assumption. We chose to impute according to a pessimistic scenario, in that it decreased the association between baseline e-cigarette use and an outcome, by imputing the smoking outcome at follow-up in the same way for both baseline ever and never e-cigarette users. At each follow-up, the highest school-specific incidences of each outcome were taken as smoking probabilities for the imputation. Significance was set to 0.05 and all tests were two tailed. All primary analyses were conducted in Mplus V7.2.

Role of funding source

The funder had no role in the study design, the collection or analysis of the data, the interpretation of data, the writing of the report or the decision to submit the article for publication.

RESULTS

Study sample

All students in the recruited schools were eligible to participate and enrolled (n=2845). Data were collected for 2839 participants at baseline, including 2185 participants who never smoked tobacco at baseline, of which 1276 (58%) and 1025 (47%) participants completed the 6 and 12-month follow-ups, respectively. The analytical sample available for analyses across waves for research question 1 is depicted in online supplemental figure 1.

Descriptive analyses

Table 1 shows that boys were more likely than girls to have ever used e-cigarettes. Furthermore, age, highest parental education and all environmental and intrapersonal factors showed significant associations with e-cigarette use at baseline.

Table 2 shows that in the combined sample of ever and never smokers, baseline e-cigarette use was positively associated with baseline use (ever) of combustible tobacco products. Prevalence of combustible tobacco products was between 5.6% (cigars) and 17.8% (cigarettes). 21.6% (n=603) of participants reported ever use of e-cigarettes; 14.7% (n=412) ever use of e-cigarettes as well as combustible tobacco; 7.1% (n=197) combustible tobacco only; and 71.4% (n=1994) never used e-cigarettes nor combustible tobacco.

Associations between baseline e-cigarette ever use and combustible tobacco ever use at follow-ups in baseline never smokers

Table 3 shows that, among never smokers, baseline ever e-cigarette users were more likely than never users to have used

Table 2 Prevalence and cross-sectional association of baseline e-cigarette use and combustible tobacco use in combined sample of baseline ever smokers and never smokers

Baseline ever use	Overall (n=2794)	Baseline e-cigarette never users (n=2191)	Baseline e-cigarette ever users (n=603)	Difference in prevalence rates	
				% (95% CI)	P value
Any combustible tobacco product, n (%)	609 (21.8)	197 (9.0)	412 (68.3)	59.3 (55.4 to 63.2)	<0.001*
Combustible cigarettes, n (%)	496 (17.8)	151 (6.9)	345 (57.2)	50.3 (46.2 to 54.4)	<0.001*
Cigars, n (%)†	157 (5.6)	28 (1.3)	129 (21.4)	20.1 (16.8 to 23.4)	<0.001*
Hookah, n (%)‡	339 (12.2)	81 (3.7)	258 (42.8)	34.1 (30.0 to 38.2)	<0.001*
Number of different combustible tobacco products, n (%)					<0.001*
0	2185 (78.2)	1994 (91.0)	191 (31.7)		
1	324 (11.6)	142 (6.5)	182 (30.2)		
2	187 (6.7)	47 (2.1)	140 (23.2)		
3	98 (3.5)	8 (0.4)	90 (14.9)		

* χ^2 test.

†Due to missing data for each respective variable, denominators are n=2793.

‡Due to missing data for each respective variable, denominators are n=2790.

Table 3 Prevalence of past 6-month combustible tobacco use at 6 and 12-month follow-ups by baseline e-cigarette ever use among baseline never smokers

Use in the prior 6 months	Baseline e-cigarette use			Difference in prevalence rates	
	Overall (n=2185)	Never use (n=1994)	Ever use (n=191)	% (95% CI)	P value
6-month follow-up					
Any combustible tobacco product, n (%) [*]	90 (7.1)	64 (5.5)	26 (23.2)	17.7 (9.8 to 25.6)	<0.001
Combustible cigarettes, n (%) [†]	63 (4.9)	43 (3.7)	20 (17.9)	14.2 (7.0 to 21.4)	<0.001
Cigars, n (%) [‡]	15 (1.2)	11 (0.9)	4 (3.6)	2.7 (−0.8 to 6.2)	0.014
Hookah, n (%) [§]	33 (2.6)	22 (1.9)	11 (9.9)	8.0 (2.4 to 13.6)	<0.001
Number of different combustible tobacco products, n (%) [†]					<0.001
0	1185 (92.9)	1099 (94.5)	86 (76.8)		
1	72 (5.6)	55 (4.7)	17 (15.2)		
2	15 (1.2)	6 (0.5)	9 (8.0)		
3	3 (0.2)	3 (0.3)	0 (0.0)		
12-month follow-up					
Any combustible tobacco product, n (%) [¶]	141 (13.8)	101 (10.8)	40 (44.4)	33.6 (23.1 to 44.1)	<0.001
Combustible cigarettes, n (%) [¶]	118 (11.5)	85 (9.1)	33 (36.7)	27.6 (17.5 to 37.7)	<0.001
Cigars, n (%) ^{**}	18 (1.8)	14 (1.5)	4 (4.4)	2.9 (−1.4 to 7.2)	0.042
Hookah, n (%) ^{**}	47 (4.6)	35 (3.8)	12 (13.3)	9.5 (2.3 to 16.6)	<0.001
Number of different combustible tobacco products, n (%) [¶]					<0.001
0	884 (86.2)	834 (89.2)	50 (55.6)		
1	106 (10.3)	74 (7.9)	32 (35.6)		
2	28 (2.7)	21 (2.2)	7 (7.8)		
3	7 (0.7)	6 (0.6)	1 (1.1)		
*n=1276. †n=1275. ‡n=1273. §n=1270. ¶n=1025. **n=1023.					

combustible tobacco at 6-month (23.2% vs 5.5%; % difference=17.7; 95% CI 9.8 to 25.6) and 12-month (44.4% vs 10.8%; % difference=33.6; 95% CI 23.1 to 44.1) follow-ups. **Table 4** displays the results after imputation. Interaction of e-cigarette use (ever) with time was significant (OR=3.03; 95% CI 1.39 to 6.63) in the unadjusted analysis. Separate analyses at both follow-ups indicated a stronger association for baseline e-cigarette use and combustible tobacco use at the 12-month follow-up (OR=11.39; 95% CI 6.04 to 21.50), compared with the 6-month follow-up (OR=3.76; 95% CI 2.12 to 6.65). Analysis for time (of follow-up) was significant for both e-cigarette ever and never users, indicating an increase in the use of combustible tobacco across time in both groups. In the adjusted model, the interaction between baseline e-cigarette use and time was also significant for any combustible tobacco product use (OR=2.99; 95% CI 1.37 to 6.50). Furthermore, baseline e-cigarette use was associated more with any combustible tobacco use at 12-month follow-up (OR=5.63; 95% CI 3.04 to 10.42), compared with the 6-month follow-up (OR=1.89; 95% CI 1.05 to 3.37).

Table 3 also shows that, among never smokers, e-cigarette use (ever vs never) at baseline was positively associated with higher odds of smoking cigarettes, cigars and hookah. In the unadjusted analyses in **table 4**, e-cigarette use (ever vs never) at baseline was also positively associated with each of these outcomes at both follow-ups. In the adjusted model, baseline ever e-cigarette use was associated with cigarette use at the 12-month follow-up and with number of tobacco products at both follow-ups. Interaction with time was not significant for hookah use and cigar use, thus associations of baseline ever use of e-cigarettes with these outcomes averaged across time were examined. The relation between e-cigarette use at baseline and hookah use (averaged

over 6 and 12-month follow-ups) was significant (OR=3.69; 95% CI 1.75 to 7.77). Of the intrapersonal factors, only smoking susceptibility was positively associated with combustible tobacco use at both follow-ups (see **table 4**).

Additional sensitivity analyses were performed (see online supplemental table 3). Findings were consistent in the adjusted models for any tobacco use, combustible cigarette use, cigar use and number of tobacco products. For hookah use, there was a significant interaction of e-cigarette use and time. Only the association between e-cigarette use and hookah use at 6-month follow-up remained significant.

Association between baseline e-cigarette use frequency and tobacco use frequency at follow-ups

Online supplemental table 1 shows that, for the unadjusted model, higher scores on the four-level baseline e-cigarette use frequency variable were associated with greater odds of higher smoking frequency averaged across both follow-ups (OR=2.11; 95% CI 1.69 to 2.64). After adjusting for covariates, this association remained significant (OR=1.63; 95% CI 1.29 to 2.06). In the unadjusted analysis, the positive association between baseline e-cigarette use and follow-up smoking frequency differed between the different baseline smoking groups (OR=0.70; 95% CI 0.52 to 0.94, p=0.02), the association being stronger among baseline non-smokers (n=2180; OR=2.53; 95% CI 1.99 to 3.23) than baseline infrequent (smoked 1–2 in the past 30 days; n=41; OR=1.84; 95% CI 1.46 to 2.31) and frequent (smoked ≥3 in the past 30 days; n=127; OR=1.33; 95% CI 0.89 to 1.99) smokers (not shown in online supplemental table 1).

Table 4 Association of baseline e-cigarette ever use and covariates to combustible tobacco use outcomes at 6 and 12-month follow-ups among baseline never smokers

	Outcome									
	Any tobacco product		Combustible cigarettes		Cigars		Hookah		Number of tobacco products	
	OR (95% CI)*	P value	OR (95% CI)*	P value	OR (95% CI)*	P value	OR (95% CI)*	P value	OR (95% CI)†	P value
Unadjusted models										
E-cigarette ever (vs never) use										
Smoking outcome at 6 months	3.76 (2.12 to 6.65)	<0.001	4.00 (2.07 to 7.71)	<0.001	3.99 (1.62 to 9.86)	0.003	5.52 (2.74 to 11.12)	<0.001	3.90 (2.22 to 6.85)	<0.001
Smoking outcome at 12 months	11.39 (6.04 to 21.50)	<0.001	9.70 (5.12 to 18.37)	<0.001	3.99 (1.62 to 9.86)	0.003	5.52 (2.74 to 11.12)	<0.001	8.40 (4.64 to 15.21)	<0.001
Time (12 vs 6 months): e-cigarette never	1.98 (1.41 to 2.78)	<0.001	2.43 (1.60 to 3.70)	<0.001	1.53 (0.73 to 3.17)	0.26	2.01 (1.20 to 3.36)	0.008	2.00 (1.43 to 2.82)	<0.001
Time (12 vs 6 months): e-cigarette ever	6.00 (2.98 to 12.11)	<0.001	5.89 (2.79 to 12.44)	<0.001	1.53 (0.73 to 3.17)	0.26	2.01 (1.20 to 3.36)	0.008	4.32 (2.15 to 8.70)	<0.001
Ever e-cigarette use × time‡	3.03 (1.39 to 6.63)	0.006	2.43 (1.07 to 5.48)	0.033	1.44 (0.25 to 8.47)	0.69	1.59 (0.48 to 5.24)	0.45	2.15 (1.01 to 4.58)	0.046
Adjusted models										
Categorical covariates										
Girls (vs boys)	0.94 (0.68 to 1.30)	0.73	0.93 (0.65 to 1.33)	0.69	0.65 (0.31 to 1.38)	0.27	0.90 (0.53 to 1.54)	0.71	0.92 (0.67 to 1.27)	0.63
Dutch (vs other) ethnicity	0.42 (0.22 to 0.79)	0.007	0.65 (0.31 to 1.37)	0.26	0.55 (0.16 to 1.88)	0.34	0.22 (0.08 to 0.55)	0.001	0.42 (0.23 to 0.78)	0.006
Lives with both parents (vs other)	0.86 (0.59 to 1.25)	0.42	0.92 (0.63 to 1.36)	0.69	0.75 (0.35 to 1.63)	0.47	0.95 (0.51 to 1.75)	0.86	0.88 (0.62 to 1.26)	0.48
Substance ever (vs never) use	1.73 (1.21 to 2.47)	0.002	1.74 (1.16 to 2.61)	0.007	1.62 (0.74 to 3.56)	0.23	1.54 (0.84 to 2.83)	0.16	1.68 (1.18 to 2.40)	0.004
Family history of smoking (yes vs no)	0.93 (0.67 to 1.29)	0.66	0.93 (0.64 to 1.34)	0.68	0.61 (0.28 to 1.31)	0.20	0.86 (0.49 to 1.50)	0.59	0.93 (0.66 to 1.32)	0.69
Continuous covariates§										
Age	1.13 (0.96 to 1.34)	0.14	1.18 (0.97 to 1.42)	0.10	1.10 (0.75 to 1.63)	0.62	1.08 (0.81 to 1.43)	0.51	1.13 (0.96 to 1.34)	0.15
Parental education	1.00 (0.85 to 1.19)	0.96	1.04 (0.88 to 1.23)	0.65	0.78 (0.54 to 1.12)	0.17	1.10 (0.85 to 1.43)	0.46	1.01 (0.86 to 1.17)	0.95
Peer smoking	1.11 (0.96 to 1.27)	0.17	1.09 (0.94 to 1.25)	0.26	1.03 (0.77 to 1.37)	0.85	1.12 (0.91 to 1.39)	0.28	1.09 (0.96 to 1.25)	0.17
CESD-Depressive symptoms	1.06 (0.91 to 1.24)	0.47	1.09 (0.91 to 1.29)	0.35	1.05 (0.75 to 1.45)	0.79	1.00 (0.80 to 1.25)	1.00	1.07 (0.92 to 1.24)	0.36
TCI-Impulsivity	0.90 (0.77 to 1.06)	0.21	0.90 (0.76 to 1.06)	0.20	0.70 (0.47 to 1.03)	0.07	0.93 (0.73 to 1.17)	0.53	0.89 (0.76 to 1.04)	0.14
Delinquent behaviour	1.03 (0.89 to 1.20)	0.65	1.04 (0.88 to 1.22)	0.66	1.18 (0.93 to 1.50)	0.17	1.15 (0.92 to 1.44)	0.23	1.05 (0.91 to 1.21)	0.53
Smoking susceptibility	1.27 (1.10 to 1.46)	0.001	1.31 (1.13 to 1.52)	<0.001	1.17 (0.85 to 1.62)	0.33	0.91 (0.71 to 1.17)	0.47	1.22 (1.07 to 1.39)	0.002
Smoking expectancies	0.96 (0.81 to 1.13)	0.62	0.90 (0.76 to 1.07)	0.22	1.14 (0.75 to 1.74)	0.54	1.06 (0.81 to 1.39)	0.68	0.97 (0.82 to 1.13)	0.67
Regressors										
E-cigarette ever (vs never) use										
Smoking outcome at 6 months	1.89 (1.05 to 3.37)	0.032	1.93 (0.98 to 3.79)	0.058	1.92 (0.75 to 4.93)	0.18	3.69 (1.75 to 7.77)	0.001	2.04 (1.16 to 3.60)	0.0124
Smoking outcome at 12 months	5.63 (3.04 to 10.42)	<0.001	4.58 (2.42 to 8.68)	<0.001					4.32 (2.40 to 7.76)	<0.001
Time (12 vs 6 months): e-cigarette never	1.98 (1.41 to 2.79)	<0.001	2.44 (1.60 to 3.72)	<0.001	1.53 (0.73 to 3.17)	0.26	2.01 (1.20 to 3.35)	0.008	2.01 (1.43 to 2.83)	<0.001
Time (12 vs 6 months): e-cigarette ever	5.92 (2.95 to 11.87)	<0.001	5.82 (2.77 to 12.19)	<0.001					4.25 (2.12 to 8.55)	<0.001
Ever e-cigarette use × time‡	2.99 (1.37 to 6.50)	0.006	2.38 (1.06 to 5.37)	0.036	1.43 (0.25 to 8.36)	0.69	1.58 (0.48 to 5.20)	0.45	2.12 (1.00 to 4.49)	0.051

All analyses include only never users of combustible tobacco products at baseline (n=2185).

*OR from repeated binary logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects.

†OR from repeated ordinal logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects, with the OR expressing the change in odds of being in a category with use of a certain amount of tobacco products versus being in a category with lower use (3 vs ≤2, ≥2 vs ≤1, ≥1 vs 0).

‡If interaction term is significant (p≤0.05) or marginally significant (p≤0.10), the effect of e-cigarette use is examined both at 6 and 12-month follow-ups, and the effect of time is examined both for never and ever users of e-cigarettes. If interaction term is not (marginally) significant, the effect of e-cigarette use is examined averaged across and is thus the same for both the 6 and 12-month follow-ups, and the effect of time is examined averaged across and is thus also the same for never and ever users of e-cigarettes.

§Continuous covariates rescaled (M=0, SD=1), such that the ORs indicate change in odds in the outcome associated with an increase in one SD unit on the covariate continuous scale.

CESD, Center for Epidemiologic Studies Depression Scale; TCI, Temperament and Character Inventory.

Association between baseline ever smoking and follow-up ever e-cigarette use at follow-ups in baseline never e-cigarette users

Online supplemental table 2 shows that among never users of e-cigarettes, smoking at baseline was positively associated with higher odds of using e-cigarettes averaged across both follow-ups for the unadjusted (OR=5.22; 95% CI 3.06 to 8.92) and adjusted (OR=3.10; 95% CI 1.58 to 6.06) analyses.

DISCUSSION

The current replication study confirms that e-cigarette use by non-smoking adolescents is associated with increased odds of subsequent combustible tobacco smoking initiation; that more frequent e-cigarette use is associated with more frequent subsequent tobacco smoking; and that the 'reverse' association applies, namely that tobacco smoking among never users of e-cigarettes is associated with greater odds of later e-cigarette use. These findings are consistent with previous studies,^{21 22 30 31} employ various adjusted analyses and sensitivity analyses and extend the findings from the US to a European context.

Collectively, this suggests that e-cigarettes may indeed act as a gateway to tobacco smoking for youth, as we see that youth who used e-cigarettes are more likely to try smoking tobacco later. We also found a dose-related relation, namely that the higher the frequency of e-cigarette use was at baseline, the higher the frequency of subsequent tobacco use was at follow-up. This may indicate a causal relation for using both products. Additionally, our analyses suggest a 'reversed' gateway of tobacco smoking to e-cigarette use. Yet, these gateways may operate through different mechanisms. E-cigarettes may be a smoother introduction to tobacco smoking⁶⁰; and reversely, the transition from tobacco smoking to e-cigarette use could point at transitioning to a less harmful behaviour. However, both behaviours may also stem from common risk factors, such as increased propensity to experiment with substances,^{14 61–63} where final choices on which specific substance is used first may depend on personal preferences, circumstances or cultural norms.³² Dual users of e-cigarettes and cigarettes, for example, share similar characteristics (eg, impulsivity) regardless of which product is used first.⁶⁴

It remains uncertain whether the found associations are causal, which the gateway hypothesis suggests, or an indication of shared risk factors for e-cigarette and tobacco use. While multiple predictors linked to tobacco smoking were addressed in our study, it remains uncertain whether all common liability risks were controlled for. Studies show that at least some youth with low propensity for tobacco smoking, nevertheless, used e-cigarettes.^{24 65 66} Another study found that relative to a propensity-matched control group without initial e-cigarette use, non-smoking adolescent e-cigarette users were less likely to become established smokers (30-day use and 100+ lifetime cigarettes).⁶⁷ This suggests that e-cigarettes do not function as a gateway to tobacco for everyone. Additionally, a study found that the relationship between e-cigarette use and subsequent smoking among adolescents may be weakened through interventions.⁶⁸ Likely, multiple mechanisms are complementary and the relation between causes and outcomes is complex and multidirectional.⁶⁹

A critique on quantitative studies on the gateway from e-cigarette to tobacco smoking is that the key question of 'why' is not addressed.⁶³ A popular explanation is that the nicotine in e-cigarettes makes individuals dependent and this may cause them to try combustible cigarettes.^{70 71} The authors of the original gateway theory even describe nicotine as a gateway drug that primes the brain for other substance use, 'whether the exposure

is from smoking tobacco, passive tobacco smoke, or e-cigarettes.'⁷² A qualitative study of youth who use(d) e-cigarettes found that e-cigarettes can be a 'smoother' introduction to the concept of smoking and that they appear to remove boundaries to smoking: 'There used to be a barrier that said either you're a smoker or a non-smoker, now I can smoke without smoking'.⁶⁰ Consequently, e-cigarette users get used to the acts and gestures of smoking which facilitates the transition to tobacco smoking. However, information in this area is still limited and more investigation is needed, including studies among tobacco smokers who started with e-cigarettes to assess reasons for taking up smoking.

Evidence and understanding of associations between e-cigarettes and tobacco and the long-term consequences of e-cigarette use among youth are still limited,²⁸ partly because of the relative novelty of e-cigarettes and still evolving technologies. Until this has been resolved, policymakers should carefully consider whether to act on the dangers (eg, a gateway to tobacco) or rather the benefits (eg, a potentially less harmful alternative to tobacco smoking) of e-cigarettes for smokers. Given the wide availability and marketing of e-cigarettes^{73 74} and that available evidence provides reasons for caution, prevention of e-cigarette use among non-smoking youth is recommended.

Limitations

The original and current study used binary and categorical outcome measures of smoking. These measures are limited, as smoking intensity (how many cigarettes per day) was not assessed and a fairly low cut-off point for frequent smoking (≥ 3 days during the past 30 days) was used because of the limited number of participants. Also, the majority (68.3%) of ever users of e-cigarettes had already smoked tobacco at baseline, which may mean that the current study sample is already too old for the main research question focusing on never users of tobacco. Furthermore, substance use was included as a dichotomous variable, meaning, for example, that students who drank alcohol and used cannabis on a weekly basis got the same score as students who had one drink in their lifetime. Additionally, the response of participating (vs invited) schools was low, which may have resulted in selection bias. However, relevant differences between students from participating schools and students at non-participating schools are not expected, as the response rate of students who were present at the time of the survey in participating school classes was 100%. Furthermore, the dropout rate from baseline to follow-up was high and may have affected the relations between e-cigarette use and tobacco use. However, the imputation of missing outcomes for research question 1 according to a pessimistic scenario (see the Data analysis section) yielded the same relations, indicating robustness of the results. Lastly, we coded Flemish (Belgian) students as Dutch in the main analyses because they are part of the (native) ethnic majority of their schools. Thus, differences between Dutch and Belgian students were not assessed, even though (cultural) differences may exist. However, preliminary analyses, in which Flemish students were coded as 'not Dutch', yielded similar results in the main analyses.

CONCLUSION

For this study, we replicated an American cornerstone study^{21 22} on the association between e-cigarette use and tobacco smoking in the Netherlands and Flanders and found similar results. High school students in the Netherlands and Flanders who used e-cigarettes at baseline were more likely to report initiation of combustible tobacco use over the next year compared with never

WHAT IS ALREADY KNOWN ON THIS SUBJECT

- ⇒ E-cigarette use is associated with subsequent tobacco smoking among youth.
- ⇒ A popular explanation of this relation is that e-cigarettes function as a gateway to tobacco smoking for youth.

WHAT IMPORTANT GAPS IN KNOWLEDGE EXIST ON THIS TOPIC

- ⇒ Most studies only investigate the relation between e-cigarette use and tobacco smoking in one direction. Less is known about the opposite relation between tobacco smoking and subsequent e-cigarette use.
- ⇒ It is not yet determined whether the gateway hypothesis or the common liability hypothesis best explains the relation between e-cigarette use and tobacco smoking.

WHAT THIS PAPER ADDS

- ⇒ This study replicated the findings from an American cornerstone study on the relationship between e-cigarette use and tobacco use. Findings indicate both the relationship between e-cigarette use and subsequent tobacco smoking and the opposite relation.
- ⇒ The association between e-cigarette use and tobacco smoking is likely bidirectional for adolescents.

users of e-cigarettes. These findings add to a growing body of studies that indicate a link between e-cigarette use and tobacco smoking in youth. The gateway hypothesis was further explored by also analysing the 'reverse' relation between baseline tobacco smoking and subsequent e-cigarette use. A similar association was found which may indicate that the gateway works in two directions, that e-cigarette use and tobacco smoking share common risk factors, or that both mechanisms apply. Different types of studies are needed to better understand *why* there are such associations and whether these may be causal relationships.

Contributors TM: writing—original draft, validation, formal analysis, data curation. MJMC: formal analysis, validation, writing—original draft. HdV, RT, CPvS: conceptualisation, methodology, writing—review and editing. VK: investigation, project administration, resources. GEN: conceptualisation, methodology, funding acquisition, writing—review and editing, supervision.

Funding This study was funded by NWO (401.16.012).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The Maastricht University Ethical Review Board approved the study (2018-0418).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Anonymised data and protocols are available upon reasonable request from the corresponding author (martinelli@ivo.nl).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given,

and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD


Thomas Martinelli <http://orcid.org/0000-0003-0055-5631>

REFERENCES

- Hajek P, Phillips-Waller A, Przulj D, *et al*. A randomized trial of e-cigarettes versus Nicotine-Replacement therapy. *N Engl J Med* 2019;380:629–37.
- McNeill A, Brose LS, Calder R. E-cigarettes : an evidence update A report commissioned by Public Health England, 2015. Available: www.gov.uk/government/uploads/system/uploads/attachment_data/file/454516/E-cigarettes_an_evidence_update_A_report_commissioned_by_Public_Health_England.pdf
- Levy DT, Borland R, Lindblom EN, *et al*. Potential deaths averted in USA by replacing cigarettes with e-cigarettes. *Tob Control* 2018;27:18–25.
- Abrams DB. Promise and peril of e-cigarettes. *JAMA* 2014;311:135.
- England LJ, Bunnell RE, Pechacek TF, *et al*. Nicotine and the developing human: a neglected element in the electronic cigarette debate. *Am J Prev Med* 2015;49:286–93.
- Tomar SL, Alpert HR, Connolly GN. Patterns of dual use of cigarettes and smokeless tobacco among US males: findings from national surveys. *Tob Control* 2010;19:104–9.
- Hanewinkel R, Isensee B. Risk factors for e-cigarette, conventional cigarette, and dual use in German adolescents: a cohort study. *Prev Med* 2015;74:59–62.
- Hajek P, Etter J-F, Benowitz N, *et al*. Electronic cigarettes: review of use, content, safety, effects on smokers and potential for harm and benefit. *Addiction* 2014;109:1801–10.
- Soneji S, Barrington-Trimis JL, Wills TA, *et al*. Association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adults: a systematic review and meta-analysis. *JAMA Pediatr* 2017;171:788–97.
- Kandel D. Stages in adolescent involvement in drug use. *Science* 1975;190:912–4.
- Kandel DB, Yamaguchi K, Klein LC. Testing the gateway hypothesis. *Addiction* 2006;101:470–2.
- Vanyukov MM, Tarter RE, Kirillova GP, *et al*. Common liability to addiction and "gateway hypothesis": theoretical, empirical and evolutionary perspective. *Drug Alcohol Depend* 2012;123:53–17.
- Hall WD, Lynskey M. Is cannabis a gateway drug? testing hypotheses about the relationship between cannabis use and the use of other illicit drugs. *Drug Alcohol Rev* 2005;24:39–48.
- Etter J-F. Gateway effects and electronic cigarettes. *Addiction* 2018;113:1776–83.
- Bell K, Keane H. All gates lead to smoking: the 'gateway theory', e-cigarettes and the remaking of nicotine. *Soc Sci Med* 2014;119:45–52.
- European Union. *Directive 2014/40/EU of the European Parliament and of the Council of 3 April 2014 on the approximation of the laws, regulations and administrative provisions of the member states concerning the manufacture, presentation and sale of tobacco and related PR*, 2014.
- Conner M, Grogan S, Simms-Ellis R, *et al*. Do electronic cigarettes increase cigarette smoking in UK adolescents? Evidence from a 12-month prospective study. *Tob Control* 2018;27:365–72.
- Levy DT, Warner KE, Cummings KM, *et al*. Examining the relationship of vaping to smoking initiation among US youth and young adults: a reality check. *Tob Control* 2019;28:629–35.
- Van Dorsselaer S, Tuithof M, Verdurmen J. Jeugd en riskant gedrag, 2020. Available: <https://www.trimbos.nl/docs/29856421-c89c-47f8-ac04-c7ce1cabad4b.pdf>
- Morgenstern M, Nies A, Goecke M, *et al*. E-Cigarettes and the use of conventional cigarettes. *Dtsch Arztebl Int* 2018;115:243–8.
- Leventhal AM, Strong DR, Kirkpatrick MG, *et al*. Association of electronic cigarette use with initiation of Combustible tobacco product smoking in early adolescence. *JAMA* 2015;314:700.
- Leventhal AM, Stone MD, Andrabi N, *et al*. Association of e-cigarette vaping and progression to heavier patterns of cigarette smoking. *JAMA* 2016;316:1918–20.
- Food and Drug Administration. *Deeming tobacco products to be subject to the federal food, drug, and cosmetic act, as amended by the family smoking prevention and tobacco control act; restrictions on the sale and distribution of tobacco products and required warning statements for Toba*. Federal Register, 2016.
- Barrington-Trimis JL, Urman R, Berhane K, *et al*. E-Cigarettes and future cigarette use. *Pediatrics* 2016;138. doi:10.1542/peds.2016-0379. [Epub ahead of print: 13 Jun 2016].
- Spindle TR, Hiler MM, Cooke ME, *et al*. Electronic cigarette use and uptake of cigarette smoking: a longitudinal examination of U.S. college students. *Addict Behav* 2017;67:66–72.
- Wills TA, Knight R, Sargent JD, *et al*. Longitudinal study of e-cigarette use and onset of cigarette smoking among high school students in Hawaii. *Tob Control* 2017;26:34–9.
- Best C, Haseen F, Currie D, *et al*. Relationship between trying an electronic cigarette and subsequent cigarette experimentation in Scottish adolescents: a cohort study. *Tob Control* 2018;27:373–8.

- 28 Khouja JN, Suddell SF, Peters SE, *et al.* Is e-cigarette use in non-smoking young adults associated with later smoking? A systematic review and meta-analysis. *Tob Control* 2020;8–15.
- 29 Treur JL, Rozema AD, Mathijssen JJP, *et al.* E-Cigarette and waterpipe use in two adolescent cohorts: cross-sectional and longitudinal associations with conventional cigarette smoking. *Eur J Epidemiol* 2018;33:323–34.
- 30 East K, Hitchman SC, Bakolis I, *et al.* The association between smoking and electronic cigarette use in a cohort of young people. *J Adolesc Heal* 2018;62:539–47.
- 31 Morello P, Pérez A, Peña L, *et al.* Prevalence and predictors of e-cigarette trial among adolescents in Argentina. *Tob Prev Cessat* 2016;2.
- 32 Wetzels JLL, Kremers SPJ, Vitoria PD, *et al.* The alcohol-tobacco relationship: a prospective study among adolescents in six European countries. *Addiction* 2003;98:1755–63.
- 33 Open Science Collaboration. Estimating the reproducibility of psychological science. *Science* 2015.
- 34 Federale overheidsdienst Volksgezondheid Veiligheid van de voedselketen en Leefmilieu. Algemene regels voor e-sigaretten. KB 28/10/2016, 2016. Available: <https://www.health.belgium.be/nl/gezondheid/zorg-voor-jezelf/alcohol-tabak/e-sigaret#Wet> [Accessed 16 Jun 2020].
- 35 Ministerie van VWS. Tabaks- en rookwarenwet, 2016. Available: <https://zoek.officialiebekendmakingen.nl/stb-2016-176.html>
- 36 Czaplinski L, Kostygina G, Kim Y, *et al.* Characterising JUUL-related posts on Instagram. *Tob Control* 2020;29:tobaccocontrol-2018-054824.
- 37 US Food and Drug Administration. FDA warns JUUL Labs for marketing unauthorized modified risk tobacco products, including in outreach to youth, 2019. Available: <https://www.fda.gov/news-events/press-announcements/fda-warns-juul-labs-marketing-unauthorized-modified-risk-tobacco-products-including-outreach-youth>
- 38 Truth Initiative. E-Cigarettes: facts, STATS and regulations, 2021. Available: <https://truthinitiative.org/research-resources/emerging-tobacco-products/e-cigarettes-facts-stats-and-regulations>
- 39 Huang J, Duan Z, Kwok J, *et al.* Vaping versus JUULing: how the extraordinary growth and marketing of JUUL transformed the US retail e-cigarette market. *Tob Control* 2019;28:146–51.
- 40 Eaton DK, Kann L, Kinchen S, *et al.* Youth risk behavior surveillance - United States, 2009. *MMWR Surveill Summ* 2010;59:1–142.
- 41 Miech R, Patrick ME, O'Malley PM, *et al.* E-Cigarette use as a predictor of cigarette smoking: results from a 1-year follow-up of a national sample of 12th grade students. *Tob Control* 2017;26:e106–11.
- 42 Audrain-McGovern J, Rodriguez D, Tercyak KP, *et al.* Identifying and characterizing adolescent smoking trajectories. *Cancer Epidemiol Biomarkers Prev* 2004;13:2023–34.
- 43 Radloff LS. The use of the center for epidemiologic studies depression scale in adolescents and young adults. *J Youth Adolesc* 1991;20:149–66.
- 44 Strong DR, Hartman SJ, Nodora J, *et al.* Predictive validity of the expanded susceptibility to smoke index. *Nicotine Tob Res* 2015;17:862–9.
- 45 Camenga DR, Kong G, Cavallo DA, *et al.* Alternate tobacco product and drug use among adolescents who use electronic cigarettes, cigarettes only, and never smokers. *J Adolesc Health* 2014;55:588–91.
- 46 Covey LS, Tam D, mood D. Depressive mood, the single-parent home, and adolescent cigarette smoking. *Am J Public Health* 1990;80:1330–3.
- 47 Tyas SL, Pederson LL. Psychosocial factors related to adolescent smoking: a critical review of the literature. *Tob Control* 1998;7:409–20.
- 48 Cardenas VM, Breen PJ, Compadre CM, *et al.* The smoking habits of the family influence the uptake of e-cigarettes in US children. *Ann Epidemiol* 2015;25:60–2.
- 49 Wills TA, Knight R, Williams RJ, *et al.* Risk factors for exclusive e-cigarette use and dual e-cigarette use and tobacco use in adolescents. *Pediatrics* 2015;135:e43–51.
- 50 Kong G, Morean ME, Cavallo DA, *et al.* Reasons for electronic cigarette experimentation and discontinuation among adolescents and young adults. *Nicotine Tob Res* 2015;17:847–54.
- 51 CDC. *National youth tobacco survey methodology report*. Atlanta, GA, 2011.
- 52 Cloninger CR, Przybeck T, Syrakis D. *The temperament and character inventory (TCI): a guide to its development and use*. St. Louis, MO, 1994.
- 53 Thompson MP, Ho C-hua, Kingree JB. Prospective associations between delinquency and suicidal behaviors in a nationally representative sample. *J Adolesc Heal* 2007;40:232–7.
- 54 Katz EC, Fromme K, D'Amico EJ. Effects of outcome expectancies and personality on young adults' illicit drug use, heavy drinking, and risky sexual behavior. *Cognit Ther Res* 2000;24:1–22.
- 55 McCulloch CE, Neuhaus JM. Generalized Linear Mixed Models. In: *Wiley StatsRef: statistics reference online*. Chichester, UK: John Wiley & Sons, Ltd, 2014.
- 56 Enders C. *Applied missing data analysis*. New York: The Guilford Press, 2010.
- 57 Raghunathan T, Lepkowski J, Hoewyk J. A multivariate technique for multiply imputing missing values using a sequence of regression models. *Surv Methodol* 2001;27:85–96.
- 58 Van BS, Brand JPL, Rubin DB. Fully conditional specification in multivariate imputation. *J Stat Comput Simul* 2007.
- 59 Graham JW, Olchowski AE, Gilreath TD. How many imputations are really needed? some practical clarifications of multiple imputation theory. *Prev Sci* 2007;8:206–13.
- 60 Akre C, Suris J-C. Adolescents and young adults' perceptions of electronic cigarettes as a gateway to smoking: a qualitative study in Switzerland. *Health Educ Res* 2017;32:448–54.
- 61 Chapman S. Unravelling gossamer with boxing gloves: problems in explaining the decline in smoking. *BMJ* 1993;307:429–32.
- 62 Etter J-F. E-Cigarettes: methodological and ideological issues and research priorities. *BMC Med* 2015;13:32–3.
- 63 Kozlowski LT, Warner KE. Adolescents and e-cigarettes: objects of concern may appear larger than they are. *Drug Alcohol Depend* 2017;174:209–14.
- 64 Conner M, Grogan S, Simms-Ellis R, *et al.* Patterns and predictors of e-cigarette, cigarette and dual use uptake in UK adolescents: evidence from a 24-month prospective study. *Addiction* 2019;114:2048–55.
- 65 Dutra LM, Glantz SA. E-Cigarettes and national adolescent cigarette use: 2004-2014. *Pediatrics* 2017;139:e20162450.
- 66 Cerrai S, Potente R, Gorini G, *et al.* What is the face of new nicotine users? 2012-2018 e-cigarettes and tobacco use among young students in Italy. *Int J Drug Policy* 2020;86:102941.
- 67 Shahab L, Beard E, Brown J. Association of initial e-cigarette and other tobacco product use with subsequent cigarette smoking in adolescents: a cross-sectional, matched control study. *Tob Control* 2021;30:212–20.
- 68 Conner M, Grogan S, Simms-Ellis R, *et al.* Evidence that an intervention weakens the relationship between adolescent electronic cigarette use and tobacco smoking: a 24-month prospective study. *Tob Control* 2020;29:425–31.
- 69 Chapman S, Bareham D, Maziak W. The gateway effect of e-cigarettes: reflections on main criticisms. *Nicotine Tob Res* 2019;21:695–8.
- 70 Farsalinos KE, Spyrou A, Tsimopoulou K, *et al.* Nicotine absorption from electronic cigarette use: comparison between first and new-generation devices. *Sci Rep* 2015;4:4133.
- 71 Tomar SL. Is use of smokeless tobacco a risk factor for cigarette smoking? The U.S. experience. *Nicotine Tob Res* 2003;5:561–9.
- 72 Kandel ER, Kandel DB. Shattuck lecture. A molecular basis for nicotine as a gateway drug. *N Engl J Med* 2014;371:932–43.
- 73 Soule EK, Sakuma K-LK, Palafox S, *et al.* Content analysis of Internet marketing strategies used to promote flavored electronic cigarettes. *Addict Behav* 2019;91:128–35.
- 74 Garrison KA, O'Malley SS, Gueorguieva R, *et al.* A fMRI study on the impact of advertising for flavored e-cigarettes on susceptible young adults. *Drug Alcohol Depend* 2018;186:233–41.

Supplementary tables and figures:

Supplementary Figure 1: Flow of Adolescent Students in Study to Assess E-Cigarette Use at Baseline and Later Use of Combustible Tobacco Products

^a Includes all combustible tobacco products.

Supplementary Table 1: Association of Baseline e-cigarette use frequency with tobacco smoking frequency at 6- and 12-month Follow-up^a

	Parameter Estimate for Association with Smoking Frequency at follow-up	
	Odds ratio (95% CI)	P-value
Unadjusted model^b		
Baseline smoking (3-level continuum):		
- Outcome at 6 months	2.06 (1.41, 3.01)	< 0.001
- Outcome at 12 months	7.11 (4.03, 12.56)	< 0.001
E-cigarette use (4-level continuum)	2.11 (1.69, 2.64)	< 0.001
E-cigarette use x smoking ^c	0.70 (0.52, 0.94)	0.02
Adjusted model^d		
Baseline smoking (3-level continuum)		
- Outcome at 6 months	1.11 (0.71, 1.73)	0.66
- Outcome at 12 months	3.52 (2.02, 6.11)	<0.001
E-cigarette use (4-level continuum)	1.63 (1.29, 2.06)	< 0.001
E-cigarette use x smoking ^c	0.83 (0.61, 1.11)	0.21

^a Ordinal logistic mixed regression of proportional odds for being at a higher smoking frequency outcome (i.e., days smoked in the past 30 days; non-smoker, 0; infrequent smokers [1-2days], 1; frequent smokers [≥3 days], 2) accounting for clustering by pupil and school in a sample with available baseline e-cigarette use and smoking frequency data (N = 2348). The baseline 4-level continuous e-cigarette use variable was categorized as never, prior (ever with no past 30-day use), infrequent (1-2 days during past 30 days), or frequent (≥3 days during past 30 days). To address missing covariate data and missing outcomes, 80 multiple imputed datasets were generated using the fully conditional specification method (with sequential regression procedure). The parameter estimates from models in each imputed data set were pooled and presented as a single estimate.

^b Unadjusted models without e-cigarette use x smoking interaction include baseline e-cigarette use and smoking frequency, and interactions of these variables with time (6 versus 12 months follow-up). In case of a significant interaction with time, parameter estimates are shown separately for 6- and 12-months follow-up, otherwise the interaction term is removed from the model.

^c Interaction term added in subsequent model. If e-cigarette use x smoking x time is not significant, the results for e-cigarette use x smoking averaged across both follow-ups are presented.

^d Adjusted models without e-cigarette use x smoking interaction include baseline e-cigarette use and smoking frequency, and interactions of these variables with time (6 versus 12 months follow-up). In case of a significant interaction with time, parameter estimates are shown separately for 6- and 12-months follow-up, otherwise the interaction term is removed from the model. Adjustment is made for demographic, environmental and psychosocial covariates as described in the main text.

Supplementary Table 2: Association of Baseline Ever Use of Any Tobacco Product and Covariates to E-cigarette Use at 6- and 12-Month Follow Ups among Baseline Never E-Cigarette Users

Baseline regressors and covariates	Ever Use of E-Cigarettes	
	OR(95%CI) ^a	P
Unadjusted Models		
Any Tobacco Product ever (vs. never) use:		
E-Cigarette use at 6 months	5.22 (3.06,8.92)	< 0.001
E-Cigarette use at 12 months	5.22 (3.06,8.92)	< 0.001
Time (12- vs. 6-month): Any Tobacco use never	1.46 (1.01,2.12)	0.04
Time (12- vs. 6-month): Any Tobacco use ever	1.46 (1.01,2.12)	0.04
Ever Tobacco Product Use × Time ^b	1.92 (0.79,4.65)	0.15
Adjusted Models		
Categorical covariates		
Girls (vs. Boys)	0.83 (0.56,1.22)	0.33
Dutch (vs. other) ethnicity	0.82 (0.33,2.06)	0.67
Lives with both parents (vs. other)	0.90 (0.59,1.39)	0.64
Substance ever (vs. never) use	2.03 (1.30,3.16)	0.002
Family history of smoking (yes vs. no)	0.97 (0.65,1.45)	0.88
Continuous covariates ^c		
Age	1.02 (0.83,1.25)	0.84
Parental education	0.99 (0.83,1.18)	0.95
Peer smoking	0.98 (0.83,1.17)	0.86
CESD-Depressive Symptoms	0.92 (0.76,1.12)	0.47
TCI-Impulsivity	0.97 (0.81,1.16)	0.71
Delinquent Behaviour	1.11 (0.96,1.29)	0.17
Smoking susceptibility	1.04 (0.85,1.26)	0.71
Smoking expectancies	0.98 (0.79,1.20)	0.82
Regressors		
Any Tobacco Product ever (vs. never) use:		
E-Cigarette use at 6 months	3.10 (1.58,6.06)	0.001
E-Cigarette use at 12 months	3.10 (1.58,6.06)	0.001
Time (12- vs. 6-month): Any Tobacco use never	1.47 (1.01,2.12)	0.04
Time (12- vs. 6-month): Any Tobacco use ever	1.47 (1.01,2.12)	0.04
Ever Tobacco Product Use × Time ^b	1.91 (0.79,4.63)	0.15

Note: All analyses include only never-users of E-Cigarettes at baseline (N = 2191).

^a OR from repeated binary logistic regression model predicting E-cigarette use from baseline ever tobacco use status (yes/no) including school fixed effects.

^b If interaction term is significant (p-value ≤ 0.05), the effect of any tobacco product use is examined both at 6- and 12 months follow up, and the effect of time is examined both for never and ever users of any tobacco product. If interaction term is not significant, the effect of any tobacco use is examined averaged across and, is thus the same for both the 6- and 12 months follow ups, and the effect of time is examined averaged across and, is thus also the same for never and ever users of tobacco products.

^c Continuous covariates rescaled (M = 0, SD = 1), such that the ORs indicate change in odds in the outcome associated with an increase in one standard deviation unit on the covariate continuous scale. CESD = Center for Epidemiologic Studies Depression Scale. TCI = Temperament and Character Inventory.

Supplementary Table 3: Sensitivity analyses of the Association of Baseline E-Cigarette Ever Use and Covariates to Combustible Tobacco Use Outcomes at 6- and 12-Month Follow Ups among Baseline Never-Smokers

Baseline regressors and covariates	Outcome									
	Any tobacco product		Combustible cigarettes		Cigars		Hookah		Number of Tobacco Products	
	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^b	P
Unadjusted Models										
E-cigarette ever (vs. never) use:										
Smoking outcome on 6 months	2.33 (1.75,3.11)	<0.001	2.62 (1.85,3.71)	<0.001	1.43 (0.90,2.25)	0.13	3.00 (1.61,5.60)	0.001	2.30 (1.73, 3.08)	<0.001
Smoking outcome on 12 months	2.33 (1.75,3.11)	<0.001	2.62 (1.85,3.71)	<0.001	1.43 (0.90,2.25)	0.13	1.28 (0.71,2.32)	0.41	2.30 (1.73,3.08)	<0.001
Time (12- vs. 6-month): E-Cigarette never	2.77 (2.33,3.30)	<0.001	3.16 (2.55,3.91)	<0.001	2.32 (1.69,3.19)	<0.001	2.34 (1.73,3.18)	<0.001	2.80 (2.35,3.33)	<0.001
Time (12- vs. 6-month): E-Cigarette ever	2.77 (2.33,3.30)	<0.001	3.16 (2.55,3.91)	<0.001	2.32 (1.69,3.19)	<0.001	1.00 (0.49,2.06)	1.00	2.80 (2.35,3.33)	<0.001
Ever e-cigarette use × Time ^c	0.95 (0.57,1.57)	0.85	0.98 (0.56,1.73)	0.96	0.71 (0.26,1.92)	0.50	0.43 (0.20,0.93)	0.033	0.85 (0.52,1.39)	0.51
Adjusted Models										
Categorical covariates										
Girls (vs. Boys)	0.97 (0.81,1.16)	0.75	0.91 (0.73,1.15)	0.44	0.91 (0.66,1.25)	0.56	1.02 (0.76,1.38)	0.89	0.95 (0.80,1.14)	0.60
Dutch (vs. other) ethnicity	0.59 (0.38,0.91)	0.02	0.68 (0.41,1.12)	0.13	1.07 (0.48,2.42)	0.87	0.33 (0.17,0.65)	0.001	0.57 (0.36,0.90)	0.02
Lives with both parents (vs. other)	1.10 (0.88,1.37)	0.40	1.27 (0.96,1.67)	0.10	0.93 (0.65,1.34)	0.71	1.11 (0.76,1.61)	0.59	1.12 (0.90,1.40)	0.31
Substance ever (vs. never) use	1.28 (1.04,1.56)	0.02	1.21 (0.95,1.56)	0.13	1.47 (1.05,2.05)	0.03	1.39 (0.99,1.94)	0.06	1.30 (1.07,1.59)	0.01
Family history of smoking (yes vs. no)	1.12 (0.92,1.37)	0.24	1.05 (0.83,1.34)	0.68	0.98 (0.70,1.38)	0.93	1.11 (0.80,1.54)	0.55	1.10 (0.90,1.33)	0.35
Continuous covariates ^d										
Age	1.32 (1.19,1.46)	<0.001	1.25 (1.10,1.41)	<0.001	1.26 (1.06,1.51)	0.01	1.24 (1.05,1.46)	0.01	1.30 (1.18,1.44)	<0.001
Parental education	0.98 (0.89,1.09)	0.76	1.04 (0.92,1.17)	0.55	0.90 (0.75,1.07)	0.23	1.03 (0.87,1.21)	0.73	0.99 (0.90,1.09)	0.85
Peersmoking	1.09 (1.00,1.19)	0.04	1.09 (0.98,1.20)	0.11	1.03 (0.91,1.17)	0.67	1.07 (0.94,1.22)	0.29	1.09 (1.00,1.18)	0.06
CESD-Depressive Symptoms	1.07 (0.97,1.17)	0.18	1.09 (0.97,1.22)	0.16	1.06 (0.90,1.25)	0.48	1.01 (0.88,1.16)	0.88	1.07 (0.98,1.17)	0.14
TCI-Impulsivity	0.84 (0.77,0.92)	<0.001	0.79 (0.71,0.88)	<0.001	0.82 (0.70,0.96)	0.01	0.93 (0.80,1.07)	0.32	0.83 (0.76,0.91)	<0.001
Delinquent Behaviour	1.05 (0.95,1.16)	0.35	1.09 (0.97,1.24)	0.15	1.00 (0.86,1.17)	0.98	1.08 (0.93,1.25)	0.32	1.06 (0.96,1.18)	0.24
Smoking susceptibility	1.12 (1.02,1.23)	0.02	1.17 (1.04,1.30)	0.01	0.98 (0.82,1.17)	0.80	0.95 (0.82,1.10)	0.46	1.10 (1.01,1.20)	0.04
Smoking expectancies	0.94 (0.86,1.03)	0.16	0.92 (0.82,1.02)	0.12	1.01 (0.85,1.20)	0.90	1.01 (0.86,1.18)	0.91	0.95 (0.87,1.04)	0.26
Regressors										
E-cigarettes ever (vs. never) use:										
Smoking outcome on 6 months	1.37 (1.01,1.84)	0.042	1.49 (1.05,2.12)	0.028	0.98 (0.59,1.61)	0.93	0.47 (0.25, 0.89)	0.02	1.36 (1.01,1.83)	0.045
Smoking outcome on 12 months	1.37 (1.01,1.84)	0.042	1.49 (1.05,2.12)	0.028	0.98 (0.59,1.61)	0.93	0.91 (0.50, 1.68)	0.77	1.36 (1.01,1.83)	0.045
Time (12- vs. 6-month): E-cigarette never	2.78 (2.34,3.31)	<0.001	3.16 (2.56,3.92)	<0.001	2.29 (1.67,3.15)	<0.001	2.35 (1.73,3.19)	<0.001	2.80 (2.35,3.32)	<0.001
Time (12- vs. 6-month): E-cigarette ever	2.78 (2.34,3.31)	<0.001	3.16 (2.56,3.92)	<0.001	2.29 (1.67,3.15)	<0.001	1.00 (0.48,2.10)	1.00	2.80 (2.35,3.32)	<0.001
Ever e-cigarette use × Time ^c	0.95 (0.57,1.58)	0.85	0.99 (0.56,1.74)	0.96	0.71 (0.26,1.93)	0.50	0.43 (0.19,0.93)	0.033	0.84 (0.51,1.37)	0.48

Note: Sensitivity analysis: for missing outcomes values were imputed assuming the prevalence for each of these outcomes at 6 and 12-month follow-up as found in schools with the highest prevalence on each of these outcomes. All analyses include only never-users of combustible tobacco products at baseline ($N = 2185$).


^a OR from repeated binary logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects.

^b OR from repeated ordinal logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects, with the OR expressing the change in odds of being in a category with use of a certain amount of tobacco products versus being in a category with lower use (3 versus ≤ 2 , ≥ 2 versus ≤ 1 , ≥ 1 versus 0).

^c If interaction term is significant (p -value ≤ 0.05) or marginally significant (p -value ≤ 0.10), the effect of e-cigarette use is examined both at 6- and 12 months follow up, and the effect of time is examined both for never and ever users of e-cigarettes. If interaction term is not (marginally) significant, the effect of e-cigarette use is examined averaged across and, is thus the same for both the 6- and 12 months followups, and the effect of time is examined averaged across and, is thus also the same for never and ever users of e-cigarettes.

^d Continuous covariates rescaled ($M=0$, $SD=1$), such that the ORs indicate change in odds in the outcome associated with an increase in one standard deviation unit on the covariate continuous scale. CESD = Center for Epidemiologic Studies Depression Scale. TCI = Temperament and Character Inventory.

Supplementary tables and figures:

Supplementary Figure 1: Flow of Adolescent Students in Study to Assess E-Cigarette Use at Baseline and Later Use of Combustible Tobacco Products

^a Includes all combustible tobacco products.

Supplementary Table 1: Association of Baseline e-cigarette use frequency with tobacco smoking frequency at 6- and 12-month Follow-up^a

	Parameter Estimate for Association with Smoking Frequency at follow-up	
	Odds ratio (95% CI)	P-value
Unadjusted model^b		
Baseline smoking (3-level continuum):		
- Outcome at 6 months	2.06 (1.41, 3.01)	< 0.001
- Outcome at 12 months	7.11 (4.03, 12.56)	< 0.001
E-cigarette use (4-level continuum)	2.11 (1.69, 2.64)	< 0.001
E-cigarette use x smoking ^c	0.70 (0.52, 0.94)	0.02
Adjusted model^d		
Baseline smoking (3-level continuum)		
- Outcome at 6 months	1.11 (0.71, 1.73)	0.66
- Outcome at 12 months	3.52 (2.02, 6.11)	<0.001
E-cigarette use (4-level continuum)	1.63 (1.29, 2.06)	< 0.001
E-cigarette use x smoking ^c	0.83 (0.61, 1.11)	0.21

^a Ordinal logistic mixed regression of proportional odds for being at a higher smoking frequency outcome (i.e., days smoked in the past 30 days; non-smoker, 0; infrequent smokers [1-2days], 1; frequent smokers [≥3 days], 2) accounting for clustering by pupil and school in a sample with available baseline e-cigarette use and smoking frequency data (N = 2348). The baseline 4-level continuous e-cigarette use variable was categorized as never, prior (ever with no past 30-day use), infrequent (1-2 days during past 30 days), or frequent (≥3 days during past 30 days). To address missing covariate data and missing outcomes, 80 multiple imputed datasets were generated using the fully conditional specification method (with sequential regression procedure). The parameter estimates from models in each imputed data set were pooled and presented as a single estimate.

^b Unadjusted models without e-cigarette use x smoking interaction include baseline e-cigarette use and smoking frequency, and interactions of these variables with time (6 versus 12 months follow-up). In case of a significant interaction with time, parameter estimates are shown separately for 6- and 12-months follow-up, otherwise the interaction term is removed from the model.

^c Interaction term added in subsequent model. If e-cigarette use x smoking x time is not significant, the results for e-cigarette use x smoking averaged across both follow-ups are presented.

^d Adjusted models without e-cigarette use x smoking interaction include baseline e-cigarette use and smoking frequency, and interactions of these variables with time (6 versus 12 months follow-up). In case of a significant interaction with time, parameter estimates are shown separately for 6- and 12-months follow-up, otherwise the interaction term is removed from the model. Adjustment is made for demographic, environmental and psychosocial covariates as described in the main text.

Supplementary Table 2: Association of Baseline Ever Use of Any Tobacco Product and Covariates to E-cigarette Use at 6- and 12-Month Follow Ups among Baseline Never E-Cigarette Users

Baseline regressors and covariates	Ever Use of E-Cigarettes	
	OR(95%CI) ^a	P
Unadjusted Models		
Any Tobacco Product ever (vs. never) use:		
E-Cigarette use at 6 months	5.22 (3.06,8.92)	< 0.001
E-Cigarette use at 12 months	5.22 (3.06,8.92)	< 0.001
Time (12- vs. 6-month): Any Tobacco use never	1.46 (1.01,2.12)	0.04
Time (12- vs. 6-month): Any Tobacco use ever	1.46 (1.01,2.12)	0.04
Ever Tobacco Product Use × Time ^b	1.92 (0.79,4.65)	0.15
Adjusted Models		
Categorical covariates		
Girls (vs. Boys)	0.83 (0.56,1.22)	0.33
Dutch (vs. other) ethnicity	0.82 (0.33,2.06)	0.67
Lives with both parents (vs. other)	0.90 (0.59,1.39)	0.64
Substance ever (vs. never) use	2.03 (1.30,3.16)	0.002
Family history of smoking (yes vs. no)	0.97 (0.65,1.45)	0.88
Continuous covariates ^c		
Age	1.02 (0.83,1.25)	0.84
Parental education	0.99 (0.83,1.18)	0.95
Peer smoking	0.98 (0.83,1.17)	0.86
CESD-Depressive Symptoms	0.92 (0.76,1.12)	0.47
TCI-Impulsivity	0.97 (0.81,1.16)	0.71
Delinquent Behaviour	1.11 (0.96,1.29)	0.17
Smoking susceptibility	1.04 (0.85,1.26)	0.71
Smoking expectancies	0.98 (0.79,1.20)	0.82
Regressors		
Any Tobacco Product ever (vs. never) use:		
E-Cigarette use at 6 months	3.10 (1.58,6.06)	0.001
E-Cigarette use at 12 months	3.10 (1.58,6.06)	0.001
Time (12- vs. 6-month): Any Tobacco use never	1.47 (1.01,2.12)	0.04
Time (12- vs. 6-month): Any Tobacco use ever	1.47 (1.01,2.12)	0.04
Ever Tobacco Product Use × Time ^b	1.91 (0.79,4.63)	0.15

Note: All analyses include only never-users of E-Cigarettes at baseline (N = 2191).

^a OR from repeated binary logistic regression model predicting E-cigarette use from baseline ever tobacco use status (yes/no) including school fixed effects.

^b If interaction term is significant (p-value ≤ 0.05), the effect of any tobacco product use is examined both at 6- and 12 months follow up, and the effect of time is examined both for never and ever users of any tobacco product. If interaction term is not significant, the effect of any tobacco use is examined averaged across and, is thus the same for both the 6- and 12 months follow ups, and the effect of time is examined averaged across and, is thus also the same for never and ever users of tobacco products.

^c Continuous covariates rescaled (M = 0, SD = 1), such that the ORs indicate change in odds in the outcome associated with an increase in one standard deviation unit on the covariate continuous scale. CESD = Center for Epidemiologic Studies Depression Scale. TCI = Temperament and Character Inventory.

Supplementary Table 3: Sensitivity analyses of the Association of Baseline E-Cigarette Ever Use and Covariates to Combustible Tobacco Use Outcomes at 6- and 12-Month Follow Ups among Baseline Never-Smokers

Baseline regressors and covariates	Outcome									
	Any tobacco product		Combustible cigarettes		Cigars		Hookah		Number of Tobacco Products	
	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^b	P
Unadjusted Models										
E-cigarette ever (vs. never) use:										
Smoking outcome on 6 months	2.33 (1.75,3.11)	<0.001	2.62 (1.85,3.71)	<0.001	1.43 (0.90,2.25)	0.13	3.00 (1.61,5.60)	0.001	2.30 (1.73, 3.08)	<0.001
Smoking outcome on 12 months	2.33 (1.75,3.11)	<0.001	2.62 (1.85,3.71)	<0.001	1.43 (0.90,2.25)	0.13	1.28 (0.71,2.32)	0.41	2.30 (1.73,3.08)	<0.001
Time (12- vs. 6-month): E-Cigarette never	2.77 (2.33,3.30)	<0.001	3.16 (2.55,3.91)	<0.001	2.32 (1.69,3.19)	<0.001	2.34 (1.73,3.18)	<0.001	2.80 (2.35,3.33)	<0.001
Time (12- vs. 6-month): E-Cigarette ever	2.77 (2.33,3.30)	<0.001	3.16 (2.55,3.91)	<0.001	2.32 (1.69,3.19)	<0.001	1.00 (0.49,2.06)	1.00	2.80 (2.35,3.33)	<0.001
Ever e-cigarette use × Time ^c	0.95 (0.57,1.57)	0.85	0.98 (0.56,1.73)	0.96	0.71 (0.26,1.92)	0.50	0.43 (0.20,0.93)	0.033	0.85 (0.52,1.39)	0.51
Adjusted Models										
Categorical covariates										
Girls (vs. Boys)	0.97 (0.81,1.16)	0.75	0.91 (0.73,1.15)	0.44	0.91 (0.66,1.25)	0.56	1.02 (0.76,1.38)	0.89	0.95 (0.80,1.14)	0.60
Dutch (vs. other) ethnicity	0.59 (0.38,0.91)	0.02	0.68 (0.41,1.12)	0.13	1.07 (0.48,2.42)	0.87	0.33 (0.17,0.65)	0.001	0.57 (0.36,0.90)	0.02
Lives with both parents (vs. other)	1.10 (0.88,1.37)	0.40	1.27 (0.96,1.67)	0.10	0.93 (0.65,1.34)	0.71	1.11 (0.76,1.61)	0.59	1.12 (0.90,1.40)	0.31
Substance ever (vs. never) use	1.28 (1.04,1.56)	0.02	1.21 (0.95,1.56)	0.13	1.47 (1.05,2.05)	0.03	1.39 (0.99,1.94)	0.06	1.30 (1.07,1.59)	0.01
Family history of smoking (yes vs. no)	1.12 (0.92,1.37)	0.24	1.05 (0.83,1.34)	0.68	0.98 (0.70,1.38)	0.93	1.11 (0.80,1.54)	0.55	1.10 (0.90,1.33)	0.35
Continuous covariates ^d										
Age	1.32 (1.19,1.46)	<0.001	1.25 (1.10,1.41)	<0.001	1.26 (1.06,1.51)	0.01	1.24 (1.05,1.46)	0.01	1.30 (1.18,1.44)	<0.001
Parental education	0.98 (0.89,1.09)	0.76	1.04 (0.92,1.17)	0.55	0.90 (0.75,1.07)	0.23	1.03 (0.87,1.21)	0.73	0.99 (0.90,1.09)	0.85
Peersmoking	1.09 (1.00,1.19)	0.04	1.09 (0.98,1.20)	0.11	1.03 (0.91,1.17)	0.67	1.07 (0.94,1.22)	0.29	1.09 (1.00,1.18)	0.06
CESD-Depressive Symptoms	1.07 (0.97,1.17)	0.18	1.09 (0.97,1.22)	0.16	1.06 (0.90,1.25)	0.48	1.01 (0.88,1.16)	0.88	1.07 (0.98,1.17)	0.14
TCI-Impulsivity	0.84 (0.77,0.92)	<0.001	0.79 (0.71,0.88)	<0.001	0.82 (0.70,0.96)	0.01	0.93 (0.80,1.07)	0.32	0.83 (0.76,0.91)	<0.001
Delinquent Behaviour	1.05 (0.95,1.16)	0.35	1.09 (0.97,1.24)	0.15	1.00 (0.86,1.17)	0.98	1.08 (0.93,1.25)	0.32	1.06 (0.96,1.18)	0.24
Smoking susceptibility	1.12 (1.02,1.23)	0.02	1.17 (1.04,1.30)	0.01	0.98 (0.82,1.17)	0.80	0.95 (0.82,1.10)	0.46	1.10 (1.01,1.20)	0.04
Smoking expectancies	0.94 (0.86,1.03)	0.16	0.92 (0.82,1.02)	0.12	1.01 (0.85,1.20)	0.90	1.01 (0.86,1.18)	0.91	0.95 (0.87,1.04)	0.26
Regressors										
E-cigarettes ever (vs. never) use:										
Smoking outcome on 6 months	1.37 (1.01,1.84)	0.042	1.49 (1.05,2.12)	0.028	0.98 (0.59,1.61)	0.93	0.47 (0.25, 0.89)	0.02	1.36 (1.01,1.83)	0.045
Smoking outcome on 12 months	1.37 (1.01,1.84)	0.042	1.49 (1.05,2.12)	0.028	0.98 (0.59,1.61)	0.93	0.91 (0.50, 1.68)	0.77	1.36 (1.01,1.83)	0.045
Time (12- vs. 6-month): E-cigarette never	2.78 (2.34,3.31)	<0.001	3.16 (2.56,3.92)	<0.001	2.29 (1.67,3.15)	<0.001	2.35 (1.73,3.19)	<0.001	2.80 (2.35,3.32)	<0.001
Time (12- vs. 6-month): E-cigarette ever	2.78 (2.34,3.31)	<0.001	3.16 (2.56,3.92)	<0.001	2.29 (1.67,3.15)	<0.001	1.00 (0.48,2.10)	1.00	2.80 (2.35,3.32)	<0.001
Ever e-cigarette use × Time ^c	0.95 (0.57,1.58)	0.85	0.99 (0.56,1.74)	0.96	0.71 (0.26,1.93)	0.50	0.43 (0.19,0.93)	0.033	0.84 (0.51,1.37)	0.48

Note: Sensitivity analysis: for missing outcomes values were imputed assuming the prevalence for each of these outcomes at 6 and 12-month follow-up as found in schools with the highest prevalence on each of these outcomes. All analyses include only never-users of combustible tobacco products at baseline ($N = 2185$).


^a OR from repeated binary logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects.

^b OR from repeated ordinal logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects, with the OR expressing the change in odds of being in a category with use of a certain amount of tobacco products versus being in a category with lower use (3 versus ≤ 2 , ≥ 2 versus ≤ 1 , ≥ 1 versus 0).

^c If interaction term is significant (p-value ≤ 0.05) or marginally significant (p-value ≤ 0.10), the effect of e-cigarette use is examined both at 6- and 12 months follow up, and the effect of time is examined both for never and ever users of e-cigarettes. If interaction term is not (marginally) significant, the effect of e-cigarette use is examined averaged across and, is thus the same for both the 6- and 12 months followups, and the effect of time is examined averaged across and, is thus also the same for never and ever users of e-cigarettes.

^d Continuous covariates rescaled ($M=0$, $SD=1$), such that the ORs indicate change in odds in the outcome associated with an increase in one standard deviation unit on the covariate continuous scale. CESD = Center for Epidemiologic Studies Depression Scale. TCI = Temperament and Character Inventory.

Supplementary tables and figures:

Supplementary Figure 1: Flow of Adolescent Students in Study to Assess E-Cigarette Use at Baseline and Later Use of Combustible Tobacco Products

^a Includes all combustible tobacco products.

Supplementary Table 1: Association of Baseline e-cigarette use frequency with tobacco smoking frequency at 6- and 12-month Follow-up^a

	Parameter Estimate for Association with Smoking Frequency at follow-up	
	Odds ratio (95% CI)	P-value
Unadjusted model^b		
Baseline smoking (3-level continuum):		
- Outcome at 6 months	2.06 (1.41, 3.01)	< 0.001
- Outcome at 12 months	7.11 (4.03, 12.56)	< 0.001
E-cigarette use (4-level continuum)	2.11 (1.69, 2.64)	< 0.001
E-cigarette use x smoking ^c	0.70 (0.52, 0.94)	0.02
Adjusted model^d		
Baseline smoking (3-level continuum)		
- Outcome at 6 months	1.11 (0.71, 1.73)	0.66
- Outcome at 12 months	3.52 (2.02, 6.11)	<0.001
E-cigarette use (4-level continuum)	1.63 (1.29, 2.06)	< 0.001
E-cigarette use x smoking ^c	0.83 (0.61, 1.11)	0.21

^a Ordinal logistic mixed regression of proportional odds for being at a higher smoking frequency outcome (i.e., days smoked in the past 30 days; non-smoker, 0; infrequent smokers [1-2days], 1; frequent smokers [≥3 days], 2) accounting for clustering by pupil and school in a sample with available baseline e-cigarette use and smoking frequency data (N = 2348). The baseline 4-level continuous e-cigarette use variable was categorized as never, prior (ever with no past 30-day use), infrequent (1-2 days during past 30 days), or frequent (≥3 days during past 30 days). To address missing covariate data and missing outcomes, 80 multiple imputed datasets were generated using the fully conditional specification method (with sequential regression procedure). The parameter estimates from models in each imputed data set were pooled and presented as a single estimate.

^b Unadjusted models without e-cigarette use x smoking interaction include baseline e-cigarette use and smoking frequency, and interactions of these variables with time (6 versus 12 months follow-up). In case of a significant interaction with time, parameter estimates are shown separately for 6- and 12-months follow-up, otherwise the interaction term is removed from the model.

^c Interaction term added in subsequent model. If e-cigarette use x smoking x time is not significant, the results for e-cigarette use x smoking averaged across both follow-ups are presented.

^d Adjusted models without e-cigarette use x smoking interaction include baseline e-cigarette use and smoking frequency, and interactions of these variables with time (6 versus 12 months follow-up). In case of a significant interaction with time, parameter estimates are shown separately for 6- and 12-months follow-up, otherwise the interaction term is removed from the model. Adjustment is made for demographic, environmental and psychosocial covariates as described in the main text.

Supplementary Table 2: Association of Baseline Ever Use of Any Tobacco Product and Covariates to E-cigarette Use at 6- and 12-Month Follow Ups among Baseline Never E-Cigarette Users

Baseline regressors and covariates	Ever Use of E-Cigarettes	
	OR(95%CI) ^a	P
Unadjusted Models		
Any Tobacco Product ever (vs. never) use:		
E-Cigarette use at 6 months	5.22 (3.06,8.92)	< 0.001
E-Cigarette use at 12 months	5.22 (3.06,8.92)	< 0.001
Time (12- vs. 6-month): Any Tobacco use never	1.46 (1.01,2.12)	0.04
Time (12- vs. 6-month): Any Tobacco use ever	1.46 (1.01,2.12)	0.04
Ever Tobacco Product Use × Time ^b	1.92 (0.79,4.65)	0.15
Adjusted Models		
Categorical covariates		
Girls (vs. Boys)	0.83 (0.56,1.22)	0.33
Dutch (vs. other) ethnicity	0.82 (0.33,2.06)	0.67
Lives with both parents (vs. other)	0.90 (0.59,1.39)	0.64
Substance ever (vs. never) use	2.03 (1.30,3.16)	0.002
Family history of smoking (yes vs. no)	0.97 (0.65,1.45)	0.88
Continuous covariates ^c		
Age	1.02 (0.83,1.25)	0.84
Parental education	0.99 (0.83,1.18)	0.95
Peer smoking	0.98 (0.83,1.17)	0.86
CESD-Depressive Symptoms	0.92 (0.76,1.12)	0.47
TCI-Impulsivity	0.97 (0.81,1.16)	0.71
Delinquent Behaviour	1.11 (0.96,1.29)	0.17
Smoking susceptibility	1.04 (0.85,1.26)	0.71
Smoking expectancies	0.98 (0.79,1.20)	0.82
Regressors		
Any Tobacco Product ever (vs. never) use:		
E-Cigarette use at 6 months	3.10 (1.58,6.06)	0.001
E-Cigarette use at 12 months	3.10 (1.58,6.06)	0.001
Time (12- vs. 6-month): Any Tobacco use never	1.47 (1.01,2.12)	0.04
Time (12- vs. 6-month): Any Tobacco use ever	1.47 (1.01,2.12)	0.04
Ever Tobacco Product Use × Time ^b	1.91 (0.79,4.63)	0.15

Note: All analyses include only never-users of E-Cigarettes at baseline (N = 2191).

^a OR from repeated binary logistic regression model predicting E-cigarette use from baseline ever tobacco use status (yes/no) including school fixed effects.

^b If interaction term is significant (p-value ≤ 0.05), the effect of any tobacco product use is examined both at 6- and 12 months follow up, and the effect of time is examined both for never and ever users of any tobacco product. If interaction term is not significant, the effect of any tobacco use is examined averaged across and, is thus the same for both the 6- and 12 months follow ups, and the effect of time is examined averaged across and, is thus also the same for never and ever users of tobacco products.

^c Continuous covariates rescaled (M = 0, SD = 1), such that the ORs indicate change in odds in the outcome associated with an increase in one standard deviation unit on the covariate continuous scale. CESD = Center for Epidemiologic Studies Depression Scale. TCI = Temperament and Character Inventory.

Supplementary Table 3: Sensitivity analyses of the Association of Baseline E-Cigarette Ever Use and Covariates to Combustible Tobacco Use Outcomes at 6- and 12-Month Follow Ups among Baseline Never-Smokers

Baseline regressors and covariates	Outcome									
	Any tobacco product		Combustible cigarettes		Cigars		Hookah		Number of Tobacco Products	
	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^a	P	OR(95%CI) ^b	P
Unadjusted Models										
E-cigarette ever (vs. never) use:										
Smoking outcome on 6 months	2.33 (1.75,3.11)	<0.001	2.62 (1.85,3.71)	<0.001	1.43 (0.90,2.25)	0.13	3.00 (1.61,5.60)	0.001	2.30 (1.73, 3.08)	<0.001
Smoking outcome on 12 months	2.33 (1.75,3.11)	<0.001	2.62 (1.85,3.71)	<0.001	1.43 (0.90,2.25)	0.13	1.28 (0.71,2.32)	0.41	2.30 (1.73,3.08)	<0.001
Time (12- vs. 6-month): E-Cigarette never	2.77 (2.33,3.30)	<0.001	3.16 (2.55,3.91)	<0.001	2.32 (1.69,3.19)	<0.001	2.34 (1.73,3.18)	<0.001	2.80 (2.35,3.33)	<0.001
Time (12- vs. 6-month): E-Cigarette ever	2.77 (2.33,3.30)	<0.001	3.16 (2.55,3.91)	<0.001	2.32 (1.69,3.19)	<0.001	1.00 (0.49,2.06)	1.00	2.80 (2.35,3.33)	<0.001
Ever e-cigarette use × Time ^c	0.95 (0.57,1.57)	0.85	0.98 (0.56,1.73)	0.96	0.71 (0.26,1.92)	0.50	0.43 (0.20,0.93)	0.033	0.85 (0.52,1.39)	0.51
Adjusted Models										
Categorical covariates										
Girls (vs. Boys)	0.97 (0.81,1.16)	0.75	0.91 (0.73,1.15)	0.44	0.91 (0.66,1.25)	0.56	1.02 (0.76,1.38)	0.89	0.95 (0.80,1.14)	0.60
Dutch (vs. other) ethnicity	0.59 (0.38,0.91)	0.02	0.68 (0.41,1.12)	0.13	1.07 (0.48,2.42)	0.87	0.33 (0.17,0.65)	0.001	0.57 (0.36,0.90)	0.02
Lives with both parents (vs. other)	1.10 (0.88,1.37)	0.40	1.27 (0.96,1.67)	0.10	0.93 (0.65,1.34)	0.71	1.11 (0.76,1.61)	0.59	1.12 (0.90,1.40)	0.31
Substance ever (vs. never) use	1.28 (1.04,1.56)	0.02	1.21 (0.95,1.56)	0.13	1.47 (1.05,2.05)	0.03	1.39 (0.99,1.94)	0.06	1.30 (1.07,1.59)	0.01
Family history of smoking (yes vs. no)	1.12 (0.92,1.37)	0.24	1.05 (0.83,1.34)	0.68	0.98 (0.70,1.38)	0.93	1.11 (0.80,1.54)	0.55	1.10 (0.90,1.33)	0.35
Continuous covariates ^d										
Age	1.32 (1.19,1.46)	<0.001	1.25 (1.10,1.41)	<0.001	1.26 (1.06,1.51)	0.01	1.24 (1.05,1.46)	0.01	1.30 (1.18,1.44)	<0.001
Parental education	0.98 (0.89,1.09)	0.76	1.04 (0.92,1.17)	0.55	0.90 (0.75,1.07)	0.23	1.03 (0.87,1.21)	0.73	0.99 (0.90,1.09)	0.85
Peersmoking	1.09 (1.00,1.19)	0.04	1.09 (0.98,1.20)	0.11	1.03 (0.91,1.17)	0.67	1.07 (0.94,1.22)	0.29	1.09 (1.00,1.18)	0.06
CESD-Depressive Symptoms	1.07 (0.97,1.17)	0.18	1.09 (0.97,1.22)	0.16	1.06 (0.90,1.25)	0.48	1.01 (0.88,1.16)	0.88	1.07 (0.98,1.17)	0.14
TCI-Impulsivity	0.84 (0.77,0.92)	<0.001	0.79 (0.71,0.88)	<0.001	0.82 (0.70,0.96)	0.01	0.93 (0.80,1.07)	0.32	0.83 (0.76,0.91)	<0.001
Delinquent Behaviour	1.05 (0.95,1.16)	0.35	1.09 (0.97,1.24)	0.15	1.00 (0.86,1.17)	0.98	1.08 (0.93,1.25)	0.32	1.06 (0.96,1.18)	0.24
Smoking susceptibility	1.12 (1.02,1.23)	0.02	1.17 (1.04,1.30)	0.01	0.98 (0.82,1.17)	0.80	0.95 (0.82,1.10)	0.46	1.10 (1.01,1.20)	0.04
Smoking expectancies	0.94 (0.86,1.03)	0.16	0.92 (0.82,1.02)	0.12	1.01 (0.85,1.20)	0.90	1.01 (0.86,1.18)	0.91	0.95 (0.87,1.04)	0.26
Regressors										
E-cigarettes ever (vs. never) use:										
Smoking outcome on 6 months	1.37 (1.01,1.84)	0.042	1.49 (1.05,2.12)	0.028	0.98 (0.59,1.61)	0.93	0.47 (0.25, 0.89)	0.02	1.36 (1.01,1.83)	0.045
Smoking outcome on 12 months	1.37 (1.01,1.84)	0.042	1.49 (1.05,2.12)	0.028	0.98 (0.59,1.61)	0.93	0.91 (0.50, 1.68)	0.77	1.36 (1.01,1.83)	0.045
Time (12- vs. 6-month): E-cigarette never	2.78 (2.34,3.31)	<0.001	3.16 (2.56,3.92)	<0.001	2.29 (1.67,3.15)	<0.001	2.35 (1.73,3.19)	<0.001	2.80 (2.35,3.32)	<0.001
Time (12- vs. 6-month): E-cigarette ever	2.78 (2.34,3.31)	<0.001	3.16 (2.56,3.92)	<0.001	2.29 (1.67,3.15)	<0.001	1.00 (0.48,2.10)	1.00	2.80 (2.35,3.32)	<0.001
Ever e-cigarette use × Time ^c	0.95 (0.57,1.58)	0.85	0.99 (0.56,1.74)	0.96	0.71 (0.26,1.93)	0.50	0.43 (0.19,0.93)	0.033	0.84 (0.51,1.37)	0.48

Note: Sensitivity analysis: for missing outcomes values were imputed assuming the prevalence for each of these outcomes at 6 and 12-month follow-up as found in schools with the highest prevalence on each of these outcomes. All analyses include only never-users of combustible tobacco products at baseline ($N = 2185$).

^a OR from repeated binary logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects.

^b OR from repeated ordinal logistic regression model predicting respective outcome from baseline ever e-cigarette use status (yes/no) including school fixed effects, with the OR expressing the change in odds of being in a category with use of a certain amount of tobacco products versus being in a category with lower use (3 versus ≤ 2 , ≥ 2 versus ≤ 1 , ≥ 1 versus 0).

^c If interaction term is significant (p-value ≤ 0.05) or marginally significant (p-value ≤ 0.10), the effect of e-cigarette use is examined both at 6- and 12 months follow up, and the effect of time is examined both for never and ever users of e-cigarettes. If interaction term is not (marginally) significant, the effect of e-cigarette use is examined averaged across and, is thus the same for both the 6- and 12 months followups, and the effect of time is examined averaged across and, is thus also the same for never and ever users of e-cigarettes.

^d Continuous covariates rescaled ($M=0$, $SD=1$), such that the ORs indicate change in odds in the outcome associated with an increase in one standard deviation unit on the covariate continuous scale. CESD = Center for Epidemiologic Studies Depression Scale. TCI = Temperament and Character Inventory.