



OPEN ACCESS

Impact of Canada's menthol cigarette ban on quitting among menthol smokers: pooled analysis of pre-post evaluation from the ITC Project and the Ontario Menthol Ban Study and projections of impact in the USA

Geoffrey T Fong ^{1,2,3}, Janet Chung-Hall ¹, Gang Meng,¹ Lorraine V Craig ¹, Mary E Thompson,⁴ Anne C K Quah,¹ K Michael Cummings ⁵, Andrew Hyland,⁶ Richard J O'Connor ⁶, David T Levy ⁷, Cristine D Delnevo,⁸ Ollie Ganz,⁸ Thomas Eissenberg ⁹, Eric K Soule ¹⁰, Robert Schwartz ¹¹, Joanna E Cohen ¹², Michael O Chaiton ¹³

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/tobaccocontrol-2021-057227>).

For numbered affiliations see end of article.

Correspondence to

Professor Geoffrey T Fong, Department of Psychology and School of Public Health Sciences, University of Waterloo, Waterloo, ON N2L 3G1, Canada; gfong@uwaterloo.ca

Received 15 December 2021
Accepted 14 April 2022
Published Online First
28 April 2022



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Fong GT, Chung-Hall J, Meng G, et al. *Tob Control* 2023;**32**:734–738.

ABSTRACT

Introduction Between 2015 and 2018, Canada banned menthol cigarettes. This study pooled data from two pre-post cohort studies (the Ontario Menthol Ban Study, and the International Tobacco Control Policy Evaluation (ITC) Canada Survey, conducted in seven provinces) to derive more precise estimates of the impact of Canada's menthol ban on quitting and to apply these estimates to project the impact of a menthol ban in the USA.

Methods Weighted multivariable logistic analyses compared post-ban quit success of menthol smokers with non-menthol smokers (for daily smokers and for all (daily + non-daily) smokers), controlling for sex, age, ethnicity, education, baseline smoking status, baseline cigarettes per day and study regions. Projections to the USA were created by multiplying the effect size of the Canadian menthol ban on quitting (percentage of increased quitting among menthol smokers) by the number of menthol smokers overall and among African Americans, from the 2019 National Survey on Drug Use and Health.

Results After the menthol cigarette ban, menthol smokers were more likely than non-menthol smokers to have quit smoking among daily smokers (difference=8.0%; 95% CI: 2.4% to 13.7%, p=0.005) and all (daily+non-daily) smokers (difference=7.3%; 95% CI: 2.1% to 12.5%, p=0.006). The projected number of smokers who would quit after a US menthol ban would be 789 724 daily smokers (including 199 732 African Americans) and 1 337 988 daily+non-daily smokers (including 381 272 African Americans).

Conclusions This pooled analysis of Canada's menthol cigarette ban provides the foundation for estimating the impact of menthol bans in the USA and other countries. Projections suggest that a US menthol cigarette ban would have a substantial impact on increasing quitting.

INTRODUCTION

Between 2015 and 2018, all Canadian provinces banned menthol in cigarettes. Two pre-post studies found that the Canadian menthol ban led

to increases in quitting among menthol smokers compared with non-menthol smokers. This study pools data from those two cohort studies—the International Tobacco Control Policy Evaluation (ITC) Project Canada Survey¹ and the Ontario Menthol Ban Study²—to derive a more precise estimate of the impact of Canada's menthol ban on quitting.

The US Food and Drug Administration (FDA) announced on 29 April 2021 that they will work toward developing product standards 'within the next year' to ban menthol as a characterising flavour in cigarettes and cigars.³ The FDA further stated that, 'This decision is based on clear science and evidence establishing the addictiveness and harm of these products and builds on important, previous actions that banned other flavored cigarettes in 2009.' More specifically, that science indicates that menthol in cigarettes increases initiation, facilitates progression to regular smoking, increases dependence and decreases the likelihood of smoking cessation, especially among African American smokers. The large declines in cigarette consumption in the USA over the last two decades have been in non-menthol cigarettes.⁴ During that same period, menthol's market share has continued to increase.⁵

We applied effect sizes from this pooled analysis of the impact of the Canadian menthol cigarette ban to project the impact of a proposed menthol ban in the USA on cessation of cigarette smoking among adults—overall and among African Americans.

METHODS

We analysed pooled pre-post data from (a) the ITC Canada Survey (the Canadian arm of the ITC Four Country Smoking and Vaping Survey) across seven provinces covering 83% of the Canadian population (Quebec, Ontario, Prince Edward Island, Newfoundland and Labrador, British Columbia, Saskatchewan, and Manitoba; n=1236 adult (18+ years) smokers, 128 pre-ban mainly menthol smokers) and (b) the Ontario Menthol Ban Study (n=1084 adult (18+ years) smokers, 295 pre-ban

Table 1 Quit success among daily and all (daily+non-daily) menthol and non-menthol adult (18+) smokers after the Canadian menthol cigarette ban, with effect sizes applied to menthol smokers (all smokers and African American smokers), based on the number of US adult (18+) menthol smokers from the 2019 National Survey on Drug Use and Health (NSDUH)

	Percentage of quitters among adult smokers from the pooled analysis (7 provinces from ITC Survey+Ontario from Ontario Study)			Effect size of Canadian menthol ban applied to all US adult menthol smokers		Effect size of Canadian menthol ban applied to African American adult menthol smokers	
	Mainly menthol	Non-menthol	Difference: menthol ban effect size (95% CI)	All smokers (95% CI)	Daily smokers only (95% CI)	All smokers (95% CI)	Daily smokers only (95% CI)
NSDUH 2019: number of US adult menthol smokers	N/A	N/A	N/A	18328597	9871550	5222907	2496650
Quit success: daily smokers	21.2%	13.2%	8.0% (2.4% to 13.7%)*	N/A	789724 (236917 to 1352402)	N/A	199732 (59920 to 342041)
Quit success: all smokers (daily+non-daily)	22.3%	15.0%	7.3% (2.1% to 12.5%)**	1337988 (384901 to 2291075)	N/A	381272 (109681 to 652863)	N/A

*P=0.005; **p=0.006.
 Mainly menthol smokers: respondents who reported smoking menthol cigarettes 'frequently' (Ontario Study) or reported a menthol brand as their usual brand (ITC Survey). Definition of 'frequent' use of menthol is provided in online supplemental table 1.
 Quit success: wave 1 smokers who reported at wave 2 that they were not smoking at all.
 Implementation of menthol cigarette bans in the seven provinces included in analysis of the ITC Canada Survey data: Quebec (26 August 2016), Ontario (1 January 2017), Prince Edward Island (1 May 2017), Newfoundland and Labrador (1 July 2017), British Columbia (2 October 2017), Saskatchewan (2 October 2017), and Manitoba (2 October 2017).
 ITC, International Tobacco Control Policy Evaluation Project; NA, not applicable.

mainly menthol smokers). Both studies used similar definitions of menthol smoking status and post-ban quitting, and were conducted within 1–2 months of each other at pre-ban (2016) and post-ban (2018) waves. The sampling strategies for each of the studies do not differ in ways that would affect our estimates. Additional details on the two surveys are provided in online supplemental table 1.

Weighted multivariable logistic analyses compared post-ban quit success (those reporting no longer smoking at post-ban) of menthol smokers with non-menthol smokers (for daily smokers and for all smokers), controlling for common covariates: sex, age, ethnicity, education, baseline smoking status, baseline cigarettes per day and study regions (ITC Survey: seven provinces; Ontario Study: Ontario as a separate region). ITC weights were rescaled to sum to the sample size to be comparable with the Ontario data set, while weights in the Ontario data set were assigned a value of 1 for each respondent. Respondents in the Ontario data set were assigned to a single stratum, which was combined with the ITC strata to account for potential design effects.

Data from the 2019 National Survey on Drug Use and Health (NSDUH) were used to obtain the total number of daily and non-daily menthol cigarette smokers in the USA, and the number of daily and non-daily African American menthol smokers. The NSDUH data for those 18 years and older were used to be consistent with the two Canadian studies. These numbers are presented in the first row of table 1. The effect size from the pooled analysis of the Canadian menthol ban—the increased percentage of menthol smokers who quit versus non-menthol smokers—was then multiplied by the number of menthol smokers (overall and among African Americans) to obtain the projected number of additional smokers who would quit in the USA, assuming that the impact of a future US menthol ban would be that found in the Canadian menthol ban. These projections were made for all smokers and for daily smokers, using the corresponding effect sizes from the pooled analysis.

RESULTS

Banning menthol cigarettes in Canada was associated with greater percentages in quitting at the post-ban survey among menthol smokers versus non-menthol smokers. Table 1 presents the quit rates for mainly menthol smokers versus non-menthol smokers from the pooled analysis for daily smokers (row 2)

and for all smokers (daily and non-daily) (row 3). The effect size of the menthol ban was 8.0 percentage points among daily smokers (95% CI: 2.4 to 13.7) and 7.3 percentage points among daily and non-daily smokers combined (95% CI: 2.1 to 12.5) (more detailed analyses (not shown) found that the impact of the menthol ban on increasing quitting among menthol smokers compared with non-menthol smokers did not differ across the seven provinces).

The projections of the potential impact of a US menthol ban are obtained by the product of each effect size and the number of corresponding menthol smokers from the NSDUH 2019 numbers of menthol smokers in row 1. Applying the Canadian results to the USA, where the percentage of current smokers who smoked menthol cigarettes was much higher (ie, 40% overall and 85% among African Americans in 2019) than in Canada prior to the menthol ban (ie, 9.5%),⁶ a US menthol ban is projected to lead to increases in quitting of 789724 daily smokers (95% CI: 236917 to 1352402) and 1337988 all smokers (daily+non-daily) (95% CI: 384901 to 2291075). Among African Americans, the projected increase in quitting would be 199732 daily smokers (95% CI: 59920 to 342041) and 381272 all smokers (95% CI: 109681 to 652863).

Further analysis to examine the impact of the Canadian menthol ban by ethnicity found a significant interaction effect, such that black Canadian mainly menthol smokers were more likely than non-black menthol smokers to have quit smoking at follow-up (p=0.029). Additional details are provided in online supplemental table 2. Because of the small sample size of black Canadian smokers (n=30) and black menthol smokers (n=4 mainly menthol smokers; n=9 occasional menthol smokers), we did not attempt to estimate post-ban quit rates among black Canadian menthol smokers and to project those estimates to US African American smokers. Instead, we used the overall effect size to project the impact of a menthol ban on African American smokers.

DISCUSSION

The pooled analysis combining the only two population-level cohort studies of the impact of the menthol cigarette ban in Canada strengthened the conclusion of each study that banning menthol cigarettes was associated with significant increases in quitting, including the new finding that the Canadian menthol ban showed similarly positive associations among all menthol

smokers (daily and non-daily menthol smokers combined)—an association that had not been statistically significant in either study alone.

Our estimates of the association of the Canadian menthol ban with cigarette smoking are more modest compared with estimates from earlier studies derived from asking smokers and young people how they might respond to a hypothetical menthol ban. In the 2010 Tobacco Use Supplement to the Current Population Survey, 39.0% of all menthol smokers and 46.8% of African American menthol smokers reported that they would stop smoking if menthol cigarettes were not available.⁷ A US survey of adolescent and adult smokers found that among the 36% who were menthol smokers, 35% reported that they would quit smoking if ‘menthol were removed from cigarettes’.⁸

A recent study using expert elicitation⁹ predicted that within 2 years after implementation of a US menthol cigarette and cigar ban, total combustible prevalence among menthol smokers aged 35–54 years would decline by about 20% from 75% to 55%, with about 10% switching to novel nicotine delivery products such as e-cigarettes. This study also projected that a menthol cigarette and cigar ban would significantly decrease initiation of smoking among youth and young adults. A related simulation analysis¹⁰ of a US menthol cigarette and cigar ban projected that cigarette smoking would decline by 14.7% in 5 years, reducing smoking-attributable and vaping-attributable life-years lost by 8.8% over a 40-year period.

Those self-reported hypothetical quit rates among US smokers are higher than the actual quit rates among Canadian menthol smokers after menthol cigarettes were banned (22.3%). However, it is important to note that the Canadian evaluation study measured the quit rate at a single time point, and thus is not directly comparable with the hypothetical quit rate of either US survey, which was temporally unbounded.

This study has significant strengths. One strength is the similarity in the timing of the surveys and the definitions of menthol smoker and measures of quitting. Internal validity was heightened by the presence of non-menthol smokers as the comparison group; this constitutes a significant challenge to alternative explanations, which would have to have increased quitting among menthol smokers to a significantly greater extent than among non-menthol smokers to have produced the observed effect. Although there were some differences in the sampling and modality of the two surveys, the effect size (difference in quit rate between menthol smokers and non-menthol smokers) did not differ between the Ontario Menthol Ban Survey and the Ontario respondents of the ITC Canada Survey ($p=0.56$), which increases the confidence in the pooled results, in accordance with the Bradford Hill principle that consistency of findings from studies employing different methodologies strengthens the likelihood of that effect.^{11 12}

There are several limitations. First, the menthol measure in NSDUH is based on self-report, which is subject to some minor misclassification,¹³ which would lead to slight differences in the number of menthol smokers that formed the basis for the projections.

Second, because of the small sample of black Canadian smokers in this study, we used the overall effect size to project quit rates among African American smokers in response to a menthol cigarette ban in the USA. We believe that applying our Canadian effect size from our pooled Canadian population cohort data to African American smokers might underestimate the possible impact of a menthol cigarette ban in the USA, given the findings of surveys asking menthol smokers what they would do if menthol cigarettes were banned, which show that African

American menthol smokers were more likely than white menthol smokers to say that they would quit,^{7 14} a pattern that is also suggested in the expert elicitation studies.^{9 10}

The most specific data regarding what quit rates might be for black menthol smokers relative to non-black menthol smokers in the USA following a menthol ban come from our pooled Canadian analysis. Despite our low sample size of only 30 black smokers, of which 4 were mainly menthol smokers, we did find that black menthol smokers had a significantly higher quit rate than non-black menthol smokers ($p=0.029$; see online supplemental table 2). But because of the very low sample size, we did not apply the *effect size* from this analysis. We instead used the general population effect size in our projections to African American menthol smokers. The studies on reactions to hypothetical bans, the projections of experts in elicitation studies and our own small sample analyses, all suggesting that quit rates among African American menthol smokers would be higher, lead us to conclude that our projection of the number of additional African American menthol smokers who would quit after a US menthol cigarette ban, based on the general population effect size, is likely to be conservative.

Third, there are uncertainties about the extent to which the Canadian findings can be generalised to the USA, as there are differences in the smoking and policy landscapes between the two countries. For example, the effect of a menthol cigarette ban could be affected by country differences in the illicit cigarette market. In Canada, interprovincial cigarette smuggling is negligible, possibly because the primary source of illicit cigarettes are First Nations reserves (most located in provinces of Ontario and Quebec) that have large distribution capacities^{15 16}; however, there was a relatively small market for menthol cigarettes prior to bans. In contrast, a large portion of the US illicit tobacco market involves cross-border sales of bootlegged tobacco from states with low taxes that is trafficked to localities with substantially higher prices.^{17 18}

Not surprisingly, tobacco companies have argued that a US menthol ban could lead to increased smuggling activity.^{19 20} Others, however, have argued that a menthol cigarette ban would cut off supply of bootlegged cigarette brands and lead to an overall decrease in the illicit cigarette market²¹—which is consistent with finding of no surge in illicit menthol cigarettes after the Canadian province of Nova Scotia implemented a menthol cigarette ban.²²

Finally, while both countries implemented a ban on non-menthol cigarettes prior to a similar ban on menthol cigarettes (Canada: July 2010,²³ before menthol cigarette bans implemented between May 2015 and October 2017^{24 25}; USA: September 2009,²⁶ before proposed menthol cigarette ban³), other important differences in the policy environments should be considered. For example, it is possible that strong restrictions on most forms of tobacco advertising and promotion in Canada led to different post-ban behaviours than would be observed in the USA, where many forms of direct tobacco advertising (eg, print media, billboards/outdoor and point of sale) are still permitted, and exposure to menthol cigarette advertising specifically is higher among African American populations than other racial or ethnic groups.²⁷

In summary, the findings from the evaluation of the actual menthol cigarette ban in Canada provides evidence that such bans can lead to significant increases in cigarette quitting. A menthol cigarette ban in the USA is projected to lead to even greater proportional impact in reducing smoking in the USA given its much higher proportion of menthol smokers, supporting the FDA's decision to move forward in the development of a rule

to ban menthol cigarettes. This pooled analysis of Canada's menthol cigarette ban also provides the foundation for estimating the impact of menthol bans on quitting among smokers in other countries and jurisdictions.

What this paper adds

What is already known on this topic

⇒ Between 2015 and 2018, all Canadian provinces banned menthol in cigarettes. Two pre–post studies found that the menthol ban—one in Ontario and one in seven Canadian provinces covering 83% of the Canadian population—led to increases in quit attempts and in quitting among menthol smokers compared with non-menthol smokers.

What this study adds

⇒ This study pools data from those two cohort studies—the International Tobacco Control Policy Evaluation (ITC) Project Canada Survey and the Ontario Menthol Ban Study—which used the similar definitions of menthol smoker and of quitting at post-ban, and were conducted within 1–2 months of each other at pre-ban (2016) and post-ban (2018) waves—to derive a more precise estimate of the impact of Canada's menthol ban on quitting and to apply this estimate to project the impact of a menthol ban in the USA, where prevalence of menthol cigarette use is high among cigarette smokers (40%), and much higher among African American smokers (85%).

⇒ The pooled analysis found that if a US menthol cigarette ban had the same impact as the Canadian menthol ban, there would be a substantial increase in quitting among menthol smokers in the USA, compared with non-menthol smokers, among all smokers and among African American smokers.

How this study might affect research, practice, or policy

⇒ This pooled analysis of Canada's menthol cigarette ban provides the foundation for estimating the impact of menthol bans in the USA and other countries.

Author affiliations

- ¹Department of Psychology, University of Waterloo, Waterloo, Ontario, Canada
²School of Public Health Systems, University of Waterloo, Waterloo, Ontario, Canada
³Ontario Institute for Cancer Research, Toronto, Ontario, Canada
⁴Department of Statistics & Actuarial Science, University of Waterloo, Waterloo, Ontario, Canada
⁵Psychiatry & Behavioral Sciences, Medical University of South Carolina, Charleston, South Carolina, USA
⁶Department of Health Behavior, Roswell Park Cancer Institute, Buffalo, New York, USA
⁷Lombardi Comprehensive Cancer Center, Georgetown University, Washington, District of Columbia, USA
⁸Rutgers Center for Tobacco Studies, Rutgers Biomedical and Health Sciences, New Brunswick, New Jersey, USA
⁹Psychology and Center for the Study of Tobacco Products, Virginia Commonwealth University, Richmond, Virginia, USA
¹⁰Department of Health Education and Promotion, East Carolina University, Greenville, North Carolina, USA
¹¹Ontario Tobacco Research Unit, University of Toronto, Dalla Lana School of Public Health, Toronto, Ontario, Canada
¹²Institute for Global Tobacco Control, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA
¹³Department of Public Health Sciences, University of Toronto, Toronto, Ontario, Canada

Twitter Michael O Chaiton @michaelchaiton

Contributors GTF is the guarantor: he accepts full responsibility for the work and/or the conduct of the study, had access to the data, and controlled the decision to publish. GTF led the conceptualisation of the study and prepared the initial draft.

MOC provided the Ontario Menthol Ban Study data, in coordination with GM, who conducted the analyses of the pooled data set. CDD and OG derived the estimates of numbers of menthol and non-menthol smokers from the 2019 NSDUH. All authors commented on drafts and approved the final version.

Funding The Canadian Arm of the ITC Four Country Smoking and Vaping Survey was supported by grants from the US National Cancer Institute (P01CA200512) and the Canadian Institutes of Health Research (FDN-148477). The Ontario Menthol Ban Study was supported by a grant from the US National Institute on Drug Abuse (R21DA04735801 and P50DA036105) and the Center for Tobacco Products of the US Food and Drug Administration; these grants also provided support for TE and EKS. Efforts for CDD and OG were provided by a Tobacco Centers of Regulatory Science award from the US National Cancer Institute and the US Food and Drug Administration (U54CA229973). Additional support to GTF was provided by a Senior Investigator Award from the Ontario Institute for Cancer Research and the O Harold Warwick Prize from the Canadian Cancer Society.

Disclaimer The content of this article is solely the responsibility of the authors and does not necessarily represent the views of any of the funding sources.

Competing interests GTF has served as a paid expert witness or consultant for governments defending their country's policies or regulations in litigation. He also served as a member of the Brazil Health Regulatory Agency (ANVISA) 2014 Working Group on Tobacco Additives. He has also served as a member of the Expert Group for Article 9 (Regulation of the contents of tobacco products) and Article 10 (Regulation of tobacco product disclosures) of the WHO Framework Convention on Tobacco Control. KMC has received payment as a consultant to Pfizer, for service on an external advisory panel to assess ways to improve smoking cessation delivery in healthcare settings. He has also served as a paid expert witness in litigation filed against cigarette manufacturers. TE is a paid consultant in litigation against the tobacco industry and also the electronic cigarette industry, and is named on a patent for a device that measures the puffing behaviour of electronic cigarette users and patent application for a smoking cessation intervention. TE and EKS are named on a patent application for a smartphone app that measures the characteristics of electronic cigarette devices and e-liquids.

Patient consent for publication Not required.

Ethics approval This study involves human participants and was approved by the University of Waterloo Office of Research Ethics (ORE#20803/30709) and University of Toronto Research Ethics Board. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Data are available upon reasonable request. In each country participating in the International Tobacco Control Policy Evaluation (ITC) Project, the data are jointly owned by the lead researcher(s) in that country and the ITC Project at the University of Waterloo. Data from the ITC Project are available to approved researchers 2 years after the date of issuance of cleaned data sets by the ITC Data Management Centre. Researchers interested in using ITC data are required to apply for approval by submitting an International Tobacco Control Data Repository (ITCDR) request application and subsequently to sign an ITCDR Data Usage Agreement. The criteria for data usage approval and the contents of the Data Usage Agreement are described online (<http://www.itcproject.org>).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Geoffrey T Fong <http://orcid.org/0000-0001-9098-6472>
 Janet Chung-Hall <http://orcid.org/0000-0002-9139-8100>
 Lorraine V Craig <http://orcid.org/0000-0001-5269-1817>
 K Michael Cummings <http://orcid.org/0000-0002-7103-7017>
 Richard J O'Connor <http://orcid.org/0000-0003-0644-182X>
 David T Levy <http://orcid.org/0000-0001-5280-3612>
 Thomas Eissenberg <http://orcid.org/0000-0002-8277-5437>
 Eric K Soule <http://orcid.org/0000-0003-2740-5633>

Robert Schwartz <http://orcid.org/0000-0001-7838-0769>
 Joanna E Cohen <http://orcid.org/0000-0002-3869-3637>
 Michael O Chaiton <http://orcid.org/0000-0002-9589-2122>

REFERENCES

- Chung-Hall J, Fong GT, Meng G. Evaluating the impact of menthol cigarette bans on cessation and smoking behaviors in Canada: longitudinal findings from the Canadian arm of the 2016-18 ITC Four Country Smoking and Vaping Surveys. *Tob Control* 2022;31:556–63.
- Chaiton MO, Nicolau I, Schwartz R, et al. Ban on menthol-flavoured tobacco products predicts cigarette cessation at 1 year: a population cohort study. *Tob Control* 2020;29:341–7.
- US Food and Drug Administration. FDA commits to evidence-based actions aimed at saving lives and preventing future generations of smokers. news release, 2021. Available: <https://www.fda.gov/news-events/press-announcements/fda-commits-evidence-based-actions-aimed-saving-lives-and-preventing-future-generations-smokers>
- US Federal Trade Commission. Federal Trade Commission cigarette report for 2020, 2021. Available: <https://www.ftc.gov/system/files/documents/reports/federal-trade-commission-cigarette-report-2020-smokeless-tobacco-report-2020/p114508fy20cigarettereport.pdf>
- Delnevo CD, Giovenco DP, Villanti AC. Assessment of menthol and Nonmenthol cigarette consumption in the US, 2000 to 2018. *JAMA Netw Open* 2020;3:e2013601.
- Canadian Tobacco, Alcohol and Drugs Survey (CTADS). Summary of results and detailed tables for 2017. Available: <https://www.canada.ca/en/health-canada/services/canadian-alcohol-drugs-survey/2017-summary.html>; <https://www.canada.ca/en/health-canada/services/canadian-alcohol-drugs-survey/2017-summary/2017-detailed-tables.html#t1>
- Hartman AM. What menthol smokers report they would do if menthol cigarettes were no longer sold. presentation to tobacco products scientific Advisory Committee, food and drug administration, 2011. Available: <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/TobaccoProductsScientificAdvisoryCommittee/UCM240176.pdf>
- O'Connor RJ, Bansal-Travers M, Carter LP, et al. What would menthol smokers do if menthol in cigarettes were banned? behavioral intentions and simulated demand. *Addiction* 2012;107:1330–8.
- Levy DT, Cadham CJ, Sanchez-Romero LM, et al. An expert elicitation on the effects of a ban on menthol cigarettes and Cigars in the United States. *Nicotine Tob Res* 2021;23:1911–20.
- Levy DT, Meza R, Yuan Z, et al. Public health impact of a US ban on menthol in cigarettes and cigars: a simulation study. *Tob Control* 2023;32:e37–44.
- Hill AB, Bradford Hill A. The environment and disease: association or causation? *Proc R Soc Med* 1965;58:295–300.
- Cochran WG, Chambers SP. The planning of observational studies of human populations. *J R Stat Soc Ser A* 1965;128:234–66.
- Villanti AC, Mowery PD, Delnevo CD, et al. Changes in the prevalence and correlates of menthol cigarette use in the USA, 2004-2014. *Tob Control* 2016;25:ii14–20.
- O'Connor RJ, Bansal-Travers M, Carter LP, et al. What would menthol smokers do if menthol in cigarettes were banned? behavioral intentions and simulated demand. *Addiction* 2012;107:1330–8.
- Non-Smokers' Rights Association. *Backgrounder on the Canadian tobacco market (Spring 2015) Technical report*, 2015.
- Schwartz R. Canada: controlling illicit tobacco trade. In: Dutta S, ed. *Confronting illicit tobacco trade: a global review of country experiences*. Washington, DC: World Bank, 2019: 51–71.
- National Research Council. *Understanding the U.S. Illicit Tobacco Market: Characteristics, Policy Context, and Lessons from International Experiences*. Washington, DC: The National Academies Press, 2015.
- Kurti M, von Lampe K, Johnson J. The intended and unintended consequences of a legal measure to cut the flow of illegal cigarettes into New York City: the case of the South Bronx. *Am J Public Health* 2015;105:750–6.
- Altria Client Services. Comment from Altria client services. *Fed Reg* 2018;83:12294.
- RAI Services Company. Comment from RAI services company. *Fed Reg* 2018;83:12294.
- Schroth KRJ, Villanti AC, Kurti M, et al. Why an FDA ban on menthol is likely to survive a tobacco industry lawsuit. *Public Health Rep* 2019;134:300–6.
- Stoklosa M. No surge in illicit cigarettes after implementation of menthol ban in nova Scotia. *Tob Control* 2019;28:702–4.
- Government of Canada. *Tobacco Reporting Regulations (SOR/2000-273)*, 2000. Available: <https://laws-lois.justice.gc.ca/eng/regulations/SOR-2000-273/>
- Canadian Cancer Society. Overview summary of federal/provincial/territorial tobacco control legislation in Canada, 2017. Available: http://convio.cancer.ca/documents/Legislative_Overview-Tobacco_Control-F-P-T-2017-final.pdf
- Government of Canada. Order amending the schedule to the tobacco act (menthol), 2017. Available: <https://canadagazette.gc.ca/rp-pr/p2/2017/2017-04-05/html/sor-dors45-eng.html>
- U.S. Food & Drug Administration. Family smoking prevention and tobacco control act, 2009. Available: <https://www.fda.gov/tobacco-products/rules-regulations-and-guidance/family-smoking-prevention-and-tobacco-control-act-table-contents>
- Mills SD, Henriksen L, Golden SD, et al. Disparities in retail marketing for menthol cigarettes in the United States, 2015. *Health Place* 2018;53:62–70.

Supplemental Tables for Fong et al. The Impact of Canada's Menthol Cigarette Ban on Quitting Among Menthol Smokers: Pooled Analysis of Pre-Post Evaluation from the ITC Project and the Ontario Menthol Ban Study and Projections of Impact in the United States

Supplemental Table 1. Comparison of ITC Canada Survey and Ontario Menthol Ban Survey

Supplemental Table 2. The association of post-ban smoking cessation and pre-ban menthol smoking status for all smokers and for daily smokers, and interactions with ethnicity (Black vs. Non-Black) for all smokers

Supplemental Table 1. Comparison of ITC Canada Survey and Ontario Menthol Ban Survey

	ITC Canada Survey	Ontario Menthol Ban Study
Sample	Nationally representative sample of 1236 adult (18+) smokers; residents of 7 provinces (Quebec, Ontario, Prince Edward Island, Newfoundland and Labrador, British Columbia, Saskatchewan, and Manitoba)	Provincially representative sample of 1084 adult (18+) smokers + additional smokers from a dedicated Ontario Smokers' Panel; residents of Ontario
Sampling frame	Random selection of respondents from a high-quality national web panel (Leger), using a combination of probability-based, or non-probability opt-in sampling frames, or a combination of these methods.	Respondents recruited using a combination of random digit dialing and convenience sample.
Survey dates	Pre-ban: July 7, 2016 to November 22, 2016 Post-ban: February 23, 2018 to July 9, 2018	Pre-ban: September to December 2016 Post-ban: January to August 2018
Menthol smoking status	<p>Pre- and post-ban: Respondents who said they had a regular brand of cigarettes were asked: 'How would you describe the flavour of your usual cigarette brand?' (response options: just tobacco, tobacco and menthol, tobacco and some other flavour, refused, don't know). Respondents were also asked if their last purchased brand is just tobacco, tobacco and menthol, or tobacco and some other flavour.</p> <p>Respondents who selected 'tobacco and menthol' flavour as their regular brand were categorised as 'mainly menthol smokers' Those who do not have a regular brand or their regular brand is not menthol but bought menthol cigarettes in their last purchase were categorised as 'occasional menthol smokers'. Those who selected 'just tobacco' or 'tobacco and some other flavour' and those who did not buy menthol cigarettes in their last purchase were categorised as 'non-menthol smokers'.</p>	<p>Pre-ban: self-reported past year use of menthol cigarettes (response options: every day, occasionally, on rare occasions, not at all). Daily smokers who used menthol cigarettes in the past year and occasional smokers who reported occasional use of menthol cigarettes in the past year were categorised as 'mainly menthol smokers'. Daily smokers who reported occasional use of menthol cigarettes in the past year were categorized as 'occasional menthol smokers'. Those who had not used menthol cigarettes in past year were categorised as 'non-menthol smokers'.</p> <p>Post-ban: self-reported current use of menthol cigarettes. Respondents who reported current use of menthol cigarettes were categorised as 'menthol smokers', and those who were not currently using menthol cigarettes were categorised as 'non-menthol smokers'.</p>
Post-ban quit success	Respondents who reported that they were not currently smoking cigarettes at post-ban.	Respondents who reported that they were not currently smoking cigarettes at post-ban.

Supplemental Table 2. The association of post-ban smoking cessation (1= current quitter; 0 = current smoker) and pre-ban menthol smoking status for all smokers and for daily smokers, including interactions with ethnicity (Black vs. Non-Black) for all smokers

Explanatory variables	Pre-ban All Smokers					Pre-ban Daily Smokers		
	Main effects			Interaction with ethnicity (Black vs. Non-Black)		N	OR (95% CI)	P
	N	OR (95% CI)	P	OR (95% CI)	P			
Pre-ban menthol smoking status								
Mainly menthol	423	1.64 (1.17 - 2.30)	0.004	21.13 (2.15 - 208.0)	0.009	308	1.90 (1.30 - 2.79)	<0.001
Occasional menthol	407	1.24 (0.84 - 1.82)	0.271	4.96 (0.47 - 52.25)	0.182	407	1.35 (0.90 - 2.02)	0.144
Non-menthol	1490	Reference		Reference		1285	Reference	
Pre-ban smoking status								
Daily	2000	0.48 (0.33 - 0.68)	<0.001	0.48 (0.34 - 0.69)	<0.001	-		
Non-daily	320	Reference		Reference		-		
Sex								
Female	1262	0.87 (0.68 - 1.10)	0.241	0.87 (0.68 - 1.11)	0.263	1089	0.76 (0.58 - 1.00)	0.046
Male	1058	Reference		Reference		911	Reference	
Age group								
18-24	247	0.79 (0.52 - 1.19)	0.262	0.79 (0.52 - 1.19)	0.257	160	0.52 (0.29 - 0.92)	0.024
25-39	510	1.20 (0.86 - 1.69)	0.287	1.20 (0.86 - 1.69)	0.287	425	1.04 (0.72 - 1.52)	0.822
40-54	754	0.86 (0.63 - 1.17)	0.332	0.86 (0.63 - 1.17)	0.327	673	0.74 (0.53 - 1.03)	0.075
55 and up	809	Reference		Reference		742	Reference	
Education								
Low	683	Reference		Reference		624	Reference	
Medium	1037	1.43 (1.06 - 1.95)	0.021	1.44 (1.06 - 1.96)	0.019	910	1.60 (1.14 - 2.24)	0.006
High	582	1.51 (1.08 - 2.12)	0.017	1.53 (1.09 - 2.14)	0.015	452	1.45 (0.98 - 2.13)	0.063
Not Stated	18	0.97 (0.26 - 3.68)	0.966	0.98 (0.26 - 3.72)	0.978	14	0.57 (0.07 - 4.51)	0.593
Pre-ban cigarettes smoked per day								
1-10 cigarettes	960	1.10 (0.71 - 1.71)	0.670	1.10 (0.71 - 1.70)	0.684	660	1.22 (0.76 - 1.96)	0.407
11-20 cigarettes	895	0.77 (0.50 - 1.20)	0.252	0.77 (0.50 - 1.20)	0.249	885	0.87 (0.55 - 1.37)	0.537
21-30 cigarettes	317	0.34 (0.19 - 0.61)	<0.001	0.34 (0.19 - 0.61)	<0.001	316	0.36 (0.20 - 0.67)	0.001
31+ cigarettes	148	Reference		Reference		139	Reference	
Study region								
ITC-BC	190	0.65 (0.36 - 1.16)	0.143	0.64 (0.36 - 1.15)	0.137	161	0.77 (0.41 - 1.44)	0.413
ITC-MB/SK/NL/PEI	143	0.71 (0.38 - 1.34)	0.290	0.71 (0.38 - 1.33)	0.287	124	0.66 (0.31 - 1.38)	0.266
ITC-ON	572	0.58 (0.40 - 0.85)	0.005	0.59 (0.40 - 0.86)	0.006	461	0.57 (0.37 - 0.88)	0.011
ITC-QC	331	0.75 (0.48 - 1.15)	0.182	0.75 (0.49 - 1.15)	0.186	277	0.78 (0.48 - 1.27)	0.313
Ontario Menthol Ban Study-ON	1084	Reference		Reference		977	Reference	

Explanatory variables	Pre-ban All Smokers					Pre-ban Daily Smokers		
	Main effects			Interaction with ethnicity (Black vs. Non-Black)		N	OR (95% CI)	P
	N	OR (95% CI)	P	OR (95% CI)	P			
Ethnicity								
Black	30	0.76 (0.27 – 2.13)	0.593	0.24 (0.05 – 1.28)	0.094	25	0.74 (0.20 – 2.70)	0.648
Non-Black	2290	Reference		Reference		1975	Reference	
Pre-ban menthol smoking status * ethnicity								
Mainly menthol, Black	4			13.08 (1.31 - 131.1)	0.029			
Mainly menthol, Non-Black	419			Reference				
Occasional menthol, Black	9			4.05 (0.38 - 42.68)	0.245			
Occasional menthol, Non-Black	398			Reference				
Non-menthol, Black	17			Reference				
Non-menthol, Non-Black	1473			Reference				

Note. The interactions with ethnicity (Black vs. non-Black) were not presented for daily smokers due to insufficient sample size.