

**Table A5** Pearson correlation coefficients between dependent and predictor variables in the model for loss<sup>1</sup>

Predictor variables	Dependent variable and predictor variables <sup>2</sup>									
	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29
V21	0.14									
V22	0.14	-0.02								
V23	-0.08	0.07	-0.06							
V24	0.09	0.16	-0.12	-0.19						
V25	0.16	0.02	-0.21	0.07	0.27					
V26	-0.23	0.07	0.18	0.06	0.13	-0.06				
V27	0.27	0.05	-0.01	0.32	-0.16	-0.16	0.10			
V28	-0.32	0.21	0.09	-0.09	0.19	-0.02	0.14	-0.10		
V29	-0.32	0.06	0.24	0.05	0.07	0.00	0.34	0.07	0.24	
V30	-0.09	0.27	0.02	0.07	0.11	-0.17	0.08	0.40	0.06	0.16

Notes

1. Sample size: 74

2. Variables were labelled as follows:

**Dependent variable**

V20 log(Loss)

**Predictor variables**

V21 Index of multiple deprivation

V22 Co-ordinator has responsibilities other than developing/running service

V23 Salary level of co-ordinator

V24 Difficulty finding good quality training courses for advisors or counsellors

V25 Central base used (almost) exclusively by smoking cessation service

V26 Satellite/outreach bases shared between service and other organisations

V27 Percentage of users receiving individual support

V28 Length (minutes) of individual sessions

V29 No. of different types of professionals receiving training from service staff

V30 Whether advisors/counsellors ever recommended bupropion